



HOKKAIDO UNIVERSITY

Title	CORRELATION OF FEATHER PULP LESIONS IN CHICKENS INOCULATED WITH HERPESVIRUS OF TURKEYS WITH RESPONSE OF MAREK'S DISEASE
Author(s)	OKADA, Hirokazu
Citation	Japanese Journal of Veterinary Research, 34(2), 151-151
Issue Date	1986-04-30
Doc URL	https://hdl.handle.net/2115/2998
Type	departmental bulletin paper
File Information	KJ00002374405.pdf



CORRELATION OF FEATHER PULP LESIONS IN CHICKENS INOCULATED WITH HERPESVIRUS OF TURKEYS WITH RESPONSE OF MAREK'S DISEASE

Hirokazu OKADA

*Department of Comparative Pathology
Faculty of Veterinary Medicine
Hokkaido University, Sapporo 060, Japan*

In order to clarify the correlation of feather pulp lesions (FPL) in chicken vaccinated with herpesvirus of turkeys (HVT) with Marek's disease (MD) response, the following four groups were compared. Group A: inoculation with HVT at hatching; group B: treatment of 3-day embryo with Testosterone propionate (TP) and inoculation with HVT at hatching; group C: inoculation of 18-day embryo with HVVT; group D: non-inoculation.

There were two kinds of FPL in the HVT-inoculated chickens 2-6 weeks of age: L lesion=perivascular small lymphocyte infiltration, which was frequently seen in 2-6-week-old chickens; P lesion=plasma cellular infiltration, which was occasionally seen in 4-6-week-old birds. Occurrence of L lesion was also noted in HVT-uninoculated chickens but was significantly higher in the HVT-inoculated groups A and B. On the contrary, incidence of L lesion in group C birds (inoculated with HVT at embryo stage) was significantly low.

The chickens were challenged with MD virus at 3 or 6 weeks of age. Fifty-five chickens were affected with MD by 15 or 18 weeks of age. MD-affected birds were classified into three groups based on necropsy findings: lymphoma-positive, 35 birds; nerve lesion-positive, 3 birds; so-called "early-mortality syndrome", 17 birds. Interestingly, incidence of L (and P) lesion, which was seen prior to challenge with MD virus, was related closely with MD response, that is, groups C and D with relatively low incidence of L lesion showed significantly higher mortality from MD than groups A and B with higher incidence of L lesion. Individually, L lesion-negative chickens (without evidence of L lesion up to 3 or 6 weeks of age) showed a higher incidence of MD-associated nuclear-inclusion formation in the feather-follicle epithelium, higher mortality and higher incidence of "early-mortality syndrome" than L lesion-positive chickens. L lesion-positive birds were relatively resistant to MD, but almost all of the MD-affected chickens had lymphoma in the viscera.