



HOKKAIDO UNIVERSITY

Title	Bacterial Genes Responsible for the Biosynthesis of Eicosapentaenoic and Docosahexaenoic Acids and Their Heterologous Expression
Author(s)	Okuyama, Hidetoshi; Orikasa, Yoshitake; Nishida, Takanori et al.
Citation	Applied and Environmental Microbiology, 73(3), 665-670 https://doi.org/10.1128/AEM.02270-06
Issue Date	2007-02
Doc URL	https://hdl.handle.net/2115/30278
Rights	Copyright © American Society for Microbiology
Type	journal article
File Information	AEM73-3.pdf



MINIREVIEW

Bacterial Genes Responsible for the Biosynthesis of Eicosapentaenoic and Docosahexaenoic Acids and Their Heterologous Expression

Hidetoshi Okuyama,^{1*} Yoshitake Orikasa,¹ Takanori Nishida,¹ Kazuo Watanabe,² and
Naoki Morita³

*Graduate School of Environmental Earth Science, Hokkaido University, Sapporo 060-0810,
Japan,¹ Sagami Chemical Research Center, Hayakawa, Ayase 252-1193, Japan,²
and Research Institute of Genome-based Biofactory, National Institute of Advanced
Industrial Science and Technology (AIST), Sapporo 062-8517, Japan³*

Running title: Bacterial genes required for EPA and DHA biosynthesis

*Corresponding author. Mailing address: *Graduate School of Environmental Earth
Science, Hokkaido University, Sapporo 060-0810, Japan.* Phone: +81-11-706-4523. Fax:
+81-11-706-2347. E-mail: hoku@ees.hokudai.ac.jp.

Until the 1990s, it was thought that, with the exception of cyanobacteria, bacteria had no polyunsaturated fatty acids (PUFAs). This was probably because the bacterial species whose physiology, biochemistry and molecular biology had been well studied until that time were mesophilic species, such as *Escherichia coli*, which have no PUFAs. It has since been found that $n-3$ long chain PUFAs such as eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) are preferentially distributed in psychrophilic bacteria that inhabit relatively unusual environments including low temperature, deep-sea environments and the intestines of sea fish (25, 31, 32). Although several PUFA-producing bacterial strains had been reported until the 1970s (15), the discovery of DHA and EPA in bacteria from deep-sea water and sediments and detailed analysis of them by DeLong and Yayanos in 1986 (2) resulted in the initiation of research into PUFA-producing bacteria.

It is interesting that the PUFAs detected in those bacteria were mostly EPA or DHA, and not C18 PUFAs such as linoleic and linolenic acids, which are most common in animals, plants, fungi and cyanobacteria. However, although EPA- and DHA-producing bacteria were discovered in the 1980s, they have not been given much attention. This can easily be imagined as microorganisms (including eukaryotes such as microalgae) with PUFAs such as EPA and DHA are widespread in marine environments (24, 25, 28). It was also thought that bacterial, as well as eukaryotic, EPA and DHA were biosynthesized by a combination of elongation and oxygen-dependent desaturation of existing fatty acids. Therefore, the successful cloning in 1996 of genes involved in the biosynthesis of EPA from *Shewanella* sp. SCRC-2738 isolated from marine fish intestines (33) was very important and could be

regarded as the first stage of research into such genes. This EPA-producing strain is presently identified as *Shewanella pneumatophori* SCRC-2738 (4). It was very surprising that the deduced proteins encoded by the EPA biosynthesis genes (*pfa* genes, see below) included no conserved sequences from fatty acid desaturase genes (7, 33), although data on fatty acid desaturases was very limited in the early 1990s (11). Five deduced proteins involved in EPA biosynthesis had domains that were conserved in fatty acid synthetase and/or polyketide (PK) synthetases (PKSs) (7). In addition, evidence has been presented to show that bacterial PUFAs can be synthesized under anaerobic conditions (7).

After the discovery of the EPA biosynthetic *pfa* genes, their homologues were cloned from various marine bacteria. The *pfa* genes of *Moritella marina* MP-1 were the first genes to be cloned from DHA-producing bacteria (27). Recent genome sequencing of various organisms, including bacteria, demonstrates that *pfa* genes are distributed abundantly in members of the bacterial genera *Shewanella* (29) and *Colwellia* (6), and that these strains have mainly been isolated from marine sources. There are some variations in the structure of *pfa* gene clusters, although they have a basic structure that is common between all types of clusters (see below). There is no doubt that all bacterial EPA and DHA are synthesized by the PKS system (7, 16).

Although the recombinant production of EPA in *E. coli* was achieved in 1996 using *pfa* genes from *S. pneumatophori* SCRC-2738 (33), no reports on the heterologous synthesis of DHA were found in the literature. This was because the *pfa* gene cluster in *M. marina* MP-1 lacked a gene corresponding to the *pfaE* gene of *S. pneumatophori* SCRC-2738. Recently, however, the *pfaE* gene was cloned from *M. marina* MP-1 (17), and the

recombinant DHA was produced (19). This might be called the second stage of the research. Here, we review the bacterial genes responsible for the biosynthesis of EPA and DHA, covering such aspects as gene cloning, characterization of the structure of the genes and their domain structures, and the recombinant production of EPA and DHA.

GENES RESPONSIBLE FOR THE BIOSYNTHESIS OF EPA AND DHA

The genes responsible for the biosynthesis of EPA were first cloned as a cosmid carrying a DNA fragment of approximately 38 kbp (33). This fragment carried at least 18 open reading frames (ORFs), of which a cluster of only five was necessary for the biosynthesis of EPA. These ORFs are now named *pfaA*, *pfaB*, *pfaC*, *pfaD* and *pfaE* (13, 17 and see Fig. 1). The clustered genes were designated the 'EPA biosynthesis gene cluster'. When various *E. coli* strains were transformed with the EPA biosynthesis gene cluster, they normally produced EPA at 1%–5% of total fatty acids (20, 33).

Genes homologous to the *pfa* genes have been cloned from various EPA- and DHA-producing bacteria. Allen and Bartlett (1) reported finding a cluster of EPA biosynthesis genes in *Photobacterium profundum* SS9. However, this cluster did not include *pfaE*, which encodes phosphopantetheinyl transferase (PPTase). A partial sequence corresponding to *pfaA* was cloned from EPA-producing *Shewanella* sp. GA-22 (3). Clustered genes homologous with those from *S. pneumatophori* SCRC-2738 were cloned from the DHA-producing deep-sea bacterial species *M. marina* MP-1 (27). This gene cluster also included only *pfaA–D* and lacked *pfaE*. Therefore, attempts at the recombinant production of DHA in *E. coli* or in other host organisms were unsuccessful until quite

recently. The *pfa* genes from *M. marina* MP-1 constitute the only DHA biosynthesis gene cluster currently cloned.

Genome sequencing of various bacteria either known or expected to produce EPA or DHA demonstrates the abundance and wide distribution of such genes. Interestingly, bacteria that have genes homologous with *pfa* genes are mostly found in marine sources. In the genome of EPA-producing *Shewanella oneidensis* MR-1 (29), a cluster of *pfaA–D* genes were found. *Colwellia psychrerythraea* 34H (6), which is expected (but not confirmed) to produce DHA, has a similar cluster of genes. All of the aforementioned *pfa* gene clusters contained *pfaE* as an ORF encoding a single protein either within the cluster or outside the cluster. The former and latter structures of *pfa* gene clusters are designated type I and type II *pfa* genes, respectively, in this review (Fig. 1). However, the structure of the type I *pfa* genes is unlikely to be essentially different from that of the type II *pfa* genes. This is known because even the gene cluster from *S. pneumatophori* SCRC-2738, which contains all five genes, has the sequential arrangement of *pfa* genes broken by the presence of two ORFs unrelated to EPA biosynthesis (Fig. 1). Unlike these structures, *Pseudoalteromonas* sp. DS-12 has a unique *pfa* gene cluster consisting of four ORFs (Dai and Zhang, unpublished; see accession no. ABF00130). In this strain, the PfaE is integrated in PfaC/E, and *pfaB* and *pfaC/E* genes are separated by one unrelated ORF (Fig. 1). This is designated the type III *pfa* gene. Gene sequences homologous with those of the *pfa* genes have also been found in eukaryotic marine microalgae producing EPA and DHA (7). Table 1 summarizes known bacterial genes homologous with the EPA biosynthesis genes *pfaA–E* of *S. pneumatophori* SCRC-2738.

STRUCTURE OF INDIVIDUAL *PFA* GENES

It is evident that only five *pfaA–E* genes are generally necessary for the biosynthesis of EPA and DHA. Although the basic structure of all *pfa* genes for EPA and DHA biosynthesis is very similar, the domain structures of some of the individual genes are slightly different (Fig. 1, Table 1). In the type I and II *pfa* gene clusters, *pfaA* is tentatively thought to encode a multifunction protein that includes domains for 3-ketoacyl synthase (KS), malonyl-CoA:acyl carrier protein (ACP) acyltransferase (AT), normally five or six ACP repeats and 3-ketoacyl-ACP reductase (KR). The *pfaC* gene encodes a protein with two KS repeats and two or three 3-hydroxydecanoyl-ACP dehydratases (HD). The second KS domain in the PfaC of *S. pneumatophori* SCRC-2738, *P. profundum* SS9, and *M. marina* MP-1 was considered to be a chain length factor (1, 17, 19). In this study, however, both the domains remained undifferentiated as information about their precise function was not available. Previously, the number of HD domains in the PfaC gene was recognized to be two (1, 7, 17); however, a very recent database search demonstrated that PfaC includes three sequential domains of HD based on *S. pneumatophori* SCRC-2738, *Shewanella marinintestina* IK-1, and *M. marina* MP-1 only (Fig. 1, Table 1). Interestingly, in the PfaC for EPA, all three HD domains were homologous with FabA, with a high degree of similarity, while the PfaC for DHA had two HD domains similar to FabA and one domain, in the center, similar to FabZ/FabA (Table 1). Genes of *pfaB* and *pfaD* encode proteins with KR and AT domains and an enoyl reductase (ER) domain, respectively (7, 20). A KS domain is included only in the PfaB gene for DHA derived from *M. marina* MP-1 (27) and *C. psychrerythraea* 34H (6). However, the active site of the KS domain of PfaB in *M.*

marina MP-1 lacked a cysteine residue, which is different from the same gene in *S. pneumatophori* SCRC-2738 and *P. profundum* SS9 (1). The domain structure of the *pfa* genes of *Pseudoalteromonas* sp. DS-12 was quite different from that of the same genes in the other bacterial strains. The PfaB gene was found to have two KS domains, and the PfaC/E gene (which was registered as a product of the *pfaD* gene in the database; Dai and Zhang, unpublished) has one PPTase domain and two HD domains (Fig. 1). At the downstream end of the *pfaC/E* gene was a 483 bp remnant DNA region (this was regarded as the *pfaD* gene in this review), which included a partial sequence similar to that of ER.

Compared with the PfaA–D genes, the domain structures of PfaE are well characterized. The PfaE (PPTase) gene for the biosynthesis of EPA or DHA can be regarded as a member of a large gene family of Sfp-type PPTases based on the sizes of the deduced protein molecules and their domain structure (17). Based on their domain structures, Orikasa et al. (17, 18) divided all Sfp-type PPTases into two groups: PPTases responsible mainly for the biosynthesis of EPA or DHA (group I), characterized by P0, P1a and P1b domains and those responsible mainly for the synthesis of PKs; and nonribosomal peptides (NRPs) (group II), characterized by the domains 1A, P1a' and P1b', which correspond to the P0, P1a and P1b domains of group I (18). Although the P2 and P3 domains are commonly conserved in the two groups, there is a higher degree of similarity in these domains within groups than between groups (18). The *pfaE* gene complementing the *pfaA–D* genes of *P. profundum* SS9 has not been cloned. However, genome sequencing of this bacterium (30) provided a candidate sequence (accession no. CAG23685) from the group II Sfp-type

PPTase (PfaE). On the other hand, the Sfp-type PPTase of *Bacillus subtilis*, Sfp itself, which is involved in the biosynthesis of surfactin (a NRP; 12), belongs to group I (18).

FUNCTIONAL COMPATIBILITY OF INDIVIDUAL PFA GENES

The compatibility of *pfa* genes involved in the biosynthesis of PUFAs has been investigated mostly using the *pfa* genes from the EPA-producing *S. pneumatophori* SCRC-2738 and DHA-producing *M. marina* MP-1. The *pfaE* gene (pETSTV::*pfaE*) from *M. marina* MP-1 complemented the pDHA3 vector carrying *pfaA–D* genes from DHA-producing *M. marina* MP-1 (19), as well as the pEPA Δ 1,2,3 vector carrying *pfaA–D* genes from EPA-producing *S. pneumatophori* SCRC-2738 (17). To examine the compatibility of *pfaE* from EPA-producing *S. pneumatophori* SCRC-2738 with pDHA3 from *M. marina* MP-1, an *E. coli* DH5 α transformant was produced that utilized pEPA Δ 5 carrying *pfaB–E* genes from *S. pneumatophori* SCRC-2738 and pDHA3 (18). Both EPA and DHA were produced in this combination, suggesting that the *pfaE* gene in pEPA Δ 5 is involved in producing DHA (18). That is, the PPTase (PfaE) from *S. pneumatophori* SCRC-2738 was able to recognize the ACP repeats (substrate) integrated into the *pfaA* gene product of *M. marina* MP-1, and the *pfaA* product of the DHA biosynthesis gene cluster played a role in EPA biosynthesis. Orikasa et al. (18, 19) obtained pDHA2 in the course of preparing pDHA3. Although pDHA2 carried *pfaA–D* genes from *M. marina* MP-1, the combined expression of pDHA2 and *pfaE* produced neither EPA nor DHA, as the *pfaA* in pDHA2 had one fatal base pair replacement inactivating the gene cluster. However, when pDHA2 was coexpressed with any of the three types of deletion clone of the pEPA clusters

from *S. pneumatophori* SCRC-2738 (that is, the clones carrying *pfaA*, *pfaC*, *pfaD* and *pfaE*; *pfaA*, *pfaB*, *pfaD* and *pfaE*; or *pfaA*, *pfaB*, *pfaC* and *pfaE*) in *E. coli*, all transformants produced both EPA and DHA. In contrast, neither EPA nor DHA was produced by a combination of pDHA2 and a deletion clone carrying *pfaB*, *pfaC*, *pfaD* and *pfaE* (Orikasa, Y., Yamada, A., Yu, A., Watanabe, K. and Okuyama, H., unpublished). All these results suggest that not only *pfaE* and *pfaA*, but also *pfaB*, *pfaC* and *pfaD*, are functionally compatible in the biosynthesis of EPA and DHA. The recombinant production of EPA and trace levels of DHA in *E. coli* was confirmed by the use of *pfa* genes (pIK814) of *S. marinintestina* IK-1 and by a combination of these genes with pDHA2 from *M. marina* MP-1 (N. Morita, Y. Yano, S. Ohgiya, and H. Okuyama, unpublished).

According to Allen and Bartlett (1), the *pfaA–D* gene cluster in *P. profundum* SS9 did not complement PPTase genes from *Shewanella* sp. SC2 and *B. subtilis*, both of which are classed in group I, in the production of EPA. No sequences similar to those PPTases, other than the deduced sequence of CAG23685, have been found in the genome (i.e., in either chromosomes 1 or 2) of *P. profundum* SS9 (30). It would be interesting to examine the compatibility of the PPTase gene from *P. profundum* SS9 with the *pfaA–D* genes from this strain and those from other EPA- or DHA-producing bacteria such as *S. pneumatophori* SCRC-2738 and *M. marina* MP-1. The functional compatibility of each *pfa* gene is summarized in Table 2.

FUTURE PERSPECTIVES

The EPA and DHA biosynthesis gene clusters were initially cloned with the aim of expressing them in various host organisms such as cyanobacteria, yeast and plants (27, 33) for the purpose of producing commercially important materials. However, only low levels of EPA were produced by the recombination of the genes in cyanobacteria (26, 34). The recombinant production of DHA in *E. coli* has been reported quite recently (18). Although fish oils are the most important source of EPA and DHA, the contamination of fish due to pollution, as well as unstable fish catches, have created a need for alternative ways to provide those PUFAs (21). The use of the PKS system to produce EPA or DHA in heterologous host organisms has some benefits, such as the need for lesser amounts of reducing equivalents such as NADPH (19, 23) and the simplicity the PUFAs have in consisting solely of EPA or DHA. Since bacterial PKS systems involved in the production of EPA and DHA are generally less active at moderate temperatures (19, 20), their genetic modification and selection in host organisms should be considered.

The PKS systems would provide a useful tool for investigating the physiological roles of EPA and DHA and their biosynthetic mechanisms. EPA levels in recombinants can be changed at random using various vectors carrying *pfa* gene(s) from *S. pneumatophori* SCRC-2738 (20). The antioxidative function of EPA was first observed using such a recombinant in *E. coli* (13, 15). Similar systems could be constructed for DHA by isolating individual *pfa* genes involved in DHA production. A biosynthetic mechanism has been proposed for EPA (and probably DHA) production, which is similar to the anaerobic pathway of unsaturated fatty acid biosynthesis (7). However, no direct evidence is available

to support this proposed mechanism. To detect intermediates in the biosynthesis of EPA or DHA, PKS recombinant systems could be used. The heterologous production of DHA in *E. coli* that had been transformed with *pfa* genes from the marine piezophilic psychrophile *M. marina* MP-1 was more active at lowered growth temperatures (19). The finding coincided with that this bacterium is psychrophilic and that it inherently formed more DHA at low temperatures (2). However, the effects of salinity of culture media and hydrostatic conditions on the recombinant production of PUFAs have not been elucidated. Transcriptional regulation of *pfaA–D* genes has been studied targeting the *pfaA–D* genes of *P. profundum* SS9 (1). More information would be produced using the recombinant systems carrying variously combinations of *pfa* genes. Normal concentrations (10–100 μ M) of cerulenin, an inhibitor of *de novo* biosynthesis of fatty acids, enhanced the synthesis of EPA and DHA in *P. profundum*, *S. marinintestina* IK-1, and *M. marina* MP-1 (1, 8). It has been demonstrated that *M. marina* MP-1 has a fatty acid biosynthetic (*fab*) gene cluster that takes part in the *de novo* synthesis of fatty acids with moderate chain lengths up to C18 (9, 10). The relationship between the PKS system and *de novo* fatty acid biosynthesis could be investigated using the PKS recombinant systems.

Menzella et al. (5) proposed combinatorial PK biosynthesis by design and rearrangement of modular PK synthase (PKS) genes. Type I PKS genes of 3–6 kbp length are similar to the *pfa* genes. Some *pfa* genes such as *pfaA* and *pfaC* that are structurally similar to PKS genes might be used for the production of novel and commercially beneficial polyketides. If five or six repeats of ACP domains operate as a cluster to enhance the biosynthesis of PUFAs, as is the case in the biosynthesis of the PK antibiotic mupirocin in *Pseudomonas*

fluorescens, in which two tandem repeats of carrier proteins are involved (22), the corresponding DNA region would become a useful tool for the enhanced production of various types of polyketides.

We were unable to identify the factor(s) that determine the final product in PKS systems that produce PUFAs. It is speculated that some cooperative interactions between domains of different Pfa proteins, rather than the activity of any single Pfa protein, might be involved in directing the final product in the system. To find an answer to this question would be the third stage in this research.

ACKNOWLEDGEMENT

All *pfa* genes from *Shewanella pneumatophori* SCRC-2738 used in this study are provided by Sagami Chemical Research Center, Japan

REFERENCES

1. **Allen, E.E., and D. H. Bartlett.** 2002. Structure and regulation of the omega-3 polyunsaturated fatty acid synthase genes from the deep-sea bacterium *Photobacterium profundum* strain SS9. Microbiology **148**:1903–1913.
2. **DeLong, E. F., and A. A. Yayanos.** 1986. Biochemical function and ecological significance of novel bacterial lipids in deep-sea procaryotes. Appl. Environ. Microbiol. **51**:730–737.

3. **Gentile, G., V. Bonasera, C. Amico, L. Giuliano, and M. M. Yakimov.** 2003. *Shewanella* sp. GA-22, a psychrophilic hydrocarbonoclastic antarctic bacterium producing polyunsaturated fatty acids. *J. Appl. Microbiol.* **95**:1124–1133.
4. **Hirota, K., Y. Nodasaka, Y. Orikasa, H. Okuyama, and I. Yumoto.** 2005. *Shewanella pneumatophori* sp. nov., eicosapentanoic-acid-producing marine bacterium isolated from pacific mackerel (*Pneumatophorus japonicus*) intestine. *Int. J. Syst. Evol. Microbiol.* **55**:2355–2359.
5. **Menzella, H.G., R. Reid, J. R. Carney, S. S. Chandran, S. J. Reisinger, K. G. Patel, D. A. Hopwood, and D. V. Santi.** 2005. Combinatorial polyketide biosynthesis by de novo design and rearrangement of modular polyketide synthase genes. *Nat. Biotechnol.* **23**:1171–1176.
6. **Méthé, B.A., K. F. Nelson, J. W. Deming, B. Momen, E. Melamud, X. Zhang, J. Moulton, R. Madupu, W. C. Nelson, R. J. Dodson, L. M. Brinkac, S. C. Daugherty, A. S. Durkin, R. T. DeBoy, J. F. Kolonay, S. A. Sullivan, L. Zhou, T. M. Davidsen, M. Wu, A. L. Huston, M. Lewis, B. Weaver, J. F. Weidman, H. Khouri, T. R. Utterback, T. V. Feldblyum, and C. M. Fraser.** 2005. The psychrophilic lifestyle as revealed by the genome sequence of *Colwellia psychrerythraea* 34H through genomic and proteomic analyses. *Proc. Natl. Acad. Sci. USA.* **102**:10913–10918.
7. **Metz, J.G., P. Roessler, D. Facciotti, C. Levering, F. Dittrich, M. Lassner, R. Valentine, K. Lardizabal, F. Domergue, A. Yamada, K. Yazawa, V. Knauf, and J. Browse.** 2001. Production of polyunsaturated fatty acids by polyketide synthases in both prokaryotes and eukaryotes. *Science* **293**:290–293.

8. **Morita, N., T. Nishida, M. Tanaka, Y. Yano, and H. Okuyama.** 2005. Enhancement of polyunsaturated fatty acid production by cerulenin treatment in polyunsaturated fatty acid-producing bacteria. *Biotechnol. Lett.* **27**:389–393.
9. **Morita, N., M. Tanaka, and H. Okuyama.** 2000. Biosynthesis of fatty acids in the docosahexaenoic acid-producing bacterium *Moritella marina* strain MP-1. *Biochem. Soc. Trans.* **28**:943–945.
10. **Morita, N., A. Ueno, M. Tanaka, S. Ohgiya, H. Hoshino, K. Kawasaki, I. Yumoto, K. Ishizaki, and H. Okuyama.** 1999. Cloning and sequencing of clustered genes involved in fatty acid biosynthesis from the docosahexaenoic acid-producing bacterium, *Vibrio marinus* strain MP-1. *Biotechnol. Lett.* **21**:641–646.
11. **Murata, N., and H. Wada.** 1995. Acyl-lipid desaturases and their importance in the tolerance and acclimatization to cold of cyanobacteria. *Biochem. J.* **308**:1–8.
12. **Nakano, M. M., N. Corbell, J. Besson, and P. Zuber.** 1992. Isolation and characterization of *sfp*: a gene that functions in the production of the lipopeptide biosurfactant, surfactin, in *Bacillus subtilis*. *Mol. Gen. Genet.* **232**:313–321.
13. **Nishida, T., Y. Orikasa, Y. Ito, R. Yu, A. Yamada, K. Watanabe, and H. Okuyama.** 2006. *Escherichia coli* engineered to produce eicosapentaenoic acid becomes resistant against oxidative damages. *FEBS Lett.* **580**:2731–2735.
14. **Nishida, T., Y. Orikasa, K. Watanabe, N. Morita, and H. Okuyama.** 2006. Evaluation of the antioxidative effects of eicosapentaenoic acid by growth inhibition testing on plates using *Escherichia coli* transformed with *pfa* genes. In C. Benning and J. Ohlrogge (ed.), *Advances in Plant Lipid Research: Proceedings of the 17th*

International Symposium on Plant Lipids, East Lansing, Michigan, July 2006 ,
Michigan State University Press, East Lansing, Michigan. in press.

15. **Oliver J. D., and R. R. Colwell.** 1973. Extractable lipids of gram-negative marine bacteria: fatty acid composition. *Int. J. Syst. Bacteriol.* **114**:897–908.
16. **Ootaki, M., N. Morita, T. Nishida, M. Tanaka, A. Hase, Y. Yano, A. Yamada, R. Yu, K. Watanabe, and H. Okuyama.** 2003. Genes and pathways involved in biosynthesis of eicosapentaenoic and docosahexaenoic acids in bacteria, p. 49–52. *In* N. Murata, M. Yamada, I. Nishida, H. Okuyama, J. Sekiya, and H. Wada (ed.), *Advanced research on plant lipids*. Kluwer Academic Publishers, Dordrecht, Netherland.
17. **Orikasa, Y., T. Nishida, A. Hase, K. Watanabe, N. Morita, and H. Okuyama.** 2006. A phosphopantetheinyl transferase gene essential for biosynthesis of *n*–3 polyunsaturated fatty acids from *Moritella marina* strain MP-1. *FEBS Lett.* **580**:4423–4429.
18. **Orikasa, Y., T. Nishida, K. Watanabe, N. Morita, and H. Okuyama.** 2006. Phosphopantetheinyl transferase genes essential for biosynthesis of polyunsaturated fatty acids and their domain structures and compatibility. *In* C. Benning and J. Ohlrogge (ed.), *Advances in Plant Lipid Research: Proceedings of the 17th International Symposium on Plant Lipids, East Lansing, Michigan, July 2006* , Michigan State University Press, East Lansing, Michigan. in press.

19. **Orikasa, Y., T. Nishida, A. Yamada, R. Yu, K. Watanabe, A. Hase, N. Morita, and H. Okuyama** 2006. Recombinant production of docosahexaenoic acid in a polyketide biosynthesis mode in *Escherichia coli*. *Biotechnol Lett.* **28**:1841–1847.
20. **Orikasa, Y., A. Yamada, R. Yu, Y. Ito, T. Nishida, I. Yumoto, K. Watanabe, and H. Okuyama.** 2004. Characterization of the eicosapentaenoic acid biosynthesis gene cluster from *Shewanella* sp. strain SCRC-2738. *Cell. Mol. Biol.* **50**:625–630.
21. **Qi, B., T. Fraser, S. Mugford, G. Dobson, O. Sayanova, J. Butler, J. A. Napier, A. T. Stobart, and C. M. Lazarus.** 2004. Production of very long chain polyunsaturated omega-3 and omega-6 fatty acids in plants. *Nat. Biotechnol.* **22**:739–745.
22. **Rahman, A. S., J. Hothersall, J. Crosby, T. J. Simpson, and C. M. Thomas.** 2005. Tandemly duplicated acyl carrier proteins, which increase polyketide antibiotic production, can apparently function either in parallel or in series. *J. Biol. Chem.* **280**:6399–6408.
23. **Ratledge, C.** 2004. Fatty acid biosynthesis in microorganisms being used for single cell oil production. *Biochimie* **86**:807–815.
24. **Russell, N. J., and D. S. Nichols.** 1999. Polyunsaturated fatty acids in marine bacteria—a dogma rewritten. *Microbiology* **145**:767–779.
25. **Singh, A., and O. P. Ward.** 1997. Microbial production of docosahexaenoic acid (DHA, C22:6). *Adv. Appl. Microbiol.* **45**:271–312.
26. **Takeyama, H., D. Takeda, K. Yazawa, A. Yamada, and T. Matsunaga.** 1997. Expression of the eicosapentaenoic acid synthesis gene cluster from *Shewanella* sp. in

- a transgenic marine cyanobacterium, *Synechococcus* sp. Microbiology **143**:2725–2731.
27. **Tanaka, M., A. Ueno, K. Kawasaki, I. Yumoto, S. Ohgiya, T. Hoshino, K. Ishizaki, H. Okuyama, and N. Morita.** 1999. Isolation of clustered genes that are notably homologous to the eicosapentaenoic acid biosynthesis gene cluster from the docosahexaenoic acid-producing bacterium *Vibrio marinus* strain MP-1. Biotechnol. Lett. **21**:939–945.
 28. **Valentine, R. C., and D. L. Valentine.** 2004. Omega-3 fatty acids in cellular membranes: a unified concept. Prog. Lipid Res. **43**:383–402.
 29. **Venkateswaran K., D. P. Moser, M. E. Dollhopf, D. P. Lies, D. A. Saffarini, B. J. MacGregor, D. B. Ringelberg, D. C. White, M. Nishijima, H. Sano, J. Burghardt, E. Stackebrandt, and K. H. Nelson.** 1999. Polyphasic taxonomy of the genus *Shewanella* and description of *Shewanella oneidensis* sp. nov. Int. J. Syst. Bacteriol. **2**:705–724.
 30. **Vezzi, A., S. Campanaro, M. D'Angelo, F. Simonato, N. Vitulo, F. M. Lauro, A. Cestaro, G. Malacrida, B. Simionati, N. Cannata, C. Romualdi, D. H. Bartlett, and G. Valle.** 2005. Life at depth: *Photobacterium profundum* genome sequence and expression analysis. Science **307**:1459-1461.
 31. **Wirsen, C. O., H. W. Jannasch, S. G. Waleham, and E. A. Canuel.** 1987. Membrane lipids of a psychrophilic and barophilic deep-sea bacterium. Curr. Microbiol. **14**:319–322.

32. **Yano, Y, A. Nakayama, H. Saito, K. and Ishihara.** 1994. Production of docosahexaenoic acid by marine bacteria isolated from deep sea fish. *Lipids* **29**:527–528.
33. **Yazawa, K.** 1996. Production of eicosapentaenoic acid from marine bacteria. *Lipids* (Supplement) **31**:S297–S300.
34. **Yu, R., A. Yamada, K. Watanabe, K. Yazawa, H. Takeyama, T. Matsunaga, and R. Kurane.** 2000. Production of eicosapentaenoic acid by a recombinant marine cyanobacterium, *Synechococcus* sp. *Lipids* **35**:1061–1064.


FIGURE LEGENDS

FIG. 1. The organization of genes responsible for bacterial EPA and DHA biosynthesis and the domain structures of individual genes. The organization of gene clusters is divided into three types. Type I, which is represented by that of *Shewanella pneumatophori* SCRC-2738, is a gene cluster including all five *pfa* genes in a similar vicinity. Type II consists of a cluster of the four genes *pfaA–D* with *pfaE* separate from the other genes. This type of cluster is represented by that of *Moriella marina* MP-1. The relative direction of *pfaE* has not been determined for this bacterium. In Type III, *pfaE* is integrated into *pfaC/E* and the cluster is considered to consist of four genes. In this review, the remnant sequence downstream of the *pfaC/E* gene was regarded as a partial sequence of *pfaD*. The original nomenclature (Dai and Zhang, unpublished; ABF00130) is shown in parenthesis. The third type of cluster has been reported in *Pseudoalteromonas* sp. DS-12 only.

Type I (*Shewanella pneumatophori* SCRC-2738)



Type II (*Moritella marina* MP-1)



Type III (*Pseudoalteromonas* sp. DS-12)





TABLE 1. A list of known bacterial *pfa* genes^a.

Organisms	Product ^b	<i>pfa</i> genes in the cluster		ACP repeats in PfaA	Domains in PfaB	HD domains in PfaC or PfaC/E	<i>pfaE</i> (PfaE)		Recombinant synthesis of the product	Ref.
		Type ^c	Cloning				Group ^d	Cloning		
<i>S. pneumatophori</i> SCRC-2738	EPA	I	Yes	6	AT	FabA-FabA-FabA	I	Yes	Yes	20
<i>S. marinintestina</i> IK-1	EPA	I	Yes	6	AT	FabA-FabA-FabA	I	Yes	Yes	Ref. ^e
<i>Shewanella</i> sp. SC2A	EPA	– ^f	No	–	–	–	I	Yes	No	1
<i>S. oneidensis</i> MR-1	EPA	I	No	4	AT	FabA-FabA ^g	I	No	No	29
<i>Shewanella</i> sp. GA-22 ^h	EPA ⁱ	–	No	–	–	–	I	–	–	3
<i>P. profundum</i> SS9	EPA	II	Yes	6	AT	FabA-FabA ^g	II	No	No	1, 30
<i>Pseudoalteromonas</i> sp. DS-12	(EPA)	III	Yes	5	KS, KS	FabA-FabA ^j	I ^k	Yes	No	Ref. ¹
<i>M. marina</i> MP-1	DHA	II	Yes	5	KS, AT	FabA-FabZ/FabA-FabA	I	Yes	Yes	17,18
<i>C. psychrerythrae</i> 34H	(DHA)	I	No	6	KS, AT	FabA-FabA ^g	I	No	No	6

^a, Nucleic acid and deduced amino acid sequences were retrieved from databases (DDBJ/GenBank/EMBL:

<http://www.ddbj.nig.ac.jp/Welcome-j.html>). The name of each domain in individual *pfa* genes is referred to **the text and FIG. 1**.

^b, In cases of (EPA) and (DHA) the production of EPA or DHA is expected but not confirmed.

^c, Type I, II, and III *pfa* genes are defined in the text **and FIG. 1**.

^d, Group I and II PPTases are defined in the text and ref. 18.

^e, Unpublished result by N. Morita, Y. Yano, S. Ohgiya, and H. Okuyama.

^f, No information is available.

^g, and ^j, Unannotated 300–400 bp and 120–150 bp sequences, respectively, are present between the two FabA-like sequences.

^h, Only a 1,624 bp partial sequence *pfaA* is deposited (AJ563807).

ⁱ, Arachidonic acid (20:4n-6) and linoleic acid (18:2n-6) are also detected.

^k, *pfaE* is included in *pfaC/E*.

^l, See accession no. ABF00130.

TABLE 2. The recombinant production of EPA and/or DHA using *pfa* genes from *Shewanella pneumatophori* SCRC-2738,

Moritella marina MP-1 and *Photobacterium profundum* SS9^a

Source and combination of <i>pfa</i> genes			Production of	Ref.
<i>S. pneumatophori</i> SCRC-2738	<i>M. marina</i> MP-1	<i>P. profundum</i> SS9	EPA or DHA	
<i>pfaA,B,C,D,E</i> ^b			EPA	20
<i>pfaA,B,C,D</i> ^c plus <i>pfaE</i> ^d			EPA	20
<i>pfaA,B,C,D</i> ^c	<i>pfaE</i> ^e		EPA	17
	<i>pfaA,B,C,D</i> ^f plus <i>pfaE</i> ^g		DHA	19
<i>pfaB,C,D,E</i> ^h	<i>pfaA,B,C,D</i> ^f		EPA and DHA	18
<i>pfaE</i> ^d	<i>pfaA</i> [*] , <i>B,C,D</i> ⁱ		None	17
	<i>pfaA</i> [*] , <i>B,C,D</i> ⁱ plus <i>pfaE</i> ^g		None	17
		<i>pfaA,B,C,D</i> ⁱ plus <i>pfaE</i> ^k	None	1

^a, This table is modified from that of (18)

^b, ^c, ^f, and ^h, *pfa* genes are harbored in pEPAΔ1, pEPAΔ1,2,3, pDHA3, and pEPAΔ5, respectively.

^d, ^e, and ^g, *pfaE* gene is harbored in pSTV::*pfaE*, pETSTV::*pfaE*, and pET21::*pfaE*, respectively.

ⁱ, *pfa* genes are harbored in pDHA2, where *pfaA*^{*} is inactive.

^j, *pfa* genes are harbored in pFOS8E1.

^k, PPTase genes from *Bacillus subtilis* and *Shewanella* sp. SC2A are used (1).