



# HOKKAIDO UNIVERSITY

Title	PREPARATION OF AN ANTIGEN OF CHLAMYDIA PSITTACI SUITABLE FOR ENZYME-LINKED IMMUNOSORBENT ASSAY
Author(s)	ITOU, Norihiko
Citation	Japanese Journal of Veterinary Research, 36(2), 155-155
Issue Date	1988-05-20
Doc URL	<a href="https://hdl.handle.net/2115/3099">https://hdl.handle.net/2115/3099</a>
Type	departmental bulletin paper
File Information	KJ00002377083.pdf



PREPARATION OF AN ANTIGEN OF *CHLAMYDIA PSITTACI*  
SUITABLE FOR ENZYME-LINKED IMMUNOSORBENT ASSAY

Norihiko ITOU

*Department of Veterinary Public Health*  
*Faculty of Veterinary Medicine*  
*Hokkaido University, Sapporo 060, Japan*

In order to utilize enzyme-linked immunosorbent assay (ELISA) for the serodiagnosis of chlamydial infection, the basic conditions necessary to prepare the antigen were evaluated. Elementary bodies (EB) of *Chlamydia psittaci* were treated with  $\beta$ -propiolactone (BPL), sodium dodecyl sulfate (SDS) and  $\text{NaClO}_4$  (a chaotropic ion). These three kinds of treated antigens were compared by ELISA for their reactivities to antibodies against *Chlamydia psittaci*. Furthermore, these antigens were used for biotin-labeled antigen sandwich (BLAS) ELISA to detect chlamydial antibodies in human serum and that from a variety of animals, regardless of the species- and class-specificities of immunoglobulin.

The results were summarized as follows.

1. The antigenic components were not solublized from EB particles by 0.2% BPL, but were completely solublized by 1% SDS, and partly by 0.5M  $\text{NaClO}_4$ .
2. BPL-treated antigen showed weak reactivity against positive sera and was judged to be unsuitable for ELISA.
3. SDS-treated antigen showed high reactivity against positive sera, caused a high level of non-specific reaction to negative sera, and was revealed to be unavailable for ELISA.
4.  $\text{NaClO}_4$ -treated antigen reacted to positive sera with a high titer, caused a low level of non-specific reaction to negative sera and was judged to be useful for ELISA.
5. These three antigens were labile with  $\text{KIO}_4$ -treatment but resistant to pronase-treatment, which suggested that the antigenic components were carried on the carbohydrate moiety.
6. The three kinds of antigens were biotinylated and were used for BLAS-ELISA, but none of the antigens were applicable for the BLAS-ELISA due to the high levels of non-specific reactions.