



HOKKAIDO UNIVERSITY

Title	Roles of the hypocretin/orexins in the regulation of sleep and wakefulness
Author(s)	Terao, Akira; Haruyama, Takashi; Kimura, Kazuhiro
Citation	Japanese Journal of Veterinary Research, 55(2-3), 75-83
Issue Date	2008-02-09
DOI	https://doi.org/10.14943/jjvr.55.2-3.75
Doc URL	https://hdl.handle.net/2115/32367
Type	departmental bulletin paper
File Information	55(2-3)_75-83.pdf



Roles of the hypocretin/orexins in the regulation of sleep and wakefulness

Akira Terao*, Takashi Haruyama and Kazuhiro Kimura

Laboratory of Biochemistry, Department of Biomedical Sciences, Graduate School of Veterinary Medicine, Hokkaido University, Sapporo 060-0818, Japan

Received for publication, January 8, 2008; accepted, January 21, 2008

Abstract

Hypocretin/orexin is produced exclusively in the dorsal and lateral hypothalamus but its projection is widespread within the brain and plays important roles. In this paper, we review the independent discoveries of the hypocretin/orexin peptides, the neuroanatomy of this system, and the link to the sleep disorder narcolepsy that has led to the idea that this system plays a crucial role in the regulation of sleep and wakefulness.

Key words: hypocretin, orexin, narcolepsy, sleep, wakefulness

Discovery of the hypocretins and the orexins

Hypocretins 1 and 2 (Hcrt 1 and Hcrt 2) are hypothalamic neuropeptides derived from a single precursor molecule by proteolytic processing^{1,2)}. The mRNA encoding these peptides was originally identified in a directional tag PCR subtraction designed to enrich for hypothalamus-specific mRNAs¹⁾. Because the cell bodies expressing this gene were restricted to an area of the hypothalamus centered around the perifornical hypothalamus (PFH) and because of a weak homology to the gut peptide secretin, these molecules were called the “hypocretins.”

Hypocretins were independently discovered by another group of investigators as ligands binding to cell lines expressing orphan G protein-coupled receptors³⁾. The chemical isolation procedure that these investigators used allowed them to deter-

mine that the longer of these two peptides was 33 amino acids in length and to define the N-terminal pyroglutamyl residue, the intrachain disulfide links, and the expected C-terminal amidation of both peptides. Since intracerebroventricular injections of these peptides increased food intake in rats, these investigators called the peptides orexin-A and orexin-B. This paper also reported functional information on the receptors for the two peptides: the orexin-1 receptor (OX₁R) was shown to preferentially bind orexin-A over orexin-B, whereas the orexin-2 receptor (OX₂R) bound both peptides with similar affinity²⁾. In the subsequent issue of *Cell*, Sakurai et al. confirmed the identity of the Hcrts and the orexins³⁾. Thus, the *prepro-hypocretin* gene is identical to *prepro-orexin*; Hcrt 2 is identical to orexin-B; and Hcrt1 is equivalent to orexin-A. In this review, we use the terms “Hcrt 1/Ox-A” and “Hcrt2/Ox-B” to denote the two

*Corresponding author: Akira Terao, Laboratory of Biochemistry, Department of Biomedical Sciences, Graduate School of Veterinary Medicine, Hokkaido University, Sapporo 060-0818, Japan
Phone: +81-11-706-5205. Fax: +81-11-757-0703. E-mail: terao@vetmed.hokudai.ac.jp

peptides and “H/O” to refer to the Hcrt/orexin-containing cells or to the *hcrtr/orexin* gene. To refer to the receptors for these peptides, “HcrtR1/OX₁R” and “Hcrt/OX₂R” will be used.

Prepro-hypocretin mRNA is encoded by a gene localized to human chromosome 17q21²¹ which consists of two exons and one intron. The first exon includes the 5'-untranslated region and the region that encodes the first 7 amino acid residues of the secretory signal sequence. Exon 2 encodes the remaining portion of the signal sequence and the precursor molecule from which Hcrt1/Ox-A and Hcrt2/Ox-B are proteolytically cleaved⁴. Very little is known about the mechanisms regulating the expression of *prepro-hypocretin* at present. A 3.2 kb fragment of the 5'-upstream region cloned from the human Hcrt gene is sufficient to direct the expression of *Escherichia coli* β -galactosidase gene in transgenic mice in the lateral hypothalamic area and adjacent regions⁴. We have demonstrated that a 450 bp of the 5'-flanking region of the Hcrt gene has the ability to promote expression of a reporter gene *in vitro* and that this expression is down regulated by the sleep promoting cytokine, α -interferon⁵. The promoters of the receptor genes have not yet been characterized.

Neuroanatomy of the hypocretin/orexin system

The distribution of H/O cells has been described both by *in situ* hybridization^{1,2} and by immunohistochemistry using antibodies against Hcrt⁶, Hcrt1/Ox-A⁷⁻¹⁰ and Hcrt2/Ox-B^{8,11}. H/O expression is restricted to a few thousand cells in the tuberal hypothalamus centered in the PFH, dorsomedial hypothalamic nuclei and the lateral hypothalamic areas, encompassing about 1 mm anterior-posteriorly in the rat brain. Cells labeled by antisera against Hcrt1/Ox-A or Hcrt2/Ox-B are medium sized (20-30 μ m diameter) and multipolar to fusiform in shape. Hcrt1/Ox-A or Hcrt2/Ox-B seem to be co-localized in these cells⁸.

Although the H/O cells are a small population

with restricted expression, their projections are widely distributed in the brain⁶⁻¹⁰ and spinal cord¹¹. Particularly abundant processes are found throughout the hypothalamus⁶⁻⁹. The densest extrahypothalamic input is to the locus coeruleus (LC)^{6,12,13}. Other projection sites include the cerebral cortex, olfactory bulb, hippocampus, amygdala, septum, diagonal band of Broca, bed nucleus of the stria terminalis, thalamus, midbrain, brain stem, and spinal cord. H/O immunoreactivity has also been reported in the enteric nervous system and pancreas¹⁴ and H/O mRNA expression has been detected in testes².

The localization of the H/O-containing cells was reminiscent of another peptide found in this region, melanin-concentrating hormone (MCH). Although the distributions of H/O and MCH cells overlap, these peptides are not present within the same cells^{6,15,16}. The majority of H/O cells express leptin receptors and are immunoreactive for the transcription factor Stat-3^{13,17}. Isolated H/O cells are galanin-immunoreactive¹⁷. An antiserum raised against ovine prolactin (oPRL) apparently labels every H/O-containing cell in the rat hypothalamus¹⁸. Since oPRL-like immunoreactivity colocalized with dynorphin B^{19,20}, bradykinin²⁰, and secretogranin II²¹, these latter substances undoubtedly co-localize with H/O.

The concentrations of Hcrt1/Ox-A and Hcrt2/Ox-B have been measured in microdissected rat brain areas by radioimmunoassay^{22,23}. Hcrt1/Ox-A and Hcrt2/Ox-B were found to be abundant in the lateral and posterior hypothalamus, ventromedial hypothalamic (VMH), paraventricular thalamic and dorsal raphe nuclei (DRN), the periaqueductal gray and the LC. Moderate levels of the H/O peptides have also been measured in the median eminence, suprachiasmatic, arcuate, supraoptic, and paraventricular nuclei (PVN) of the hypothalamus, the substantia nigra, and the nucleus tractus solitarius. Low concentrations of Hcrt1/Ox-A and Hcrt2/Ox-B have been found in the cerebral cortex. In these studies, Hcrt2/Ox-B contents have always been 2-3 fold greater than the Hcrt1/Ox-A contents. Hcrt1/Ox-A has also been detected in cerebrospi-

nal fluid²⁴⁾ and in plasma²⁵⁾, but not in peripheral tissue or in the pineal and pituitary glands²⁶⁾. The Hcrt1/Ox-A distribution in hypothalamus apparently differs between males and females²⁶⁾.

HcrtR1/OX₁R and HcrtR1/OX₂R mRNAs are differentially distributed throughout the brain. In the hypothalamus, high levels of HcrtR1/OX₁R mRNA were initially reported in the VMH while HcrtR2/OX₂R mRNA was found to be highly expressed in the PVN²⁷⁾. More recent studies^{28,29)} have shown moderate to high levels found throughout the hypothalamus although some nuclei, such as the arcuate, mammillary and tuberomammillary nuclei (TMN), apparently express only the HcrtR2/OX₂R. High levels of HcrtR1/OX₁R mRNA have also been reported in the tenia tecta, the bed nucleus of the stria terminalis, the CA1 and CA2 fields of the hippocampus, the amygdalohippocampal area, the lateral dorsal tegmental and pedunculo pontine nuclei (LDT and PPT) and the LC. HcrtR2/OX₂R has mainly been found in olfactory tubercle, shell of the nucleus accumbens, the lateral and medial septal nuclei, central medial thalamic nuclei, the subthalamic and anterior pretectal nucleus, and the CA3 region of the hippocampus. In the cerebral cortex, layers III, V and VI also express HcrtR1/OX₁R mRNA whereas the HcrtR2/OX₂R mRNA is found in layers II and VI. Some brain regions such as the paraventricular thalamic nucleus, the DRN and ventral tegmental area (VTA) express the mRNA for both receptors in roughly equal abundance. The expression of H/O receptors has also been reported in the adrenal gland³⁰⁾, enteric nervous system, and pancreas¹⁴⁾.

The hypocretin/orexins and food intake regulation

As indicated above, H/O expression is restricted to the PFH, an area classically associated with the regulation of food intake. Since the original description of the orexigenic effects of the H/O peptides²⁾, other investigators have confirmed that

administration of Hcrt1/Ox-A into the lateral ventricles to fed rats stimulated food consumption³¹⁻³³⁾. Fasting has been shown to upregulate *prepro-H/O* mRNA levels²⁾. Injection of neutralizing anti Hcrt-antibodies significantly suppressed feeding in fasted rats³⁴⁾. Peripheral administration of a novel HcrtR1 antagonist inhibited natural and stimulated feeding for several days³⁵⁾.

The H/O neurons are in close proximity to the cells that contain MCH, a potent promoter of food consumption^{36,37)}. The composite distribution of the HcrtR1/OX₁R and HcrtR2/OX₂R mRNAs summarized above strongly resembles the distribution of MCH receptor mRNAs^{38,39)}, with the highest receptor levels for both systems being present within the hypothalamus, the LC, and the hippocampal formation. Despite the similarities between these two systems, H/O-and MCH-containing neurons apparently differ in their response to stimuli. H/O cells increase firing upon application of glucose⁴⁰⁾ whereas MCH cells are insensitive to deoxyglucose⁴¹⁾. Administration of agouti-related protein (AGRP) increases MCH but not *H/O* gene expression⁴²⁾, suggesting that the promoters of these genes respond to different signals that regulate food intake. Nonetheless, the similarity in the localization of the cell bodies for these two peptide systems and the overlapping distributions of their receptors strongly suggests an interaction between the two systems in some homeostatic functions³⁹⁾.

As indicated above, the widespread anatomical projections of the H/O neurons suggest that they may be involved in multiple functions. In addition to food intake regulation, some of the functions in which this system has been implicated include neuroendocrine⁴³⁻⁴⁷⁾, cardiovascular^{48,49)}, water balance⁵⁰⁾, and gastrointestinal⁵¹⁾ control.

The hypocretin/orexins and the sleep disorder narcolepsy

Two independent lines of research converged in 1999 to implicate the H/O system in the sleep disorder narcolepsy. Narcolepsy is a neurological

disorder that specifically affects the generation and organization of sleep⁵²⁾. Narcoleptic patients have persistent abnormal daytime sleepiness and fragmented sleep at night^{4,52-54)}. The pathognomic symptom of narcolepsy is cataplexy, the sudden loss of skeletal muscle tone for periods that can vary from seconds to minutes that is usually precipitated by strong emotion^{52,54)}. Other symptoms include sleep paralysis, hypnagogic hallucinations and sleep onset rapid eye movement (REM) periods: while normal individuals begin their rest with non-REM sleep, narcoleptic patients go directly into REM sleep, often accompanied by dreamlike hallucinations and muscle tone paralysis⁵⁴⁾. Because of these traits, it has been hypothesized that the symptoms of narcolepsy result from the malfunction of some aspects of the REM sleep control mechanism⁵⁵⁾.

A genetic component for narcolepsy has been established in both humans and dogs. In narcoleptic dogs, the *canarc-1* mutation is transmitted in Doberman pinschers and Labrador retrievers as an autosomal recessive trait with full penetrance. The first link between the H/O system and narcolepsy was identification of *canarc-1* as a deletion mutation in the *Hcrtr2/OX₂R* gene which results in a truncated, nonfunctional protein⁵⁶⁾. In a remarkable convergence, *H/O* null mutant mice were found to exhibit periods of "behavioral arrest" that strongly resemble the cataplectic attacks and sleep-onset REM periods characteristic of narcolepsy in other species⁵⁷⁾. These mice also have an altered sleep architecture, as evidenced by increased levels of both REM and non-REM sleep, short latency REM periods, and decreased sleep bout lengths, primarily during the dark (active) period. Together, these two papers indicate that dysfunction of either the H/O ligand or one of its receptors can result in narcolepsy.

Although mutations in genes of the hypocretin system are directly responsible for the etiology of narcolepsy in animals, human narcolepsy appears to be different and, in most cases, seems to be transmitted as a multifactorial trait. Genetic studies suggest a relationship between human

narcolepsy and the immune system⁵⁸⁻⁶⁰⁾. Across ethnic groups, expression of the MHC class II allele HLA-DQB1*0602 confers increased susceptibility to narcolepsy²²⁾. Studies of familial narcolepsy also demonstrate that environmental factors acting on a specific genetic background may contribute to susceptibility to narcolepsy⁶¹⁾. The high degree of discordance in identical twins further implicates environmental influences. Nonetheless, at least one human narcoleptic bears a mutation in the *H/O* gene⁶²⁾ but no receptor mutations have been reported to date. The likely cause of narcolepsy in the majority of human cases was recently described. An abnormality in H/O neurotransmission in narcoleptic humans was first suggested by undetectable levels of Hcrtr1/Ox-A in cerebrospinal fluid (CSF) from 7 of 9 narcoleptic patients²⁴⁾. Subsequent studies show that human narcolepsy is associated with loss of H/O-containing cell bodies in the PFH. In postmortem analyses, *H/O* mRNA was undetectable in two narcoleptic brains, although mRNA for MCH was readily detectable in both controls and narcoleptics⁶²⁾. Immunohistochemistry showed an 85-95% reduction in the number of H/O-containing cells in narcoleptic brains with no evident change in the number of MCH cells⁶³⁾. These studies suggest that degeneration of the H/O cells may be the cause of human narcolepsy. Increased staining for glial fibrillary acid protein in the PFH combined with the previously described genetic link with the HLA system suggests that the H/O cells may degenerate by an autoimmune mechanism⁶³⁾.

The hypocretin/orexins and sleep/wake regulation

The involvement of H/O system in the pathophysiology of narcolepsy suggests a role for this system in 'normal' sleep/wake regulation. The region of the posterior hypothalamus (PH) containing the H/O cells has long been implicated in arousal state control. Subsequent to World War I, a viral pandemic resulted in *encephalitis lethar-*

gica in which patients experienced persistent sleepiness that was attributed to injury of the PH and rostral midbrain⁶⁴). Experimental lesions of the posterior lateral hypothalamus also causes persistent hypersomnia in monkeys⁶⁵, rats⁶⁶ and cats⁶⁷). The H/O cells in the PH project to the monoaminergic and cholinergic groups historically implicated in arousal state regulation. Intracerebroventricular injections of Hcrt1/Ox-A into rats at light onset (the major sleep period) increases arousal and locomotor activity accompanied by a marked reduction in REM sleep and deep slow wave sleep⁶⁸). A diurnal variation in Hcrt1/Ox-A immunoreactivity has been reported in the pons and in the preoptic/anterior hypothalamic region and in *prepro-hcrt* mRNA in the hypothalamus²⁶), further supporting involvement of the H/O system in arousal state regulation. On the other hand, *prepro-hcrt* mRNA levels were unaffected in either rats and mice by a period of sleep deprivation sufficient to cause elevation of EEG delta power activity during recovery sleep after sleep deprivation⁶⁹).

Any influence of the H/O system on sleep and wakefulness is likely to be manifest through projections of the H/O cells onto the monoaminergic and cholinergic cell groups⁷⁰). In addition to intense innervation of the LC mentioned above⁶), numerous synaptic contacts have been observed between H/O boutons and noradrenergic cells of the LC¹³). H/O terminals are also in close proximity to histaminergic cells of the TMN, serotonergic cells of the DRN and cholinergic cells in both the pons (LDT and PPT) and basal forebrain (diagonal band of Broca and the medial septum)⁵⁷). These cholinergic areas are thought to be involved in the EEG desynchronization characteristic of waking and REM sleep.

Diagnosis/drug treatment of narcolepsy

Current treatment for narcoleptic patients employ monoaminergic amphetamine-like and antidepressant compounds⁷¹). These compounds have many side effects and only partially control the

condition. The loss of H/O neurons in the hypothalamus^{62,63}), reduction of Hcrt1/Ox-A levels in the CSF of narcoleptic patients²⁴), and the existence of a mutation in the *H/O* gene of a child with severe symptoms⁶²) strongly supports a central role for H/O deficiency in the human narcolepsy and suggests several possibilities for improved diagnosis and treatments. First, monitoring CSF Hcrt1/Ox-A levels will be useful for diagnosis of narcolepsy. Second, supplementation of H/O transmission with receptor agonists will likely be an effective treatment strategy, at least for the sleepiness aspects of narcolepsy since, fortunately, human narcoleptics do not have the receptor mutation found in narcoleptic canines⁶²). Conversely, selective antagonists for H/O receptors might be also useful for sleep promotion, since H/O system seems to be important for maintenance of wakefulness. These compounds should be stable *in vivo* to have a long lasting effect and also, should cross blood-brain barrier so that the drug can be given peripherally. The first orexin receptor antagonist for insomnia treatment was recently reported⁷²) and more compounds will follow in the near future. Regulation of the activity of the H/O neurons is another potential therapeutic avenue.

References

- 1) de Lecea, L., Kilduff, T. S., Peyron, C., Gao, X.-B., Foye, P. E., Danielson, P. E., Fukuhara, C., Battenberg, E. L. F., Gautvik, V. T., Bartlett II, F. S., Frankel, W. N., van den Pol, A. N., Bloom, F. E., Gautvik, K. M. and Sutcliffe, J. G. 1998. The hypocretins: hypothalamus-specific peptides with neuroexcitatory activity. *Proc. Natl. Acad. Sci. U.S.A.*, **95**: 322-327.
- 2) Sakurai, T., Amemiya, A., Ishii, M., Matsuzaki, I., Chemelli, R. M., Tanaka, H., Williams, S. C., Richardson, J. A., Kozlowski, G. P., Wilson, S., Arch, J. R., Buckingham, R. E., Haynes, A. C., Carr, S. A., Annan, R. S., McNulty, D. E., Liu, W. S., Terrett, J. A., Elshourbagy, N. A., Bergsma, D. J. and Yanagisawa, M. 1998. Orexins and

- orexin receptors: a family of hypothalamic neuropeptides and G protein-coupled receptors that regulate feeding behavior. *Cell*, **92**: 573-585.
- 3) Sakurai, T., Amemiya, A., Ishii, M., Matsuzaki, I., Chemelli, R. M., Tanaka, H., Williams, S. C., Richardson, J. A., Kozlowski, G. P., Wilson, S., Arch, J. R., Buckingham, R. E., Haynes, A. C., Carr, S. A., Annan, R. S., McNulty, D. E., Liu, W. S., Terrett, J. A., Elshourbagy, N. A., Bergsma, D. J. and Yanagisawa, M. 1998. Orexins and orexin receptors: a family of hypothalamic neuropeptides and G protein-coupled receptors that regulate feeding behavior. *Cell*, **92**: 1 page following 696.
 - 4) Sakurai, T., Moriguchi, T., Furuya, K., Kajiwara, N., Nakamura, T., Yanagisawa, M. and Goto, K. 1999. Structure and function of human prepro-orexin gene. *J. Biol. Chem.*, **274**: 17771-17776.
 - 5) Waleh, N. S., Apte-Deshpande, A., Terao, A., Ding, J. and Kilduff, T. S. 2001. Modulation of the promoter region of prepro-hypocretin by alpha-interferon. *Gene*, **262**: 123-128.
 - 6) Peyron, C., Tighe, D. K., van den Pol, A. N., de Lecea, L., Heller, H. C., Sutcliffe, J. G. and Kilduff, T. S. 1998. Neurons containing hypocretin (orexin) project to multiple neuronal systems. *J. Neurosci.*, **18**: 9996-10015.
 - 7) Chen, C. T., Dun, S. L., Kwok, E. H., Dun, N. J. and Chang, J. K. 1999. Orexin A-like immunoreactivity in the rat brain. *Neurosci. Lett.*, **260**: 161-164.
 - 8) Date, Y., Ueta, Y., Yamashita, H., Yamaguchi, H., Matsukura, S., Kangawa, K., Sakurai, T., Yanagisawa, M. and Nakazato, M. 1999. Orexins, orexigenic hypothalamic peptides, interact with autonomic, neuroendocrine and neuroregulatory systems. *Proc. Natl. Acad. Sci. U. S. A.*, **96**: 748-753.
 - 9) Nambu, T., Sakurai, T., Mizukami, K., Hosoya, Y., Yanagisawa, M. and Goto, K. 1999. Distribution of orexin neurons in the adult rat brain. *Brain Res.*, **827**: 243-260.
 - 10) Harrison, T. A., Chen, C. T., Dun, N. J. and Chang, J. K. 1999. Hypothalamic orexin A-immunoreactive neurons project to the rat dorsal medulla. *Neurosci. Lett.*, **273**: 17-20.
 - 11) van den Pol, A. N. 1999. Hypothalamic hypocretin (orexin): robust innervation of the spinal cord. *J. Neurosci.*, **19**: 3171-3182.
 - 12) Hagan, J. J., Leslie, R. A., Patel, S., Evans, M. L., Wattam, T. A., Holmes, S., Benham, C. D., Taylor, S. G., Routledge, C., Hemmati, P., Munton, R. P., Ashmeade, T. E., Shah, A. S., Hatcher, J. P., Hatcher, P. D., Jones, D. N., Smith, M. I., Piper, D. C., Hunter, A. J., Porter, R. A. and Upton, N. 1999. Orexin A activates locus coeruleus cell firing and increases arousal in the rat. *Proc. Natl. Acad. Sci. U. S. A.*, **96**: 10911-10916.
 - 13) Horvath, T. L., Peyron, C., Diano, S., Ivanov, A., Aston-Jones, G., Kilduff, T. S. and van Den Pol, A. N. 1999. Hypocretin (orexin) activation and synaptic innervation of the locus coeruleus noradrenergic system. *J. Comp. Neurol.*, **415**: 145-159.
 - 14) Kirchgeßner, A. L. and Liu, M. 1999. Orexin synthesis and response in the gut. *Neuron*, **24**: 941-951.
 - 15) Broberger, C., De Lecea, L., Sutcliffe, J. G. and Hokfelt, T. 1998. Hypocretin/orexin-and melanin-concentrating hormone-expressing cells form distinct populations in the rodent lateral hypothalamus: relationship to the neuropeptide Y and agouti gene-related protein systems. *J. Comp. Neurol.*, **402**: 460-474.
 - 16) Elias, C. F., Saper, C. B., Maratos-Flier, E., Tritos, N. A., Lee, C., Kelly, J., Tatro, J. B., Hoffman, G. E., Ollmann, M. M., Barsh, G. S., Sakurai, T., Yanagisawa, M. and Elmquist, J. K. 1998. Chemically defined projections linking the mediobasal hypothalamus and the lateral hypothalamic area. *J. Comp. Neurol.*, **402**: 442-459.
 - 17) Hakansson, M., de Lecea, L., Sutcliffe, J. G., Yanagisawa, M. and Meister, B. 1999. Leptin receptor-and STAT3-immunoreactivities in hypocretin/orexin neurones of the lateral hypothalamus. *J. Neuroendocrinol.*, **11**: 653-663.

- 18) Risold, P. Y., Griffond, B., Kilduff, T. S., Sutcliffe, J. G. and Fellmann, D. 1999. Preprohypocretin (orexin) and prolactin-like immunoreactivity are coexpressed by neurons of the rat lateral hypothalamic area. *Neurosci. Lett.*, **259**: 153-156.
- 19) Griffond, B., Deray, A., Fellmann, D., Ciofi, P., Croix, D. and Bugnon, C. 1993. Colocalization of prolactin-and dynorphin-like substances in a neuronal population of the rat lateral hypothalamus. *Neurosci. Lett.*, **156**: 91-95.
- 20) Griffond, B., Deray, A., Jacquemard, C., Fellmann, D. and Bugnon, C. 1994. Prolactin immunoreactive neurons of the rat lateral hypothalamus: immunocytochemical and ultrastructural studies. *Brain Res.*, **635**: 179-186.
- 21) Griffond, B., Grillon, S., Duval, J., Colard, C., Jacquemard, C., Deray, A. and Fellmann, D. 1995. Occurrence of secretogranin II in the prolactin-immunoreactive neurons of the rat lateral hypothalamus: an in situ hybridization and immunocytochemical study. *J. Chem. Neuroanat.*, **9**: 113-119.
- 22) Mondal, M. S., Nakazato, M., Date, Y., Murakami, N., Yanagisawa, M. and Matsukura, S. 1999. Widespread distribution of orexin in rat brain and its regulation upon fasting. *Biochem. Biophys. Res. Commun.*, **256**: 495-499.
- 23) Mondal, M. S., Nakazato, M., Date, Y., Murakami, N., Hanada, R., Sakata, T. and Matsukura, S. 1999. Characterization of orexin-A and orexin-B in the microdissected rat brain nuclei and their contents in two obese rat models. *Neurosci. Lett.*, **273**: 45-48.
- 24) Nishino, S., Ripley, B., Overeem, S., Lammers, G. J. and Mignot, E. 2000. Hypocretin (orexin) deficiency in human narcolepsy. *Lancet*, **355**: 39-40.
- 25) Arihara, Z., Takahashi, K., Murakami, O., Totsune, K., Sone, M., Satoh, F., Ito, S. and Mouri, T. 2001. Immunoreactive orexin-A in human plasma. *Peptides*, **22**: 139-142.
- 26) Taheri, S., Mahmoodi, M., Opacka-Juffry, J., Ghatei, M. A. and Bloom, S. R. 1999. Distribution and quantification of immunoreactive orexin A in rat tissues. *FEBS Lett.*, **457**: 157-161.
- 27) Trivedi, P., Yu, H., Mac Neil, D. J., Van der Ploeg, L. H. and Guan, X. M. 1998. Distribution of orexin receptor mRNA in the rat brain. *FEBS Lett.*, **438**: 71-75.
- 28) Lu, X. Y., Bagnol, D., Burke, S., Akil, H. and Watson, S. J. 2000. Differential distribution and regulation of OX1 and OX2 orexin/hypocretin receptor messenger RNA in the brain upon fasting. *Horm. Behav.*, **37**: 335-344.
- 29) Marcus, J. N., Aschkenasi, C. J., Lee, C. E., Chemelli, R. M., Saper, C. B., Yanagisawa, M. and Elmquist, J. K. 2001. Differential expression of orexin receptors 1 and 2 in the rat brain. *J. Comp. Neuro.*, **1435**: 6-25.
- 30) Malendowicz, L. K., Tortorella, C. and Nussdorfer, G. G. 1999. Orexins stimulate corticosterone secretion of rat adrenocortical cells, through the activation of the adenylate cyclase-dependent signaling cascade. *J. Steroid Biochem. Mol. Biol.*, **70**: 185-188.
- 31) Edwards, C. M., Abusnana, S., Sunter, D., Murphy, K. G., Ghatei, M. A. and Bloom, S. R. 1999. The effect of the orexins on food intake: comparison with neuropeptide Y, melanin-concentrating hormone and galanin. *J. Endocrinol.*, **160**: R7-12.
- 32) Haynes, A. C., Jackson, B., Overend, P., Buckingham, R. E., Wilson, S., Tadayyon, M. and Arch, J. R. 1999. Effects of single and chronic intracerebroventricular administration of the orexins on feeding in the rat. *Peptides*, **20**: 1099-1105.
- 33) Yamanaka, A., Kunii, K., Nambu, T., Tsujino, N., Sakai, A., Matsuzaki, I., Miwa, Y., Goto, K. and Sakurai, T. 2000. Orexin-induced food intake involves neuropeptide Y pathway. *Brain Res.*, **859**: 404-409.
- 34) Yamada, H., Okumura, T., Motomura, W., Kobayashi, Y. and Kohgo, Y. 2000. Inhibition of food intake by central injection of anti-orexin antibody in fasted rats. *Biochem. Biophys. Res. Commun.*, **267**: 527-531.
- 35) Arch, J. 2000. Orexins, feeding and the big

- picture. *Br. J. Nutr.*, **84**: 401-403.
- 36) Qu, D., Ludwig, D. S., Gammeltoft, S., Piper, M., Pelleymounter, M. A., Cullen, M. J., Mathes, W. F., Przypek, R., Kanarek, R. and Maratos-Flier, E. 1996. A role for melanin-concentrating hormone in the central regulation of feeding behaviour. *Nature*, **380**: 243-247.
 - 37) Shimada, M., Tritos, N. A., Lowell, B. B., Flier, J. S. and Maratos-Flier, E. 1998. Mice lacking melanin-concentrating hormone are hypophagic and lean. *Nature*, **396**: 670-674.
 - 38) Saito, Y., Cheng, M., Leslie, F. M. and Civelli, O. 2001. Expression of the melanin-concentrating hormone (MCH) receptor mRNA in the rat brain. *J. Comp. Neurol.*, **435**: 26-40.
 - 39) Kilduff, T. S. and de Lecea, L. 2001. Mapping of the mRNAs for the hypocretin/orexin and melanin-concentrating hormone receptors: networks of overlapping peptide systems. *J. Comp. Neurol.*, **435**: 1-5.
 - 40) Bayer, L., Colard, C., Nguyen, N. U., Risold, P. Y., Fellmann, D. and Griffond, B. 2000. Alteration of the expression of the hypocretin (orexin) gene by 2-deoxyglucose in the rat lateral hypothalamic area. *Neuroreport*, **11**: 531-533.
 - 41) Bayer, L., Poncet, F., Fellmann, D. and Griffond, B. 1999. Melanin-concentrating hormone expression in slice cultures of rat hypothalamus is not affected by 2-deoxyglucose. *Neurosci. Lett.*, **267**: 77-80.
 - 42) Hanada, R., Nakazato, M., Matsukura, S., Murakami, N., Yoshimatsu, H. and Sakata, T. 2000. Differential regulation of melanin-concentrating hormone and orexin genes in the agouti-related protein/melanocortin-4 receptor system. *Biochem. Biophys. Res. Commun.*, **268**: 88-91.
 - 43) van den Pol, A. N., Gao, X. B., Obrietan, K., Kilduff, T. S. and Belousov, A. B. 1998. Presynaptic and postsynaptic actions and modulation of neuroendocrine neurons by a new hypothalamic peptide, hypocretin/orexin. *J. Neurosci.*, **18**: 7962-7971.
 - 44) Pu, S., Jain, M. R., Kalra, P. S. and Kalra, S. P. 1998. Orexins, a novel family of hypothalamic neuropeptides, modulate pituitary luteinizing hormone secretion in an ovarian steroid-dependent manner. *Regul. Pept.*, **78**: 133-136.
 - 45) Tamura, T., Irahara, M., Tezuka, M., Kiyokawa, M. and Aono, T. 1999. Orexins, orexigenic hypothalamic neuropeptides, suppress the pulsatile secretion of luteinizing hormone in ovariectomized female rats. *Biochem. Biophys. Res. Commun.*, **264**: 759-762.
 - 46) Mitsuma, T., Hirooka, Y., Mori, Y., Kayama, M., Adachi, K., Rhue, N., Ping, J. and Nogimori, T. 1999. Effects of orexin A on thyrotropin-releasing hormone and thyrotropin secretion in rats. *Horm. Metab. Res.*, **31**: 606-609.
 - 47) Lopez, M., Senaris, R., Gallego, R., Garcia-Caballero, T., Lago, F., Seoane, L., Casanueva, F. and Dieguez, C. 1999. Orexin receptors are expressed in the adrenal medulla of the rat. *Endocrinology*, **140**: 5991-5994.
 - 48) Shirasaka, T., Nakazato, M., Matsukura, S., Takasaki, M. and Kannan, H. 1999. Sympathetic and cardiovascular actions of orexins in conscious rats. *Am. J. Physiol.*, **277**: R1780-1785.
 - 49) Samson, W. K., Gosnell, B., Chang, J. K., Resch, Z. T. and Murphy, T. C. 1999. Cardiovascular regulatory actions of the hypocretins in brain. *Brain Res.*, **831**: 248-253.
 - 50) Kunii, K., Yamanaka, A., Nambu, T., Matsuzaki, I., Goto, K. and Sakurai, T. 1999. Orexins/hypocretins regulate drinking behaviour. *Brain Res.*, **842**: 256-261.
 - 51) Takahashi, N., Okumura, T., Yamada, H. and Kohgo, Y. 1999. Stimulation of gastric acid secretion by centrally administered orexin-A in conscious rats. *Biochem. Biophys. Res. Commun.*, **254**: 623-627.
 - 52) Aldrich, M. S. 1993. The neurobiology of narcolepsy-cataplexy. *Prog. Neurobiol.*, **41**: 533-541.
 - 53) Nishino, S. and Mignot, E. 1997. Pharmacological aspects of human and canine narcolepsy. *Prog. Neurobiol.*, **52**: 27-78.

- 54) Aldrich, M. S. 1998. Diagnostic aspects of narcolepsy. *Neurology*, **50**: S2-7.
- 55) Siegel, J. M. 1999. Narcolepsy: a key role for hypocretins (orexins). *Cell*, **98**: 409-412.
- 56) Lin, L., Faraco, J., Li, R., Kadotani, H., Rogers, W., Lin, X., Qiu, X., de Jong, P. J., Nishino, S. and Mignot, E. 1999. The sleep disorder canine narcolepsy is caused by a mutation in the hypocretin (orexin) receptor 2 gene. *Cell*, **98**: 365-376.
- 57) Chemelli, R. M., Willie, J. T., Sinton, C. M., Elmquist, J. K., Scammell, T., Lee, C., Richardson, J. A., Williams, S. C., Xiong, Y., Kisanuki, Y., Fitch, T. E., Nakazato, M., Hammer, R. E., Saper, C. B. and Yanagisawa, M. 1999. Narcolepsy in orexin knockout mice: molecular genetics of sleep regulation. *Cell*, **98**: 437-451.
- 58) Matsuki, K., Grumet, F. C., Lin, X., Gelb, M., Guilleminault, C., Dement, W. C. and Mignot, E. 1992. DQ (rather than DR) gene marks susceptibility to narcolepsy. *Lancet*, **339**: 1052.
- 59) Mignot, E., Lin, X., Arrigoni, J., Macaubas, C., Olive, F., Hallmayer, J., Underhill, P., Guilleminault, C., Dement, W. C. and Grumet, F. C. 1994. DQB1*0602 and DQA1*0102 (DQ 1) are better markers than DR2 for narcolepsy in Caucasian and black Americans. *Sleep*, **17**: S60-67.
- 60) Mignot, E., Tafti, M., Dement, W. C. and Grumet, F. C. 1995. Narcolepsy and immunity. *Adv. Neuroimmunol.*, **5**: 23-37.
- 61) Mignot, E. 1998. Genetic and familial aspects of narcolepsy. *Neurology*, **50**: S16-22.
- 62) Peyron, C., Faraco, J., Rogers, W., Ripley, B., Overeem, S., Charnay, Y., Nevsimalova, S., Aldrich, M., Reynolds, D., Albin, R., Li, R., Hungs, M., Pedrazzoli, M., Padigar, M., Kucherlapati, M., Fan, J., Maki, R., Lammers, G. J., Bouras, C., Kucherlapati, R., Nishino, S. and Mignot, E. 2000. A mutation in a case of early onset narcolepsy and a generalized absence of hypocretin peptides in human narcoleptic brains. *Nat. Med.*, **6**: 991-997.
- 63) Thannickal, T., Moore, R., Y., Nienhuis, R., Ramanathan, L., Gulyani, S., Aldrich, M., Cornford, M. and Siegel, J. M. 2000. Reduced number of hypocretin neurons in human narcolepsy. *Neuron*, **27**: 469-474.
- 64) von Economo, C. 1930. Sleep as a problem of localization. *J. Nerv. Ment. Dis.*, **71**: 249-259.
- 65) Ranson, S. W. 1939. Somnolence caused by hypothalamic lesions in the monkey. *Arch. Neurol. Psychiat.*, **41**: 1-23.
- 66) Nauta, W. J. H. 1946. Hypothalamic regulation of sleep in rats. An experimental study. *J. Neurophysiol.*, **9**: 285-316.
- 67) Swett, C. and Hobson, J. 1968. The effects of posterior hypothalamic lesions on behavioral and electrographic manifestations of sleep and waking in cats. *Arch. Ital. Biol.*, **106**: 270-282.
- 68) Piper, D. C., Upton, N., Smith, M. I. and Hunter, A. J. 2000. The novel brain neuropeptide, orexin-A, modulates the sleep-wake cycle of rats. *Eur. J. Neurosci.*, **12**: 726-730.
- 69) Terao, A., Peyron, C., Ding, J., Wurts, S. W., Edgar, D. M., Heller, H. C. and Kilduff, T. S. 2000. Prepro-hypocretin (prepro-orexin) expression is unaffected by short-term sleep deprivation in rats and mice. *Sleep*, **23**: 867-874.
- 70) Kilduff, T. S. and Peyron, C. 2000. The hypocretin/orexin ligand-receptor system: Implications for sleep and sleep disorders. *Trends Neurosci.*, **23**: 359-365.
- 71) Guilleminault, C., Heinzer, R., Mignot, E. and Black, J. 1998. Investigations into the neurologic basis of narcolepsy. *Neurology*, **50**: S8-15.
- 72) Brisbare-Roch, C., Dingemans, J., Koberstein, R., Hoeber, P., Aissaoui, H., Flores, S., Mueller, C., Nayler, O., van Gerven, J., de Haas, S. L., Hess, P., Qiu, C., Buchmann, S., Scherz, M., Weller, T., Fischli, W., Clozel, M. and Jenck, F. 2007. Promotion of sleep by targeting the orexin system in rats, dogs and humans. *Nat. Med.*, **13**: 150-155.