



HOKKAIDO UNIVERSITY

Title	TWO CALCIUM REGULATORY SYSTEMS IN SQUID MANTLE MUSCLE
Author(s)	Konno, Kunihiro
Degree Grantor	北海道大学
Degree Name	博士(水産学)
Dissertation Number	乙第2120号
Issue Date	1981-06-30
Doc URL	https://hdl.handle.net/2115/32710
Type	doctoral thesis
File Information	2120.pdf



TWO CALCIUM REGULATORY SYSTEMS IN SQUID MANTLE MUSCLE

by

Kunihiko KONNO

B. F., Hokkaido University, Hakodate, Japan, 1974

M. F., Hokkaido University, Hakodate, Japan, 1976

A DISSERTATION

submitted in partial fulfilment of
the requirements for the degree

DOCTOR OF FISHERIES SCIENCE

Laboratory of Biochemistry

Department of Food Science

Faculty of Fisheries

Hokkaido University, Hakodate, Japan

1981

CONTENTS

I	Introduction	1
II	Methods and materials	3
1	Preparative methods for muscular proteins	3
1-(A)	Myosin B and myosin from squid mantle muscle	3
1-(B)	Actin from squid mantle muscle	5
1-(C)	Native tropomyosin from squid mantle muscle	5
1-(D)	Troponin and tropomyosin from squid mantle muscle	6
1-(E)	Myosin, desensitized myosin and EDTA-LC from scallop adductor muscle	7
1-(F)	Light chains from squid mantle muscle	7
1-(G)	Myosin and actin from rabbit skeletal muscle	10
1-(H)	Glycerinated muscle fibers from scallop adductor muscle	10
2	Assay method for biological and physico-chemical properties	11
2-(A)	Assay for ATPase activity	11
2-(B)	Measurement of superprecipitation	12
2-(C)	SDS gel, urea gel and disc gel electrophoreses and densitometry	12

2-(D)	UV absorption spectra of light chains and divalent cation induced difference spectra of them	13
2-(E)	Calcium binding of myosin and light chain	13
2-(F)	Quantative determination of protein concentration	14
2-(G)	Tension determination of glycerinated muscle fibers of scallop muscle	14
3	Materials and reagents	16
III	Results	18
1	Actin-linked Ca regulation	20
1-(A)	Native tropomyosin in squid muscle	20
1-(A)-a)	Ca sensitivity of squid acto-skeletal myosin	20
1-(A)-b)	Isolation of native tropomyosin from squid muscle	23
1-(A)-c)	A comparison between native tropo- myosin from rabbit and squid muscles	25
1-(A)-d)	Binding of squid native tropomyosin to rabbit F-actin	27
1-(B)	Troponin and tropomyosin in squid muscle	32
1-(B)-a)	Isolation of troponin and tropomyosin from squid muscle	32
1-(B)-b)	SDS gel electrophoretic patterns of squid troponin and tropomyosin	32

1-(B)-c)	Effect of squid troponin, tropomyosin and their complex on superprecipitation of skeletal actomyosin	36
1-(B)-d)	A comparison between troponins from rabbit and squid muscles	38
1-(B)-e)	Binding of squid troponin and tropo- myosin to F-actin	41
2	Myosin-linked Ca regulation	44
2-(A)	Ca sensitive myosin from squid mantle muscle	44
2-(A)-a)	SDS gel electrophoretic patterns of squid myosin and myosin B	44
2-(A)-b)	Ca sensitivity of skeletal acto- squid myosin	44
2-(A)-c)	EDTA-treatment of squid myosin	49
2-(B)	A comparison of Ca and Sr requirements for Mg-ATPase activity of myosin-linked and actin-linked regulations	51
2-(B)-a)	Calcium or strontium ion requirement for squid myosin B Mg-ATPase	51
2-(B)-b)	Effects of squid and rabbit native tropomyosin on the Ca requirement of Mg-ATPase of squid myosin B	53
2-(C)	Regulatory light chain from squid myosin	55
2-(C)-a)	Isolation of highly purified squid light chains	55

2-(C)-b)	Effect of 2-mercaptoethanol on the patterns of squid light chain in disc gel electrophoresis	59
2-(C)-c)	Effect of calcium on the patterns of squid light chains in disc gel electrophoresis	59
2-(C)-d)	UV absorption spectra of isolated squid light chains and their changes induced by calcium or strontium ions	62
2-(C)-e)	UV absorption spectrum of scallop EDTA-LC and its change induced by calcium or strontium ions	66
2-(D)	Desensitization and resensitization of scallop myosin	69
2-(D)-a)	Desensitization of scallop myosin by EDTA-treatment	69
2-(D)-b)	Binding of squid light chains to scallop desensitized myosin	69
2-(D)-c)	Resensitization of scallop desensitized myosin by squid light chain	71
2-(D)-d)	Relationship between EDTA-LC(or squid LC-2) content of myosin and Ca sensitivity of acto-scallop myosin	75
2-(D)-e)	Competitive binding of EDTA-LC and squid LC-2 to scallop desensitized myosin	78

2-(E)	Calcium binding of myosin and regulatory light chain	80
2-(E)-a)	Calcium binding of isolated squid LC-2 and scallop EDTA-LC	80
2-(E)-b)	Calcium binding of squid and scallop myosin	82
2-(E)-c)	Effect of desensitization and resensi- tization on the calcium binding of scallop myosin	84
2-(F)	Calcium sensitivity of squid myosin alone	87
2-(F)-a)	Ca requirement for activation of Mg-ATPase of squid myosin alone	87
2-(F)-b)	Effects of temperature and pH on Ca sensitivity of squid myosin	91
2-(F)-c)	Effect of KCl concentration on Ca sensitivity of squid myosin	91
2-(F)-d)	ITP as substrate for squid myosin	95
2-(F)-e)	Effect of urea on the Ca sensitivity of squid myosin	97
2-(F)-f)	Effect of tryptic digestion of squid myosin on Ca sensitivity in its ATPase	102
2-(F)-g)	Ca sensitivity of scallop myosin alone and loss of it by EDTA-treatment	107
2-(F)-h)	Effects of desensitization and resensi- tization on superprecipitation ability of acto-scallop myosin	111

2-(G)	ATP concentration dependence of ATPase and superprecipitation	113
2-(G)-a)	ATP concentration dependence of super- precipitation and ATPase of squid and scallop myosin Bs	113
2-(G)-b)	ATP concentration dependence of Mg-ATPase activity of squid and scallop myosin	116
2-(G)-c)	ATP concentration dependence of Mg-ATPase activity of desensitized and resensitized scallop myosin and of actomyosin from them	116
2-(H)	Tension development of glycerinated muscle fiber of scallop adductor muscle	119
2-(H)-a)	Effect of EDTA-treatment of glycerinated fibers on its Ca sensitivity in tension development	119
2-(H)-b)	Relationship between EDTA-LC content in fibers and Ca sensitivity of tension development in the presence of calcium	122
2-(H)-c)	Effect of regulatory light chain on Ca sensitivity and on tension development in the presence of calcium	124
IV	Discussion	127
V	Summary	136
VI	Acknowledgements	142
VII	References	143

I INTRODUCTION

It is well known that two different control systems regulate the activity of various muscles(1,2). In vertebrate muscles, troponin and tropomyosin are apparently the only regulatory proteins and the control is therefore actin-linked(3). However, the mechanism of regulation in molluscan muscles differs from that in vertebrate muscles. In 1970, Kendrick-Jones et al.(2) found myosin-linked Ca regulation in scallop muscle, and also found that one of the light chain components of scallop myosin acts as regulatory subunit(4,5).

In both types of regulation, contraction is triggered by small amounts of calcium ions. The resting state is maintained because actin and myosin are unable to interact in the absence of calcium, and this occurs by the blocking the sites either on actin or on myosin. Despite this similarity in function, the interaction between actin and myosin is prevented differently in the two regulatory systems, and two systems contained different components. These two types of component(troponin in actin-linked and regulatory light chain in myosin-linked) cannot related to each other in any simple fashion, and it is very unlikely that one regulatory system could have evolved directly from the other.

A comparative study may give insights into the way of the two regulatory mechanisms evolved and also certain

functional differences between various muscles. Lehman and Szent-Györgyi(6) extensively investigated the distribution of regulatory systems, and categorized vertebrates muscles as having a group of actin-linked regulation system, and molluscan muscles as having a myosin-linked regulation system.

In the present report, a study is described on the Ca regulation system in the mantle muscle of squid and in the adductor muscle of scallop. The author was able to isolate not only Ca sensitive squid myosin but also squid troponin as well as native tropomyosin in a pure form. It can be, therefore, concluded that squid mantle muscle has both a myosin-linked regulatory system and a actin-linked regulatory system, although myosin-linked regulation appears to be predominant in preparation of squid myosin B.

A further study is conducted on the isolation of regulatory subunit in squid myosin(regulatory light chain) and on its characterization. In the process of functional characterization of squid regulatory light chain, the author found some important differences from those described by Kendrick-Jones et al.(5) in the functional role of regulatory light chain.

II Methods and Materials

1 Preparative methods for muscular proteins

1-(A) Myosin B and myosin from squid mantle muscle

Preparative methods for squid myosin B and for myosin were outlined in Chart I. The mantle muscle was quickly excised from a live squid, blended with 0.16 M KCl, 20 mM Tris-HCl(pH 7.5), 0.1 mM PMSF and 10 mM 2-mercaptoethanol and centrifuged to remove the supernatant. The muscle residue was washed several times with the same solution, then extracted with 3 vols of Weber-Edsall solution containing 1 mM ATP, 0.1 mM PMSF, and 10 mM 2-mercaptoethanol at 0°C for 24 h. The extract was obtained by centrifugation, and diluted with 4 vols of 1 mM NaHCO₃. The precipitate thus formed was collected by centrifugation, redissolved in 0.2 M MgCl₂, 5 mM ATP, 1 mM EGTA, 10 mM 2-mercaptoethanol, and 10 mM phosphate buffer(pH 6.4), and centrifuged at 10,000 x g for 30 min to remove insoluble materials. ATP(5 mM) was added to the clear supernatant, and the mixture was strongly centrifuged at 80,000 x g for 90 min, thus removing actin precipitates. The supernatant was diluted with 13 vols of 1 mM NaHCO₃. The myosin precipitate was collected by centrifugation, redissolved in 0.25 M KCl, 10 mM Tris-maleate(pH 7.0), and 10 mM 2-mercaptoethanol, and centrifuged at 100,000 x g for 60 min, removing insoluble materials. The supernatant was diluted with 4 vols of 1 mM NaHCO₃, and 10 mM

Chart I. Preparative method of myosin from squid muscle

Squid mantle muscle

- mince at 0 - 4°C
- wash with 0.16 M KCl, 0.1 mM PMSF, pH 7.5
- extract with Weber-Edsall solution containing
1 mM ATP, 0.1 mM PMSF, 0.1 mM DTT for 2 h
at 0 - 4°C
- centrifuge at 14,000 x g for 20 min

Supernatant(myosin B)

- dilute KCl to 50 mM by adding 1 mM NaHCO₃
- centrifuge at 10,000 x g for 10 min

Precipitate

- suspend in 0.2 M MgCl₂, 1 mM EGTA, 10 mM sodium
phosphate, pH 6.4
- add 10 mM ATP
- centrifuge at 80,000 x g for 90 min

Supernatant

- dilute MgCl₂ to 15 mM by adding 1 mM NaHCO₃
- centrifuge at 10,000 x g for 10 min

Precipitate

- dissolve into 0.25 M KCl, 0.1 mM DTT, pH 7.5
- centrifuge at 80,000 x g for 60 min

Supernatant

- dilute KCl to 50 mM by adding 1 mM NaHCO₃
- centrifuge at 14,000 x g for 10 min

Supernatant

- add 10 mM MgCl₂
- keep at 0 - 4°C for 30 min
- centrifuge at 14,000 x g for 10 min

Precipitate

- dissolve into 0.5 M KCl, 0.1 mM DTT, pH 7.5
- dialyze against 0.5 M KCl, 0.1 mM DTT, 20 mM
Tris-maleate, pH 7.5
- centrifuge at 100,000 x g for 60 min

Supernatant(myosin)

MgCl₂ was then added(7). The myosin precipitate was collected by centrifugation, dissolved in 0.5 M KCl, 20 mM Tris-maleate(pH 7.5), and 0.2 mM DTT, and stored at 0°C.

1-(B) Actin from squid mantle muscle

Actin was prepared from the mantle muscle of squid by the method of Spudich and Watt(8). Crude G-actin was extracted from the acetone-dried muscle powder with 20 vols of 0.5 mM ATP, 0.5 mM 2-mercaptoethanol, 0.2 mM CaCl₂, and 2 mM Tris-HCl(pH 7.6) for 30 min at 0°C. G-actin in the extract was polymerized by adding 0.6 M KCl, 2 mM MgCl₂, and 10 mM Tris-HCl(pH 7.6) and leaving the mixture to stand for 1 h at room temperature(25°C). F-actin was collected by ultracentrifugation at 80,000 x g for 180 min. The F-actin pellet was then dialyzed against the extraction solution, thus depolymerizing F-actin into G-actin. The G-actin solution was clarified by centrifugation at 100,000 x g for 120 min, and the clear supernatant of G-actin was repolymerized into F-actin.

1-(C) Native tropomyosin from squid mantle muscle

Squid native tropomyosin was prepared from mantle muscle by the method of Ebashi(3). The squid mantle muscle was blended with 0.16 M KCl, 20 mM Tris-HCl(pH 7.5) containing 0.1 mM PMSF and 10 mM 2-mercaptoethanol, and

centrifuged to obtain muscle residue. This washing procedure was repeated several times. The washed residue was extracted with 10 mM 2-mercaptoethanol and 0.1 mM PMSF for a day at 0°C. The viscous supernatant of native tropomyosin was obtained by centrifugation, and fractionated by ammonium sulfate salting-out. The fraction precipitated between 45 - 65 % saturation was collected by centrifugation, dissolved in a small volume of 1 mM NaHCO₃, and dialyzed against the same solution. The dialyzed solution of native tropomyosin was then clarified by centrifugation at 100,000 x g for 60 min.

1-(D) Troponin and tropomyosin from squid mantle muscle

Squid native tropomyosin was separated into troponin and tropomyosin by the method of Ebashi(9). The squid native tropomyosin solution in 0.4 M LiCl was brought to pH 4.7 with 0.5 N HCl at 0°C, and then centrifuged. The supernatant containing troponin was adjusted back to pH 7.5. For further purification, the supernatant was chromatographed on a column of DEAE-cellulose. Troponin was eluted with KCl between 0.16 M and 0.18 M. Squid tropomyosin was prepared from the precipitate by placing native tropomyosin in 0.4 M LiCl at pH 4.7. It was purified by repeating the isoelectric precipitation at pH 4.7 in 0.4 M LiCl, and by precipitation with

ammonium sulfate between 50 and 55 % saturation.

1-(E) Myosin, desensitized myosin and EDTA-LC from
scallop adductor muscle

Scallop myosin(myosin B or myofibrils) was prepared from scallop adductor striated muscle by the method used for squid myosin preparation(10, also see 2). Scallop desensitized myosin was prepared by the method of Szent-Györgyi et al.(4) with a slight modification. Scallop myosin was kept suspended for 15 min in an ice-cooled solution of 10 mM EDTA, 30 mM KCl, and 10 mM Tris-HCl(pH 7.5). The suspension was then centrifuged at 6,000 x g for 10 min, and the supernatant and the sediment were separately obtained. The EDTA-treatment procedure was repeated several times. The pellet of desensitized myosin was dissolved in 0.5 M KCl, 20 mM Tris-maleate(pH 7.5) and 1 mM DTT. The supernatants were combined and used to isolate scallop EDTA-light chain by ammonium sulfate salting-out. The fraction precipitating between 50 and 100 % saturation with ammonium sulfate was collected.

1-(F) Light chains from squid mantle muscle

Two types of light chain subunits of squid myosin were dissociated from large polypeptide chains(heavy chains) by the method of Holt and Lowey(11). Preparative method of squid light chains was outlined in Chart II.

Chart II. Preparative method for squid light chain

Squid myosin

- 4 M urea, 0.25 M KCl, 2.5 mM EDTA, 2.5 mM DTT,
and 25 mM Tris-HCl, pH 8.0 for 30 min at 25°C
- dilution to 50 mM KCl
- centrifugation

Supernatant

DEAE-cellulose

- 20 mM Tris-HCl, pH 8.0, 0.1 mM DTT
- elution with 1 M KCl
- dialysis against 0.15 M KCl, 20 mM Tris-HCl, pH 8.0,
0.1 mM DTT
- centrifugation

Supernatant

- ammonium sulfate salting-out

Precipitate

- between 50 and 100 % saturation
- dialysis against 0.15 M KCl, 20 mM Tris-HCl, pH 8.0,
0.1 mM DTT

Dialyzate

Sephadex G-75(2.5 x 90 cm)

- 0.15 M KCl, 20 mM Tris-HCl, pH 8.0, 0.1 mM DTT

light chains fraction

DEAE-Sephadex A25(1 x 80 cm)

- 0.15 - 0.3 M KCl, 20 mM Tris-HCl, pH 8.0, 0.5 mM DTT

LC-2(→ 0.23 M KCl)

LC-1(→ 0.25 M KCl)

Squid myosin(10 - 20 mg/ml) was kept at 25°C for 30 min in 4 M urea, 0.25 M KCl, 25 mM Tris-HCl(pH 8.0), and 20 mM 2-mercaptoethanol. The heavy chains precipitated on dilution with 10 vols of cold water, and the light chains were collected in the supernatant on centrifugation. In order to concentrate the light chains, the supernatant was applied to a DEAE-cellulose column, removing the unretained fraction, and the column was eluted with a small volumes of 1 M KCl containing 20 mM Tris-HCl(pH 8.0) and 10 mM 2-mercaptoethanol. The concentrated light chain solution thus obtained was dialyzed against 0.15 M KCl, 20 mM Tris-HCl(pH 8.0), and 10 mM 2-mercaptoethanol, and insoluble materials formed were removed by centrifugation at 20,000 x g for 30 min. The supernatant containing light chains was then fractionated by ammonium sulfate salting-out. The fraction precipitating between 50 and 100 % saturation with ammonium sulfate was collected. This fraction was fairly free from paramyosin, actin, tropomyosin, and troponin. If necessary, especially when it was obtained from myosin B or myofibrils, the light chain fraction was further purified by Sephadex G-75 gel filtration in 0.15 M KCl and 20 mM Tris-HCl(pH 8.0). In the gel filtration, paramyosin, actin, and troponin-tropomyosin complex were eluted at the void volume, and light chains were retained, thus being separated. Light chains thus obtained in a pure form were chromatographed

on a DEAE-Sephadex A25 column(12) to obtain the two types of light chains separately. A long column of 1 x 80 cm, a shallow gradient of KCl concentration(from 0.15 M to 0.3 M) and a slow rate of flow(10 ml/h) were used to facilitate the separation.

1-(G) Myosin and actin from rabbit skeletal muscle

Skeletal myosin was prepared from rabbit back and leg muscle by the method of Connell(12) with a slight modification. Rabbit skeletal actin was prepared by the method of Spudich and Watt(8).

1-(H) Glycerinated muscle fibers from scallop adductor muscle

Glycerinated muscle fiber bundles of scallop adductor muscle were prepared by routine method of Szent-Györgyi et al.(14). Strips of scallop adductor muscle, approximately 2 mm x 2 mm in section and 2 - 3 cm in length, were dissected out, tied with cotton threads onto glass rods at the appropriate length, and immersed in a solution containing 50 % glycerol, 5 mM phosphate buffer(pH 7.0), 4 mM $MgCl_2$, 2 mM EGTA and 1 mM 2-mercaptoethanol at 0°C. After overnight incubation at 0°C with exchange of the solution, the preparation immersed in the new solution were stored at - 20°C for at least a month before use.

2 Assay method for biological and physico-chemical properties

2-(A) Assay for ATPase activity

The actin-activated Mg-ATPase activity of myosin or that of myosin alone were assayed at 25°C by measuring Pi liberation. Phosphate was determined by the method of Fiske and Subbarow(15), except that p-methylaminophenol sulfate was used in place of 1, 2, 4-aminonaphtol sulphonic acid as a reducing reagent. In the ATP concentration dependence, an ATP regenerating system of pyruvate kinase and phosphoenol pyruvate(1 mM) was used to maintain the ATP concentration, and ATPase activity was determined by the measuring liberated pyruvate(16). The standard reaction medium contained 20 mM Tris-maleate(pH 6.8), 30 mM KCl, 2 mM MgCl₂, 1 mM ATP, and 0.1 mM CaCl₂ or 0.5 mM EGTA. Ca sensitivity was defined as follow:

$$\text{Ca sensitivity} = \frac{A_{\text{Ca}} - A_{\text{E}}}{A_{\text{Ca}}} \times 100, \text{ where } A_{\text{Ca}} \text{ and } A_{\text{E}}$$

were Mg-ATPase activity in the presence of 0.1 mM CaCl₂ and 0.5 mM EGTA, respectively. The desired concentration of the free calcium ions was obtained by addition of 0.2 mM CaCl₂ and appropriate concentrations of EGTA. Desired concentrations of free strontium ions were obtained by adding 0.5 mM EGTA and appropriate concentrations of SrCl₂. The stability constants adopted for estimation of the free ion concentrations were $5 \times 10^5 \text{ M}^{-1}$ for the

Ca-EGTA complex and $6 \times 10^2 \text{ M}^{-1}$ for the Sr-EGTA complex(17).

2-(B) Measurement of superprecipitation

Superprecipitation of actomyosin was induced at 20°C by adding 0.5 mM ATP to reaction medium containing 30 mM KCl, 20 mM Tris-maleate(pH 6.8), 2 mM MgCl_2 , and 0.1 mM CaCl_2 or 0.5 mM EGTA. The absorbance at 550 nm was measured with a Shimadzu UV 210A spectrophotometer.

2-(C) SDS gel, urea gel and disc gel electrophoreses and densitometry

SDS gel electrophoresis was carried out as described by Weber and Osborn(18). Urea gel electrophoresis was conducted essentially as described by Perrie et al(12). Gels contained 7.5 % polyacrylamide, 8 M urea, and 0.375 M Tris-HCl(pH 8.9). The sample protein was kept at 37°C for 1 h in a solution of 8 M urea, 0.1 M Tris-HCl(pH 8.9), 5 % 2-mercaptoethanol, and Bromophenol Blue as a marker. The running buffer contained 0.05 M Tris and 0.19 M glycine(pH 9.4). Disc gel electrophoresis was conducted as described by Davis(19). Gels contained 7.5 % polyacrylamide, 0.375 M Tris-HCl(pH 8.9), 10 % 2-mercaptoethanol, and 1 mM CaCl_2 (+Ca) or 1 mM EGTA(-Ca). The running buffer was the same as that used for urea gel electrophoresis. Gels were stained with 0.25 % Coomassie brilliant blue or 0.5 % Fast Green dissolved in 50 %

methanol and 9.2 % acetic acid(26). Destaining was performed with 50 % methanol and 9.2 % acetic acid. The density of Fast Green stained protein bands was measured at 640 nm with Shimadzu CS-900 dual-wavelength TLC scanner.

2-(D) UV absorption spectra of light chains and divalent cation induced difference spectra of them

The ultraviolet absorption spectra of myosin light chains and the calcium- or strontium-induced change in the UV absorption of light chains were determined at 20°C in a medium containing 0.1 M KCl, 20 mM histidine(pH 6.8), using a Shimadzu 210 A spectrophotometer. In obtaining UV difference spectra, the reference cell contained 1 mM EGTA and the sample cell contained Ca-EGTA buffer or Sr-EGTA buffer(21).

2-(E) Calcium binding of myosin and light chain

Calcium bindings to myosin and to regulatory light chains were measured by an equilibrium dialysis method, using ^{45}Ca (21). The equilibrium dialysis was conducted at 20°C in 0.1 M KCl and 20 mM histidine buffer(pH 6.8), or in 0.3 M KCl and 20 mM histidine buffer(pH 6.8). The free calcium concentration was set by using Ca-EGTA buffer (17).

2-(F) Quantative determination of protein concentration

Protein concentrations were determined by the biuret method, using bovine serum albumine fraction V as a standard(22).

2-(G) Tension determination of glycerinated muscle fibers of scallop muscle

The isometric tension of muscle fibers bundle were determined as described by Simmons and Szent-Györgyi(23). The glycerinated muscle strip was shredded into a muscle bundle with farceps with thin and well fitted tips. The muscle bundle was immediately applied to tensiometer. One end of the fiber bundle was fixed to an immobilized glass bar and other to transducer(UL-2, Shinkoh Tsushin) which was connected to a amplifier-recorder system. Thus attached fibers were rinsed in a solution containing 0.1 M KCl, 20 mM imidazole(pH 6.8), 5 mM $MgCl_2$, 2 mM EGTA, and 0.3 mM DTT.

The two types of solutions were used in this study: a contracting solution(0.1 M KCl, 20 mM imidazole(pH 6.8), 5 mM $MgCl_2$, 4 mM ATP, 0.2 mM $CaCl_2$, 0.1 mM EGTA and 0.3 mM DTT) and a relaxing solution(0.1 M KCl, 20 mM imidazole(pH 6.8), 5 mM $MgCl_2$, 4 mM ATP, 1 mM EGTA, and 0.3 mM DTT). Contraction was induced by the changing the solution, in which fibers were immersed, from relaxing solution to contracting solution. The temperature of

the cell in which the fibers set in was held at 20°C by circulating water under the cell.

For the EDTA-treatment of fiber bundle attached to tensiometer, fibers were soaked in a solution containing 0.1 M KCl, 20 mM Tris-HCl(pH 7.5), 10 mM EDTA and 0.3 mM DTT at 20°C for various intervals. In order to resensitize the muscle fibers, thus treated with EDTA, were soaked in relaxing solution containing approximately 1.8 mg/ml of scallop EDTA-LC or squid LC-2 for various intervals.

3 Materials and reagents

The abbreviations used were as follows:

ATP(adenosine 5' triphosphate)

ITP(inosine 5' triphosphate)

EDTA(ethylenediamine tetraacetate)

EGTA(ethyleneglycol bis(β -aminoethyl)-N,N' tetraacetate)

DTT(dithiothreitol)

DTNB(5, 5' dithiobis(2-nitrobenzoate))

SDS(sodium dodecyl sulfate)

PMSF(phenylmethylsulfonyl fluoride)

DEAE-cellulose(diethylaminoethyl-cellulose)

UV(ultraviolet)

OD(optical density)

LC(light chain component of myosin)

HC(heavy chain component of myosin)

NTM(native tropomyosin)

TN(troponin)

TM(tropomyosin)

The animals used were as follows: rabbit(Oryctolagus cuniculus domesticus), squid (Todarodes pacificus), and scallop(Patinopecten yessoensis).

The reagents purchased were as follows:

ATP(Daiichi pure chemicals Co. Ltd.)

SDS(Nakarai chemical Ltd.)

EDTA, EGTA(Dojin chemical Co. Ltd.)

DEAE-cellulose(Brown Co.)

Sephadex G-200, Sephadex G-75, DEAE-Sephadex A-25

(Pharmacia fine chemicals)

ITP, PMSF, PEP trypsin and trypsin inhibitor(Sigma chemicals Co.)

DTT, 2-mercaptoethanol, urea, ammonium sulfate, Fast Green, acrylamide(Wako pure chemical industry Ltd.)

⁴⁵Ca(Japan isotope center)

III Results

A competitive test devised by Lehman et al(6, 24) was applied to the squid myosin B to determine whether the Ca regulatory system is myosin-linked or actin-linked. As shown in Table I, the ATPase activity of the squid myosin B preparation($\mu\text{mol Pi/min/mg}$ myosin B) was 0.559 in the presence of 0.1 mM $\text{CaCl}_2(+\text{Ca})$, and 0.003 in the presence of 0.5 mM EGTA($-\text{Ca}$). The two activities were unaffected by the addition of a large amount of rabbit skeletal F-actin(34.6 μg actin to 44.6 μg of squid myosin B). The Ca sensitivity defined as described in "Methods and materials" was unaffected by the addition of a large amount of skeletal myosin(40.6 μg myosin to 44.6 μg of squid myosin B). It thus appeared that both myosin-linked and actin-linked Ca regulatory systems are present in squid myosin B. This view conflicts with that of Lehman and Szent-Györgyi(6), who categorized squid muscle in a group of muscle having only myosin-linked Ca regulation on the basis of the same competitive test. Tsuchiya et al.(25) also suggested the presence of actin-linked regulation in squid muscle on the basis of the same test.

Table I Effect of skeletal F-actin and skeletal myosin on the Mg-ATPase of squid myosin B.

Mg-ATPase was assayed at 25°C in a medium containing 30 mM KCl, 20 mM Tris-maleate(pH 6.8), 2 mM MgCl₂, 1 mM ATP, and 0.1 mM CaCl₂(+ Ca) or 0.5 mM EGTA(-Ca), and 89 µg/ml of squid myosin B(A), or 44 µg/ml of squid myosin B and 35 µg/ml of skeletal F-actin(B), or 44µg/ml of squid myosin B and 41 µg/ml of skeletal myosin(C). In (B) and (C), F-actin and myosin were mixed with myosin B in 0.5 M KCl, 20 mM Tris-maleate(pH 7.5) before making up the assay medium.

	Mg-ATPase(µmol Pi/min.mgmyosinB)		
	+ Ca	- Ca	Ca.sens.
(A) <u>squid myosin B</u>	<u>.559</u>	<u>.003</u>	<u>99</u>
(B) + rabbit actin	.559	.038	93
(C) <u>+ rabbit myosin</u>	<u>1.053</u>	<u>.292</u>	<u>73</u>

1 Actin-linked Ca regulation

1-(A) Native tropomyosin in squid muscle

1-(A)-a) Ca sensitivity of squid acto-skeletal myosin

The competitive test shown in Table I suggested that squid myosin B contains a actin-linked regulatory system as well as a myosin-linked system. Goldberg and Lehman(26) also recently reported the presence of actin-linked Ca regulatory complex. The author, therefore, investigated the actin-linked Ca regulation in squid muscle. Three different squid actin preparations were prepared for this purpose. They were: (A) pure F-actin obtained by the method of Spudich and Watt, (C) crude F-actin i.e., the whole extract from squid acetone-dried muscle powder polymerized by 0.1,M KCl,(pH 7.6) and 1 mM $MgCl_2$, and (B) partially purified F-actin, precipitated by ultracentrifugation of (B). Each of three preparations of squid actin was combined with rabbit skeletal myosin, which is Ca insensitive. Actomyosin thus reconstituted by using the F-actin(A) showed no Ca sensitivity in Mg-ATPase assay(Fig. 1A), whereas that reconstituted by using F-actin(C) showed definite Ca sensitivity of 74 %(Fig. 1C). That obtained using F-actin(B) also showed Ca sensitivity of 50 %(Fig. 1B).

SDS gel electrophoretic patterns of the three actin preparations are shown in Fig. 2. F-actin(A) was free from other proteins, crude F-actin(C) contained many





Fig. 1. Ca sensitivity of squid acto-skeletal myosin

Actomyosins were reconstituted from rabbit skeletal myosin (82 $\mu\text{g/ml}$) and squid actin preparation of three different grades of purity; (A), actin (85 $\mu\text{g/ml}$) purified by the method of Spudich and Watt; (C), crude actin (106 $\mu\text{g/ml}$) comprising whole extracts from acetone-dried squid muscle powder with 2 mM Tris-HCl (pH 7.6), 0.5 mM ATP, 0.5 mM 2-mercaptoethanol, and 0.2 mM CaCl_2 , polymerized with 0.1 M KCl, 1 mM MgCl_2 ; (B), partially purified actin (98 $\mu\text{g/ml}$) precipitated by ultracentrifugation of (C). Mg-ATPase was assayed in the medium described in Table I.



(A) (B) (C)

Fig. 2. SDS gel electrophoretic patterns of squid actin preparations

A, B, and C were the actin preparations used in the experiments shown in Fig. 1A, B, and C, respectively. Approximately 10 μ g(A) or 20 μ g(B and C) of protein was loaded on each gel.

other proteins, and F-actin(B) showed 5 protein bands including one actin band. These result revealed that four protein components were capable of associating with F-actin, thus precipitating with F-actin on ultracentrifugation. Accordingly, it is suggested that the four protein components may correspond on the constituents of squid native tropomyosin.

1-(A)-b) Isolation of native tropomyosin from squid muscle

Native tropomyosin was isolated from the squid mantle muscle. The method for preparing rabbit skeletal native tropomyosin was successfully employed to the squid muscle, though 10 mM 2-mercaptoethanol and 0.1 mM PMSF were present in the extracting medium, and the fraction precipitated between 45 and 65 % saturation of ammonium sulfate. The SDS gel electrophoretic pattern of native tropomyosin thus isolated is shown later(Fig. 7). Four major bands with a few minor bands were detectable.

Squid native tropomyosin thus isolated was tested for its effect on the superprecipitation of rabbit skeletal actomyosin. As shown in Fig. 3A, skeletal actomyosin alone showed no Ca sensitivity. The time required to reach half-maximal turbidity(superprecipitation) ($\tau_{1/2}$) was 3.5 min in 0.1 mM CaCl_2 (●) as well as in 0.5 mM EGTA (○). On the other hand, in the presence of squid native tropomyosin(55 μg), $\tau_{1/2}$ for the superprecipitation




Fig. 3. Effect of squid native tropomyosin on the superprecipitation of reconstituted actomyosin

Actomyosin was reconstituted from skeletal myosin (128 $\mu\text{g/ml}$) and skeletal actin (57 $\mu\text{g/ml}$) (A), or skeletal myosin (128 $\mu\text{g/ml}$) and squid actin (46 $\mu\text{g/ml}$) (B). ($\bullet$ $\circ$), actomyosin alone, ($\blacktriangle$ $\triangle$), squid native tropomyosin (55 $\mu\text{g/ml}$) was added. The reaction was carried out at 20°C in a medium containing 30 mM KCl, 20 mM Tris-maleate (pH 6.8), 2 mM MgCl_2 , 0.5 mM ATP, and 0.1 mM CaCl_2 (closed) or 0.5 mM EGTA (open).

of skeletal actomyosin(128 μ g myosin plus 57 μ g actin) was 5.5 min in 0.1 mM CaCl_2 ($\blacktriangle$) and 9 min in 0.5 mM EGTA($\triangle$), thus showing Ca sensitivity. Purified squid actin was capable of replacing skeletal actin in the above test, that is, hybrid actomyosin(squid actin/skeletal myosin) alone showed no Ca sensitivity, τ 1/2 being 4.5 min ($\bullet$ $\circ$). However, in the presence of squid native tropomyosin(55 μ g), the hybrid(128 μ g myosin plus 46 μ g actin) showed Ca sensitivity, τ 1/2 being 8 min in 0.1 mM CaCl_2 ($\blacktriangle$) and 24 min in 0.5 mM EGTA($\triangle$). It is, therefore, concluded that our preparation of squid native tropomyosin has the same Ca sensitizing action as skeletal native tropomyosin.

1-(A)-c) A comparison between native tropomyosins from rabbit and squid muscles

The biological activities of squid native tropomyosin were compared with those of rabbit one by measuring the inhibitory effect of them on skeletal actomyosin Mg-ATPase activity. As shown in Fig. 4, skeletal actomyosin alone was not sensitive to calcium. Addition of rabbit native tropomyosin conferred Ca sensitivity to this actomyosin as previously reported(27), and squid native tropomyosin also conferred Ca sensitivity to actomyosin. Furthermore, squid native tropomyosin was distinguishable from that of rabbit in the following respect that Mg-ATPase activity




Fig. 4. Effects of native tropomyosins from squid and rabbit on the Mg-ATPase activity of actomyosin


The ATPase activity of skeletal actomyosin was assayed in the medium containing 30 mM KCl, 20 mM Tris-maleate(pH 6.8), 2 mM MgCl₂, 1 mM ATP and either 0.1 mM CaCl₂(closed) or 0.5 mM EGTA(open). (■ □), actomyosin alone, (● ○), squid native tropomyosin, and (▲ △), rabbit native tropomyosin.

of actomyosin was inhibited by squid native tropomyosin not only in the absence of calcium but also in the presence of calcium. It was characteristic to squid native tropomyosin and was not found in rabbit native tropomyosin (Fig. 4).

1-(A)-d) Binding of squid native tropomyosin to
rabbit F-actin

Binding ability of squid native tropomyosin to rabbit F-actin was measured by centrifugation method. Squid native tropomyosin was mixed with F-actin in a medium of 0.1 M KCl, 20 mM Tris-maleate(pH 7.5) and maintained at 0°C for an hour. The mixture was then centrifuged down at 100,000 x g for three hours, and thus obtained F-actin pellet was dissolved in SDS solution and applied to SDS gel electrophoresis. The results obtained were shown in Fig. 5. Beside actin band, four major protein bands were found to be identical to those of native tropomyosin. This result indicates that squid native tropomyosin is capable of binding to F-actin. The stoichiometric binding of native tropomyosin to F-actin was further studied. As shown in Fig. 6, binding of native tropomyosin from squid or rabbit to F-actin was reached maximum at around 30 % addition(w/w) of native tropomyosin to F-actin. Amounts of subunits of native tropomyosin bound to

F-actin were estimated from the stain density in SDS gel electrophoretic patterns. Table II showed that the amounts of 28,000 dalton and 24,000 dalton components in squid native tropomyosin was found to be very small in the pellet precipitating with F-actin. Tropomyosin and 52,000 dalton component were found in the pellet in an amount of about 1 mol per 7 mol of actin monomer. In addition, it was found that intact myofibrils from squid mantle muscle contained small amounts of 28,000 dalton and 24,000 dalton components. This fact suggested that the selective loss of these components did not occur during preparation of squid native tropomyosin. These results, therefore, strongly indicated that squid native tropomyosin was composed of different(or irregular) subunit composition from that of rabbit native tropomyosin.



(A) (B) (C)

Fig. 5. Binding of squid native tropomyosin to F-actin

Squid native tropomyosin was mixed with rabbit F-actin in a weight ratio of 0.3 : 1 for native tropomyosin : F-actin in the medium of 0.1 M KCl, 20 mM Tris-maleate (pH 7.5), and maintained for an hour in ice-cold water. The mixture was then ultracentrifuged at 100,000 x g for 3 hr, the pellet thus obtained was applied on SDS gel rod. (C), squid native tropomyosin, (B), pellet obtained after ultracentrifugation, and (A), rabbit F-actin alone.




Fig. 6. Stoichiometric binding of native tropomyosin to F-actin

Binding of native tropomyosin to F-actin was measured by the same method as in Fig. 5, except that various amounts of native tropomyosin were mixed. F-actin pellet combining native tropomyosin was separated and applied on SDS gel and stained with Fast Green. The intensity of each subunit was determined. (A) squid native tropomyosin; (Δ), total, ($\bullet$), tropomyosin, ($\square$), 52,000 dalton component, ($\circ$), 28,000 dalton component and ($\blacktriangle$), 24,000 dalton component, (B) rabbit native tropomyosin, (Δ), total, ($\bullet$), tropomyosin, ($\square$), troponin-T, ($\circ$), troponin-I, and ($\blacktriangle$), troponin-C.

Table II. Amounts of subunits in squid native tropomyosin bound to skeletal F-actin

The amounts of subunits bound to F-actin were estimated from the data in Fig. 6.

	SUBUNIT	moles/7moles of ACTIN
Rabbit	TM	1.0
	TN-T	1.4
	-I	1.4
	-C	1.0
Squid	TM	1.0
	TN-52,000	1.0
	-28,000	0.3
	-24,000	0.3

1-(B) Troponin and tropomyosin in squid muscle

1-(B)-a) Isolation of troponin and tropomyosin from squid muscle

Ebashi's method for the preparation of skeletal troponin was used with a slight modification to isolate squid troponin from squid native tropomyosin. Squid native tropomyosin was dissolved in 0.4 M LiCl, and the pH of the solution was adjusted to 4.7 by adding 0.5 N HCl. The supernatant(crude preparation of squid troponin) was collected by centrifugation. The crude preparation was further purified by DEAE-cellulose chromatography. The elution pattern is shown in Fig. 7. A major peak was eluted with KCl at a concentration around 0.17 M. This major peak was identified as squid troponin, on the bases of its SDS gel electrophoretic pattern(Fig. 8) and its Ca sensitizing activity. Squid tropomyosin was also isolated from native tropomyosin.

1-(B)-b) SDS gel electrophoretic patterns of squid troponin and tropomyosin

SDS gel electrophoretic patterns of purified troponin, tropomyosin, and original native tropomyosin thus obtained from squid muscle are shown in Fig. 8. Purified squid troponin was identical with skeletal troponin in showing three protein bands, though squid troponin was different in mobility from rabbit skeletal troponin(28 - 30) or from carp dorsal troponin(27). It is rather similar to lobster




Fig. 7. DEAE-cellulose chromatography of squid troponin.

A solution of native tropomyosin in 0.4 M LiCl was brought to pH 4.7. Tropomyosin precipitated was removed by centrifugation. The supernatant was readjusted to pH 7.5, dialyzed against 10 mM Tris-HCl(pH 7.5), and then applied(25.5 mg) to a column(0.8 x 15 cm) of DEAE-cellulose pre-equilibrated with the same buffer. The column was eluted with a linear gradient of KCl concentration. The flow rate was approximately 6 ml/h, and fractions of 3.8 ml were collected.

NTM TN TM




Fig. 8. SDS gel electrophoretic patterns of native tropomyosin, troponin and tropomyosin prepared from squid muscle

NTM, native tropomyosin(20 μ g); TN, troponin(15 μ g); TM, tropomyosin(5 μ g).




Fig. 9. Elution profile of squid troponin from Sephadex G-200 gel column

Squid troponin in 0.1 M KCl, 20 mM Tris-HCl(pH 7.5) was applied to a Sephadex G-200 gel column pre-equilibrated with the same buffer. About 20 mg of troponin was applied and 4.8 ml/tube was collected. Void volume and bed volume were determined by using Blue dextran and ATP, respectively.

troponin(31); the three bands of squid troponin corresponded to 52,000, 28,000, and 24,000 daltons.

It seems necessary to study whether three subunits detected in SDS gel of squid troponin form a complex or not. Squid troponin was dissolved in 0.1 M KCl, 20 mM Tris-HCl(pH 7.5) and applied to a Sephadex G-200 gel column pre-equilibrated with the same buffer solution. The elution profile was shown in Fig. 9. The most of protein applied was eluted as a single fraction after the void volume and molecular weight was estimated to be 1×10^5 daltons. These results suggested that three subunits in squid troponin formed a complex under the above conditions.

1-(B)-c) Effect of squid troponin, tropomyosin and their complex on superprecipitation of skeletal actomyosin

The Ca sensitizing activity of squid troponin was examined by studying its effect on the superprecipitation of actomyosin in the presence of tropomyosin. Actomyosin was reconstituted from rabbit skeletal myosin(157 μ g) and squid purified actin(38 μ g). As shown in Fig. 10A, no Ca sensitivity was detectable with actomyosin alone: $\tau_{1/2}$ was 4 min in the presence of 0.1 mM CaCl_2 (●) as well as in its absence (○). Addition of squid troponin (51 μ g) alone did not confer Ca sensitivity (▲ △).




Fig. 10. Effect of squid troponin-tropomyosin complex on the superprecipitation of reconstituted actomyosin. (A); Actomyosin reconstituted from rabbit skeletal myosin (157 µg/ml) and squid purified actin (38 µg/ml) with no addition (● ○), or with squid troponin (51 µg/ml) (▲ △), or with squid tropomyosin (47 µg/ml) (■ □). (B); Squid troponin (51 µg/ml)-tropomyosin (47 µg/ml) complex was added to the reconstituted actomyosin (● ○). The reaction medium was as described in Fig. 3. Closed and open symbols denote 0.1 mM CaCl₂ (● ▲ ■) and 0.5 mM EGTA (○ △ □) media, respectively.

Addition of squid tropomyosin(47 μ g) alone also failed to confer Ca sensitivity though it slightly reduced the rate of superprecipitation: $\tau_{1/2}$ was 9 min in the presence of calcium(■) and 11.5 min in its absence(□). On the other hand, Fig. 10B shows that addition of both troponin(51 μ g) and tropomyosin(47 μ g) could confer Ca sensitivity: $\tau_{1/2}$ was 16 min in 0.1 mM CaCl_2 (●) and 31 min in 0.5 mM EGTA(○). It can, therefore, be concluded that the Ca sensitizing effect of squid troponin-tropomyosin complex was similar in general features to that of the skeletal troponin-tropomyosin system, but the Ca sensitizing action of squid native tropomyosin troponin-tropomyosin complex is less effective.

1-(B)-d) A comparison between troponins from rabbit and squid muscles.

The Ca sensitizing effects of hybrid and homologue troponin-tropomyosin complex formed between troponin and tropomyosin from squid and rabbit were then compared. The effect was estimated by measuring the inhibition of actomyosin Mg-ATPase activity with these complexes. Figure 11 showed the effect of complexes using squid troponin and squid tropomyosin or rabbit tropomyosin. Neither of them alone did not confer Ca sensitivity to actomyosin(Fig. 11A). Squid troponin-tropomyosin complex conferred Ca sensitivity but its extent was very small.




Fig. 11. Effects of hybrid and homologue complexes of troponins and tropomyosins from squid and rabbit muscle on the Mg-ATPase of actomyosin

The ATPase was assayed in the same medium as in Fig. 4. (A); (● ○), squid troponin alone, (▲ △), squid tropomyosin alone, and (■ □), rabbit tropomyosin alone. (B); squid troponin plus squid tropomyosin(1/1 by weight ratio). (C); squid troponin plus rabbit tropomyosin(1/1 by weight ratio).




Fig. 12. Effect of hybrid and homologue complexes of troponins and tropomyosins from squid and rabbit muscles. on the Mg-ATPase of actomyosin

The ATPase was assayed in the medium as in Fig. 11.

(A) (● ○) rabbit troponin alone, (▲ △) squid tropomyosin alone, and (■ □) rabbit tropomyosin alone, (B) rabbit tropomyosin plus rabbit troponin (1/1 by weight ratio), (C) squid tropomyosin plus rabbit troponin (1/1 by weight ratio).

It should be noted that Mg-ATPase activity was reduced even when the calcium was present in the system with troponin-tropomyosin complex(Fig. 11B). These effects of complex were similar to that obtained with native tropomyosin from squid mantle muscle. Replacement of squid tropomyosin by rabbit tropomyosin unaffected the general trends(Fig. 11C); the Ca sensitivity was slightly reduced with decreasing ATPase activity in the presence of calcium.

On the contrary, as shown in Fig. 12, the troponin-tropomyosin complex constituted using rabbit troponin well conferred Ca sensitivity to actomyosin. This result indicated that squid tropomyosin was highly functional as well as rabbit tropomyosin. Therefore, the author concluded that the relatively lower Ca sensitivity conferred by squid native tropomyosin was stemmed from properties of squid troponin.

1-(B)-e) Binding of squid troponin and tropomyosin to F-actin

The binding ability of squid troponin, tropomyosin, and their complex to skeletal F-actin was studied. Proteins were mixed in the medium of 0.1 M KCl, 20 mM Tris-HCl(pH 7.5), maintained at 0°C for an hour, centrifuged down at 100,000 x g for 3 h, and then subunit compositions of the pellet thus obtained were analyzed

by SDS gel electrophoresis. Figure 13 showed that squid tropomyosin as well as rabbit tropomyosin bound to skeletal F-actin. Squid troponin alone did not bind to F-actin, while it bound to F-actin only when tropomyosin either from squid or rabbit was present in the system. It is, therefore, clear that squid troponin preparation did not contain any subunit, which was able to bind to F-actin, and that squid troponin initially bind to tropomyosin, forming the complex which was subsequently bound to F-actin. These binding features of troponin and tropomyosin were essentially identical with those of rabbit troponin and tropomyosin system(29).




Fig. 13. Binding of squid troponin and tropomyosin to rabbit F-actin

Binding of squid troponin and tropomyosin to rabbit F-actin was measured by the same method as described in Fig. 5. (A), Squid troponin preparation, (B), squid tropomyosin, (C), rabbit tropomyosin, (D), F-actin pellet with squid troponin and squid tropomyosin, (E), F-actin pellet with squid troponin and rabbit tropomyosin, (F), F-actin pellet with squid troponin, (G), F-actin pellet with squid tropomyosin, and (H), F-actin pellet with rabbit tropomyosin.

2 Myosin-linked Ca regulation

2-(A) Ca sensitive myosin from squid mantle muscle

The presence of myosin-linked Ca regulation in molluscan muscle, especially in scallop muscle was already suggested by Kendrick-Jones et al.(2). In the present study, the presence of myosin-linked Ca regulation system in squid myosin B was also strongly suggested by competitive test of squid myosin B with skeletal F-actin.

2-(A)-a) SDS gel electrophoretic patterns of squid myosin and myosin B

SDS gel electrophoretic patterns of squid myosin B, myosin, and actin used in the present study are shown in Fig. 14. This preparation of myosin was free from actin contamination, though it contained a small amount of paramyosin. Two light chain components with molecular weights of 15,000 and 17,000 were found in this preparation of squid myosin(32, 33). Squid actin was virtually free from any other protein bands, and showed the same mobility as skeletal actin in SDS gel electrophoresis(data not shown)(34).

2-(A)-b) Ca sensitivity of skeletal acto-squid myosin

As shown in Fig. 15, actomyosin reconstituted from squid myosin(202 $\mu\text{g/ml}$) and skeletal purified F-actin

Myo B Myo Act




Fig. 14. SDS gel electrophoretic patterns of myosin B, myosin, and actin prepared from squid mantle muscle

Ten % polyacrylamide gels containing 0.1 % SDS were prepared. Electrophoresis was carried out according to the method of Weber and Osborn. Approximately 60 μ g of myosin B(Myosin B), 50 μ g of myosin(Myosin), and 30 μ g of actin(Act) were applied to each gel rod(10 cm long and 0.5 cm thick).

(175 $\mu\text{g/ml}$) exhibited Ca sensitivity. Superprecipitation of skeletal acto-squid myosin in the presence of 0.1 mM CaCl_2 (●) occurred immediately after the addition of 0.5 mM ATP, but its occurrence in the presence of 0.5 mM EGTA (○) was greatly delayed (over 60 min). The addition of CaCl_2 to the EGTA medium (indicated by an arrow in Fig. 15) induced immediate superprecipitation. Essentially the same result was obtained when skeletal actin was replaced by purified squid F-actin (▲△). This squid F-actin was Ca insensitive, as shown in Fig. 1A.

Ca sensitivity of actomyosin reconstituted from squid myosin (87.0 $\mu\text{g/ml}$) and skeletal F-actin (45.8 $\mu\text{g/ml}$) was also demonstrated by Mg-ATPase assay. Figure 16 shows that P_i liberation proceeds rapidly in the presence of 0.1 mM CaCl_2 (●) but very slowly in its absence (○) and that, unlike actomyosin reconstituted from gizzard myosin and actin (35), P_i liberation in the presence of CaCl_2 was decreased by a subsequent addition of 2 mM EGTA (indicated by the arrow in Fig. 16). It is already known that chicken gizzard actomyosin is phosphorylated in the presence of calcium, so that the rate of liberation by its Mg-ATPase is unaffected by the addition of EGTA to this system, whereas treatment of the same actomyosin with phosphatase reduces the rate of P_i liberation from ATP. These results suggest that phosphorylation and dephosphorylation of light chain component are not




Fig. 15. Superprecipitation of hybrid and squid actomyosins.

Superprecipitation of reconstituted actomyosin was followed by measuring the absorbance at 550 nm. Actomyosin was reconstituted from squid myosin(202 $\mu\text{g}/\text{ml}$) and skeletal actin(107 $\mu\text{g}/\text{ml}$)(● ○) or from squid myosin(202 $\mu\text{g}/\text{ml}$) and squid actin(175 $\mu\text{g}/\text{ml}$)(▲ △). The reaction was carried out at 20°C in a medium containing 30 mM KCl, 20 mM Tris-maleate(pH 6.8), 2 mM MgCl_2 , 0.5 mM ATP, and 0.1 mM CaCl_2 (● ▲) or 0.5 mM EGTA(○ △). 0.5 mM CaCl_2 was added to the EGTA medium as indicated by the arrows.




Fig. 16. Ca dependent Pi liberation by hybrid acto-myosin Mg-ATPase

ATPase activity was assayed in the medium described in Table I, except that skeletal actin(45.8 $\mu\text{g/ml}$) and squid myosin(87.0 $\mu\text{g/ml}$) were used. As indicated by the arrows, 2 mM EGTA was added to the 0.1 mM CaCl_2 medium (●) and 2 mM CaCl_2 to the 0.5 mM EGTA medium(○).

involved in the myosin-linked Ca regulation in squid muscle.

2-(A)-c) EDTA-treatment of squid myosin

Szent-Györgyi et al.(4) reported that treatment of scallop myosin with EDTA in a low-salt medium caused release of one of the light chain components, named EDTA-LC, and resulted in loss of its Ca sensitivity. However, the author found that EDTA-treatment had no effect on the Ca sensitivity of squid myosin. Squid myosin was treated 16 times, each time with 10 vols of 5 mM EDTA, 30 mM KCl, and 10 mM Tris-HCl(pH 7.5). Actomyosin was reconstituted from the EDTA-treated myosin(108 µg/ml) and skeletal actin(60 µg/ml), which was free from troponin and tropomyosin. The hybrid actomyosin thus reconstituted was found to be highly Ca sensitive(Fig. 17), that is, its Mg-ATPase activity(µmol Pi/min/mg myosin) was 0.222 in the presence of 0.1 mM CaCl₂(●), whereas it was 0.02 in the presence of 0.5 mM EGTA(○). These activity values were essentially the same as those obtained with actomyosin reconstituted from untreated squid myosin and actin. No change in the SDS gel electrophoretic pattern of squid myosin was detectable upon EDTA-treatment(data not shown). The result suggesting that squid myosin was not able to desensitize by EDTA-treatment, was also agreed with that reported by Szent-Györgyi et al.(4), using Loligo siphone muscle myosin.




Fig. 17. Effect of EDTA treatment on the Ca sensitivity of squid myosin

Squid myosin was washed 16 times with 10 vols of 5 mM EDTA, 30 mM KCl, and 10 mM Tris-HCl(pH 7.5). Acto-myosin was then reconstituted from EDTA-treated myosin (108 $\mu\text{g/ml}$) and skeletal actin(60 $\mu\text{g/ml}$). Mg-ATPase activity was assayed in the presence of 0.1 mM CaCl_2 (●) or 0.5 mM EGTA(○) in the reaction medium described in Table I.

2-(B) A comparison of Ca and Sr requirements for Mg-ATPase
activity of myosin-linked and actin-linked
regulations

2-(B)-a) Calcium or strontium ion requirement in squid
myosin B Mg-ATPase

The question arose to which of the two regulatory systems is predominant in squid muscle. Kendrick-Jones et al.(2) reported that scallop myosin B Mg-ATPase(myosin-linked) was fully activated only at higher calcium concentrations than rabbit skeletal myosin B Mg-ATPase. Initially, the Ca and Sr requirements for Mg-ATPase of squid myosin B were studied in comparison with those of rabbit myosin B and two hybrid actomyosin: rabbit actin/squid myosin (myosin-linked regulatory system) and crude squid actin/rabbit myosin(actin-linked regulatory system). As shown in Fig. 18, the Ca and Sr requirements of squid myosin B (●) were identical with those of the former hybrid (myosin-linked)(△) rather than the latter(actin-linked)(□). The Ca and Sr requirements of the actin-linked regulatory system did not appear in squid myosin B, and it seems that the myosin-linked regulatory system is more closely associated with squid myosin B. It should be noted that the Sr requirement of squid crude actin/rabbit myosin(actin-linked)(□) was identical to that of rabbit myosin B(○), which is actin-linked. It is, therefore, likely that a myosin-linked regulatory system




Fig. 18. Mg-ATPase activity of squid myosin B and reconstituted actomyosin as a function of calcium ion or strontium ion concentration

(●), Squid myosin B(95 $\mu\text{g/ml}$); (Δ), squid myosin (110 $\mu\text{g/ml}$) and skeletal actin(49 $\mu\text{g/ml}$); ($\square$), squid crude actin(139 $\mu\text{g/ml}$) and skeletal myosin(93 $\mu\text{g/ml}$); ($\circ$), rabbit skeletal myosin B(102 $\mu\text{g/ml}$). The ATPase assay was conducted as described in Table I. The maximal and minimal ATPase activities obtained under the present conditions were scaled as 100 and 0 %, respectively.

is predominant in squid myosin B.

It should also be pointed out(Fig. 18) that the concentration of calcium or strontium required for Mg-ATPase of squid mantle myosin B is higher than that required for Mg-ATPase of skeletal myosin B; half maximal activation of Mg-ATPase was obtained at 0.8 μ M calcium ions or 28 μ M strontium ions with rabbit myosin B and at 2.5 μ M calcium ions or 140 μ M strontium ions with squid myosin B.

2-(B)-b) Effects of squid and rabbit native tropomyosins
on the Ca requirement for Mg-ATPase of squid
myosin B

The results described above strongly suggested that myosin-linked Ca regulation was operating in squid myosin B. In order to establish this findings, the author tested whether addition of large amounts of native tropomyosin to squid myosin B shifted the Ca requirement of myosin B ATPase from myosin-linked to actin-linked or not. As shown in Fig. 19, neither squid native tropomyosin nor rabbit one shifted Ca requirement of squid myosin B.




Fig. 19. Effect of native tropomyosin from squid or rabbit on the Ca requirement of Mg-ATPase of squid myosin B

ATPase assay was conducted as described in Fig. 18, (○) squid myosin B alone(108 $\mu\text{g/ml}$), (△) squid myosin B(108 $\mu\text{g/ml}$) with rabbit native tropomyosin(23 $\mu\text{g/ml}$), and (□) squid myosin B(108 $\mu\text{g/ml}$) with squid native tropomyosin(30 $\mu\text{g/ml}$).

2-(C) Regulatory light chain from squid muscle

The author found that myosin-linked regulation rather than actin-linked regulation is dominantly operating in squid myosin B. Therefore, it is necessary to study on the myosin-linked Ca regulation mechanism in squid mantle muscle.

At first, the author attempted to isolate two types of small polypeptide chain(so-called light chain) from squid myosin, which was considered to be regulatory subunit in myosin-linked regulation, and to study on the functional role of light chain in myosin-linked Ca regulation in detail. For the purpose of this experiment, scallop adductor myosin has to be employed by the simple reason that squid myosin was not desensitized by EDTA-treatment or by any other techniques tried up to now.

2-(C)-a) Isolation of highly purified squid light chains

As reported by Kendrick-Jones et al.(32), squid myosin showed two light chain bands in SDS gel and a single band in urea gel, whereas scallop myosin showed a single band in SDS gel and two bands in urea gel. These results indicate that squid light chains differ in size but have a similar net charge and scallop light chains differ in net charge but have a similar size. The two types of




Fig. 20. Sephadex G-75 gel filtration of squid light chains

Squid light chains were precipitated with ammonium sulfate between 50 and 60 % saturation, and the precipitate was applied to a column of Sephadex G-75(2.5 x 90 cm) in 0.15 M KCl, 20 mM Tris-HCl(pH 8.0), and 0.5 mM DTT. Fractions of 5 ml were collected. The inset photos are of SDS gel electrophoretic patterns of the fractions obtained. SDS gel electrophoresis was conducted as described in Fig. 14.

light chains of squid myosin were designated as LC-1 (17,000 daltons) and LC-2(15,000 daltons), and those of scallop myosin as EDTA-LC and SH-LC.

Squid light chains were not released from myosin either by EDTA-treatment(10, 32) or by DTNB-treatment(36) (data not shown), but they were released by 4 M urea(11). Chart II outlines the procedures adopted for the isolation of squid light chains(see also "Methods and materials"). The fractions obtained by ammonium sulfate salting-out were examined by SDS-gel electrophoresis. Only a very small amount of light chains was precipitated at 50 % saturation, and thus the fraction precipitating between 50 and 100% saturation with ammonium sulfate was collected. The fraction was dialyzed against a solution containing 0.15 M KCl, 20 mM Tris-HCl(pH 8.0) and 0.1 mM DTT, and centrifuged to remove insoluble materials. The supernatant was subjected to gel filtration, using the same solution as for dialysis. As shown in Fig. 20, proteins with molecular sizes larger than those of light chains appeared at around the void volume((a) in Fig. 20), and light chains in a pure form(see SDS gel electrophoretic pattern (b) in Fig. 20) were retained and thus appeared in a separate fraction.

Various types of chromatography have been reported to be effective in separating the myosin light chains. The method of Perrie et al.(12) with some modification,




Fig. 21. Separation of light chains of squid myosin by DEAE-Sephadex A25 chromatography

A solution (approximately 30 mg/100 ml) of the light chain mixture obtained by Sephadex G-75 gel filtration was applied to a column (1 x 80 cm) of DEAE-Sephadex A25 pre-equilibrated with 0.15 M KCl, 20 mM Tris-HCl (pH 8.0), and 0.5 mM DTT. A linear gradient of KCl, as indicated by the dashed line, was applied from 0.15 M to 0.3 M in 20 mM Tris-HCl (pH 8.0), and 0.5 mM DTT. The flow rate was 10 ml/h, and the volume of each fraction was 5 ml. The SDS gel electrophoretic patterns of the light chain-1 fractions (b) and the light chain-2 fractions (a) are shown in the insets (10 % polyacrylamide, 0.1 % SDS).

satisfactory to separate LC-1 and LC-2 of squid myosin. As shown in Fig. 21, LC-2(15,000 daltons) was eluted with approximately 0.22 M KCl, and LC-1(17,000 daltons) with approximately 0.25 M KCl.

2-(C)-b) Effect of 2-mercaptoethanol on the patterns of squid light chain in disc gel electrophoresis

Disc gel electrophoresis was conducted in the presence of various concentrations of 2-mercaptoethanol. Figure 22A shows that the squid LC-2 preparation gave a single band which showed no change in mobility at any of the concentrations of 2-mercaptoethanol tested. In contrast, Fig. 22B shows that the squid LC-1 preparation gave three bands (one major and two minor) in the absence of 2-mercaptoethanol, two bands in the presence of 0.25 % 2-mercaptoethanol, and finally a single band when the concentration of 2-mercaptoethanol was increased to 10 to 25 %. It seems therefore reasonable to conclude that squid LC-1 is equivalent to scallop SH-LC.

2-(C)-c) Effect of calcium on the patterns of squid light chains in disc gel electrophoresis

Head et al(37) studied the calcium-binding abilities of troponin-C preparations from various sources by conducting urea gel electrophoresis in the presence of




Fig. 22. Effect of 2-mercaptoethanol on the patterns of squid light chains in disc gel electrophoresis

Disc gel electrophoresis was carried out by the method described by Davis. A light chain sample was dissolved in 0.1 M Tris-HCl (pH 8.9) and various concentrations of 2-mercaptoethanol. It was applied to a gel rod of 7.5 % polyacrylamide; a solution of 0.05 M Tris and 0.19 M glycine (pH 9.4) was used in the electrophoretic run. Electrophoresis was performed at 2 mA/rod, and the temperature was maintained at around 10°C by circulating cold water. (A), Squid LC-2, (B), squid LC-1. (a) 0 %, (b) 0.0025 %, (c) 0.025 %, (d) 0.25 %, (e) 2.5 %, (f) 10 %, and (g) 25 % of 2-mercaptoethanol was present.




Fig. 23. Calcium-induced change in the mobility of squid light chain in disc gel electrophoresis

Disc gel electrophoresis was carried out as described in Fig. 22, except that 10 % 2-mercaptoethanol was present in this experiment. +Ca and -Ca denote the presence of 1 mM CaCl_2 and 1 mM EGTA, respectively. BPB was used as front marker.

calcium(1 mM CaCl_2) or in its absence(5 mM EGTA). The author found that in urea gel electrophoresis, calcium had no effect on the mobility of either LC-1 or LC-2.

On the other hand, as shown in Fig. 23, the author found that in disc gel electrophoresis, calcium caused a decrease in the mobility of LC-2 but no change in that of LC-1. It is therefore suggested that squid LC-2 can bind calcium. This view was supported, as described in the next section, by the finding that calcium can induce changes in UV absorption spectrum of squid LC-2.

2-(C)-d) UV absorption spectra of isolated squid light chains and their changes induced by calcium or strontium ions

Figure 24 shows the UV absorption spectra of squid LC-1 and LC-2; they are very similar to each other, and also to those of rabbit skeletal light chains(38), and scallop light chains(22), in that absorption peaks characteristic of phenylalanine residues are noticeable.

Figure 25 shows that 1 mM CaCl_2 induced practically no change in the UV absorption spectrum of squid LC-1, but a large changes in that of squid LC-2, indicating that calcium does not bind to LC-1 but binds to LC-2. Three positive peaks and three negative peaks were seen, and the largest negative changes appeared at 287 nm(tyrosine residue) and at 293 nm(tryptophan residue).




Fig. 24. Ultraviolet absorption spectra of isolated squid light chains

UV absorption spectra of isolated light chains were measured in 0.1 M KCl, and 20 mM histidine-KOH(pH 6.8).

-----, light chain-1(0.98 mg/ml); ———, light chain-2(1.15 mg/ml). $E_{280\text{ nm}}^{1\%} = 5.4$ for LC-1, and 6.9 for LC-2.




Fig. 25. Calcium-induced spectral change of isolated squid light chains

The absorption difference spectra were measured in 0.1 M KCl, and 20 mM histidine-KOH(pH 6.8). 1 mM CaCl_2 was present in the sample cell, and 1 mM EGTA in reference cell. Base lines are indicated by dashed lines. (A) Light chain-1(0.98 mg/ml), and (B) light chain-2(1.15 mg/ml).




Fig. 26. Calcium and strontium binding to squid light chain-2

The relative magnitude of the difference spectrum at 293 nm was plotted as a function of the free calcium or strontium ion concentration. The concentrations of light chain-2 employed were the same as in Fig. 25. The concentrations of free calcium and strontium ions were maintained by using Ca-EGTA and Sr-EGTA buffers. Stability constants of $5 \times 10^5 \text{ M}^{-1}$ and $2 \times 10^2 \text{ M}^{-1}$ were used for the Ca-EGTA and Sr-EGTA complexes, respectively. The reference cell always contained 1 mM EGTA. Open and closed symbols represent different light chain-2 preparations (● ○) Calcium-induced difference spectra, (▲ △) strontium induced difference spectra. The Ca requirement(---) and Sr requirement(-.-.-) of skeletal acto-squid myosin ATPase activity are also shown.

The change at 293 nm was measured as a function of calcium or strontium ion concentration, and the results are shown in Fig. 26. The half-maximal change was obtained with 6.6 μ M free calcium ions or with 600 μ M free strontium ions. These values are somewhat larger than those obtained previously(10) for the half-maximal activation of actomyosin ATPase: 2 μ M free calcium and 140 μ M free strontium.

2-(C)-e) UV absorption spectrum of scallop EDTA-LC
and its change induced by calcium or strontium
ions

It was also found that divalent cations induce a spectral change of scallop EDTA-LC(Fig. 27). Figure 27A shows the UV absorption spectrum of scallop EDTA-LC, which is identical to that reported by Kendrick-Jones et al.(4). Figure 27B shows that 1 mM CaCl_2 induced a change in the UV absorption spectrum of scallop EDTA-LC; it is different from that of squid LC-2 in lacking the negative change at 293 nm(tryptophan residue).

However, as shown in Fig.28, the divalent cation concentrations required to induce the spectral changes were too high for this process to account for the sensitivity of scallop myosin B ATPase to divalent cations. The calcium concentration required for half-maximal activation of scallop myosin B ATPase was 2.5 μ M free calcium ions,




Fig. 27. Ultraviolet absorption spectrum and calcium-induced spectral change of scallop EDTA-light chain

(A) The UV absorption spectrum of EDTA-light chain (2.2 mg/ml) was obtained in a medium containing 0.1 M KCl, 20 mM histidine-KOH(pH 6.8). (B) The calcium-induced spectral change of EDTA-light chain was obtained in the medium described above, except that the sample cell also contained 1 mM CaCl_2 and the reference cell 1 mM EGTA. The dashed line indicates the base line.




Fig. 28. The calcium and strontium requirements for producing difference spectra in scallop EDTA-light chain and for activation of scallop myosin B ATPase

The relative magnitude of the difference spectrum at 287 nm was plotted as a function of the free calcium or strontium concentration. The experimental conditions were as described in Fig. 25. The Mg-ATPase activities of scallop myosin B were measured at 25°C in a medium containing 30 mM KCl, 20 mM Tris-maleate(pH 6.8), 2 mM MgCl_2 , 1 mM ATP, and various concentrations of free calcium or strontium ions. ($\bigcirc$) Calcium-induced difference spectrum, (Δ) strontium-induced difference spectrum, ($\bullet$) Ca requirement for Mg-ATPase, and ($\blacktriangle$) Sr requirement for Mg-ATPase.

whereas that required for the half-maximal change in the induced spectral change was 320 μM free calcium ions; the strontium concentration required for half-maximal activation of myosin B ATPase was 100 μM free strontium, whereas that required for the half-maximal change in the induced spectral change was 4.0 mM free strontium ions.

2-(D) Desensitization and resensitization of scallop myosin

2-(D)-a) Desensitization of scallop myosin by EDTA-treatment

As reported previously(10), EDTA-treatment did not remove any light chains from squid myosin and did not reduce the Ca sensitivity of acto-squid myosin ATPase. Thus, desensitized scallop myosin was used to study the functional role of squid light chains.

EDTA-treatment of scallop myosin was repeated five times at 0 - 4°C; as shown in Fig. 29, EDTA-LC were selectively and completely removed from scallop myosin.

2-(D)-b) Binding of squid light chains to scallop desensitized myosin

The abilities of squid LC-1 and of squid LC-2 to bind to desensitized(EDTA-treated) scallop myosin((b) in Fig. 29) were tested in the following way(see also "Methods




Fig. 29. Binding of squid light chains to scallop desensitized myosin

Scallop desensitized myosin was prepared by EDTA-treatment as described in Fig. 31. Squid light chains were added to scallop desensitized myosin in 0.5 M KCl, 20 mM Tris-maleate(pH 7.5), and 2 mM $MgCl_2$. Ten volumes of 1 mM $MgCl_2$ were then added, and the mixture was centrifuged. Unbound light chains in the supernatant were removed. The myosin pellet was dissolved in 8 M urea, 0.1 M Tris-HCl(pH 8.9), and 5 % 2-mercaptoethanol, and then applied to a 7.5 % polyacrylamide gel rod containing 8 M urea and 0.375 M Tris-HCl(pH 8.9). Urea-gel electrophoresis was carried out by a modification of the method of Perrie *et al.* The running buffer contained 0.05 M Tris and 0.19 M glycine(pH 9.4). Approximately 50 μ g of protein was loaded on each gel. (a) Untreated scallop myosin, (b) desensitized scallop myosin, (c) scallop desensitized myosin incubated with squid LC-2, (d) squid LC-2, (e) scallop desensitized myosin incubated with squid LC-1, and (f) squid LC-1.

and materials"): 0.6 mg/ml of LC-1 or LC-2 was mixed with 6 mg/ml of desensitized scallop myosin in a solution containing 0.5 M KCl, 20 mM Tris-maleate(pH 7.5), 2 mM MgCl_2 , and 1 mM DTT, and the mixture was kept overnight in ice-water. It was then diluted with 10 vols of 1 mM MgCl_2 , and centrifuged to remove unbound light chains in the supernatant. The myosin sediment was washed with 0.1 M KCl, 20 mM Tris-maleate(pH 7.5), 1 mM MgCl_2 , and 1 mM DTT, then dissolved in 8 M urea, 0.1 M Tris-HCl(pH 8.9), and 5 % 2-mercaptoethanol. Figure 29 shows that desensitized scallop myosin took up squid LC-2 in an amount approximately equal to its own SH-LC, which remained unchanged during the desensitization and resensitization procedures, and that it took up practically no squid LC-1 (see also Fig. 32).

2-(D)-c) Resensitization of scallop desensitized myosin by squid light chain.

The resensitizing effects of squid light chains and scallop EDTA-LC were examined, and the results obtained are shown in Fig. 30. Various amounts of scallop EDTA-LC, squid LC-1, or squid LC-2 were added to desensitized (EDTA-treated) scallop myosin in the presence of magnesium ion, and then combined with rabbit skeletal actin in a weight ratio of 1/3 for actin/desensitized myosin. The Mg-ATPase activity of the reconstituted actomyosin was measured in the presence and absence of calcium. Mg-ATPase




Fig. 30. Resensitization of scallop desensitized myosin by squid light chains and by scallop EDTA-light chain

Various amounts of squid light chains or scallop EDTA-light chain were added to scallop desensitized myosin in the medium described in Fig. 29. Mg-ATPase activities of the desensitized myosin were assayed at 25°C in the presence of skeletal F-actin in a medium containing 30 mM KCl, 20 mM Tris-maleate(pH 6.8), 2 mM MgCl_2 , 1 mM ATP, and either 0.1 mM CaCl_2 (closed) or 0.5 mM EGTA(open). 100 - 110 $\mu\text{g/ml}$ of desensitized myosin and 35 $\mu\text{g/ml}$ of skeletal F-actin were used. (■ □) Squid LC-1, (● ○) squid LC-2, (▲ △) scallop EDTA-LC.

of acto-desensitized myosin showed no Ca sensitivity. Squid LC-1, which did not bind to desensitized scallop myosin(Fig. 29), had no effect on the ATPase activity either in the presence or absence of calcium. On the other hand, squid LC-2 increased the activity in the presence of calcium and decreased it in its absence, thus conferring Ca sensitivity on acto-desensitized myosin. Moreover, the resensitizing effect of squid LC-2 was quantitatively equal to that of scallop EDTA-LC.

These results with scallop myosin are somewhat different from those reported by Szent-Györgyi *et al.*(4). For instance, they reported that only one mol of EDTA-LC can readily be removed by EDTA-treatment, with complete loss of Ca sensitivity, and that removal of the remaining EDTA-LC requires DTNB-treatment, resulting in incomplete resensitization. Based mainly on these results, they proposed that cooperation between either two head(S-1) of scallop myosin or two EDTA-LC is involved in the molecular mechanism of Ca regulation in scallop adductor muscle. However, the author found(Fig. 29) that EDTA-treatment removes all the EDTA-LC from scallop adductor myosin. The author therefore conducted further studies on the desensitization and resensitization of scallop adductor myosin. Removal of EDTA-LC was evaluated by measuring the staining densities in urea gel electrophoretic patterns, and Ca sensitivity was determined by measuring Mg-ATPase activities in the presence and absence of calcium.




Fig. 31. EDTA-treatment of scallop myosin and myofibrils

Scallop myosin(A) and myofibrils(B) were suspended in 10 mM EDTA, 30 mM KCl, and 10 mM Tris-HCl(pH 7.5) for 15 min at 0 - 4°C or at 20°C, then centrifuged at 6,000 x g for 10 min, thus obtaining the sediments. This procedure was repeated 5 times. The contents of EDTA-LC in myosin and myofibrils were determined by urea-gel electrophoresis, and expressed as the staining density ratio of EDTA-LC band to SH-LC band. The Ca sensitivity was determined in terms of Mg-ATPase activity and is expressed as $(1 - A_{EG}/A_{Ca}) \times 100$, where A_{EG} and A_{Ca} are the activity in the presence of EGTA and that in the presence of $CaCl_2$, respectively. (A) Scallop myosin, (B) scallop myofibrils, (● ○) EDTA-treatment at 0 - 4°C, (▲ △) EDTA-treatment at 20°C, (● ▲) Ca sensitivity, and (○ △) EDTA-LC content.

Since Szent-Györgyi et al. stated that reversibility of desensitization-resensitization is better with myofibrils than with myosin alone, the author also carried out EDTA-treatment with myofibrils. It was thus confirmed (Fig. 31) that EDTA-treatment removed all the EDTA-LC from scallop myosin or myofibrils. Moreover, Fig. 31 shows that scallop desensitized myosin retaining one mol of EDTA-LC per mol of myosin also retained a definite Ca sensitivity, and that removal of not half the total amount of EDTA-LC but of all of it is required to achieve a complete loss of Ca sensitivity.

2-(D)-d) Relationship between EDTA-LC(or squid LC-2) content of myosin and Ca sensitivity of acto-scallop myosin.

Using 5-times-EDTA-treated scallop myosin, which was completely free from EDTA-LC and from Ca sensitivity, its recombination with isolated scallop EDTA-LC, squid LC-1 and LC-2 as well as its resensitization in terms of the Mg-ATPase activity of reconstituted actomyosin was studied. The results obtained are shown in Fig. 32. The amount of light chains bound to scallop desensitized myosin was expressed as the ratio of the Fast Green staining density of the recombined light chain band to that of the SH-LC band retained in desensitized myosin. The symbol(●) in Fig. 32A indicates the density ratio of EDTA-LC to SH-LC in untreated scallop myosin. The




Fig. 32. Recombination and resensitization of scallop desensitized myosin

(A) Recombination of light chains with scallop desensitized myosin was determined by urea-gel electrophoresis. The amount of light chains bound to desensitized myosin was determined by densitometry at 640 nm with Fast Green staining, and is expressed in terms of (light chain bound)/(scallop SH-LC) $\times$ 100. (X) Squid light chain-1, (●) squid light chain-2, (■) scallop EDTA-LC, and (▲) a mixture of squid light chain-2 and scallop EDTA-LC (1 : 1, w/w). (⊙) The EDTA-LC content in untreated scallop myosin. (B) Resensitization of scallop desensitized myosin by squid light chains or by scallop EDTA-LC was studied by measuring the Mg-ATPase activities in combination with skeletal F-actin. The recovery of Ca sensitivity was estimated by measuring Mg-ATPase activities (see Figs. 31 and 29). (X) Squid light chain-1, (○) squid light chain-2, (□) scallop EDTA-LC, and (△) a mixture of squid light chain-2 and scallop EDTA-LC (1 : 1, w/w). (⊙) Ca sensitivity of untreated myosin.

Ca sensitivity was expressed in terms of $(1 - A_E/A_{Ca}) \times 100$, where A_E and A_{Ca} are the ATPase activity in the presence of EGTA and that in the presence of calcium, respectively. The symbol($\odot$) in Fig. 32B indicates the Ca sensitivity of untreated scallop myosin.

Dashed lines in Fig. 32 indicate that both the recombination and resensitization were saturated when the amount of either EDTA-LC or squid LC-2 added reached approximately 7.5 % weight ratio to scallop desensitized myosin, where the amount of the light chains bound was practically equal to that of EDTA-LC present in untreated scallop myosin. If one assumes that one mol of scallop myosin(480,000 daltons) is composed of two mol of heavy chains($2 \times 206,000$ daltons), two mol of EDTA-LC($2 \times 17,000$ daltons), and two mol of SH-LC($2 \times 17,000$ daltons), then the weight ratio of EDTA-LC to desensitized myosin is expected to be $17/(206 + 17) = 0.076$. The numerical agreement between the expected value and the observed value is remarkably good. It should also be noted that a small amount of squid LC-1 preparation can bind to desensitized myosin but does not confer any Ca sensitivity on desensitized myosin. The results shown in Fig. 32 indicate that the restoration of Ca sensitivity was proportional to the amount of scallop EDTA-LC or squid LC-2 bound to scallop desensitized myosin. It therefore appears that cooperation of two mol of EDTA-LC in scallop myosin is not involved in myosin-linked Ca regulation(4).

2-(D)-e) Competitive binding of EDTA-LC and squid
LC-2 to scallop desensitized myosin

The recombination and resensitization were also studied by adding a mixture of scallop EDTA-LC and squid LC-2 in various weight ratios, while the total amount of light chains added was kept at 10 % (w/w) of desensitized myosin. As shown in Fig. 33, as the ratio of squid LC-2 to EDTA-LC increased in the mixture added, the amount of EDTA-LC bound decreased and that of LC-2 increased, thus indicating competitive binding between EDTA-LC and squid LC-2. Meanwhile, the total amount of light chains bound (EDTA-LC + squid LC-2), as well as the Ca sensitivity, remained nearly unchanged. It is therefore concluded that squid LC-2 is functionally identical with scallop EDTA-LC, and that scallop desensitized myosin free from EDTA-LC retains its original abilities to bind light chains and to respond to calcium ions.




Fig. 33. Competitive binding of squid light chain-2 and scallop EDTA-LC to scallop desensitized myosin

A mixture of squid light chain-2 and scallop EDTA-LC in a various ratios(the total amount of light chains added was kept at 10 % of myosin by weight) was added to scallop desensitized myosin in the medium described in Fig. 30. The amount of light chains bound was determined by urea-gel electrophoresis(see Fig 32). The amount of squid light chain-2 bound was expressed as a ratio to the total amount of light chain bound(○). The total amount of light chain bound was also expressed as the ratio to the amount of scallop SH-LC(□). The recovery of Ca sensitivity (Δ) was estimated by measuring Mg-ATPase activities(see Figs. 29 and 31).

2-(E) Calcium binding of myosin and of regulatory
light chain

2-(E)-a) Calcium binding of isolated squid LC-2 and
scallop EDTA-LC

The effects of calcium and strontium ions on the UV absorption spectra of light chain subunits isolated from squid myosin and from scallop adductor myosin were studied. The author then found (a) that the divalent cations induced difference spectra in regulatory light chains(squid LC-2 and scallop EDTA-LC) but not in the second type of light chains(LC-1 and SH-LC), and (b) that the strontium and calcium concentrations required for inducing difference spectra were much higher than those required for activation of the Mg-ATPase of myosin B. In other words, the results obtained were qualitatively in favor of, but quantitatively inconsistent with, the possibility that the divalent cation-induced change in the regulatory light chain conformation may be the first step in a series of reactions, which lead to "contraction" of actomyosin.

In the present study, the author measured calcium binding with regulatory light chains in an isolated form and with myosins(light chains in a bound form). Isolated regulatory light chains were purified by Sephadex G-75 gel filtration. The molecular size of scallop EDTA-LC estimated by the gel filtration was about 5×10^4 daltons




Fig. 34. Calcium binding to isolated regulatory light chains from squid and scallop muscles

Calcium binding was measured at 20°C by using ^{45}Ca in equilibrium dialysis in a medium containing 0.1 M KCl, 20 mM histidine (pH 6.8), and no (○) or 2.5 mM MgCl_2 (●). Free calcium concentration was set by using a Ca-EGTA buffer system. (A) Squid LC-2, (B) scallop EDTA-LC.

and that of squid LC-2 was about 2×10^4 daltons. Scallop EDTA-LC was therefore suggested to be in an aggregated form. Calcium binding to regulatory light chain was then measured by using ^{45}Ca in equilibrium dialysis(21).

As shown in Fig. 34, isolated light chains had two types of calcium binding site: calcium binding with the first type of sites occurred at about $1 - 10 \mu\text{M}$ free calcium whereas that with the second type occurred at about $100 - 1,000 \mu\text{M}$ free calcium. The amount of calcium bound to the first type of sites was small: approximately 0.4 mol of calcium per mol of squid LC-2 (1.5×10^4 daltons), and 0.15 mol of calcium per mol of scallop EDTA-LC(1.7×10^4 daltons). Moreover, the amount of calcium bound was greatly reduced when 2.5 mM MgCl_2 was present.

Lower calcium binding ability of isolated scallop EDTA-LC was also reported by Szent-Györgyi et al.(4). In the present study, the author found that squid regulatory light chain retained its calcium binding ability even if it was released from heavy chain. In this respect, squid regulatory light chain is clearly different from EDTA-LC of scallop.

2-(E)-b) Calcium binding of squid and scallop myosin

On the other hand, as shown in Fig. 35, calcium




Fig. 35. Calcium binding to myosins from squid and scallop muscles

Calcium binding was measured under the conditions described in Fig. 34, except that 0.3 M KCl was used instead of 0.1 M KCl. The same symbols as in Fig. 34 are used.

(A) Squid myosin. (B) Scallop myosin.

binding with myosins was very different from that with isolated light chains. The amount of calcium bound with myosin was about two mol per mol of myosin(4.8×10^5 daltons), which is equivalent to one mol of calcium per mol of light chain(in a bound form). Only one type of binding site was detectable, and the half-maximal binding occurred at about $2.0 \mu\text{M}$ free calcium with squid myosin, and at about $7.9 \mu\text{M}$ free calcium with scallop myosin. These concentrations of free calcium ions were about equal to those required for activation of the ATPase of acto-squid myosin and for that of acto-scallop myosin (see Figs. 37 and 50A). Therefore, the difference between the calcium concentration required for activating myosin B ATPase and that required for inducing difference spectra of isolated light chains may be understood as the difference in calcium binding between a bound form of light chains and an isolated form of light chains(see Figs. 34 and 35).

2-(E)-c) Effects of desensitization and resensitization
on the calcium binding of scallop myosin

Moreover, it was also able to be shown(Fig. 36) that the regulatory light chain is responsible for the calcium binding described above. The author studied the effect of desensitization and resensitization of myosin on its calcium binding ability. Figure 36 shows that desensitized scallop myosin, which was completely free from EDTA-LC,

lost its calcium binding ability(A), and that the lost ability was fully recovered by adding either scallop EDTA-LC(C) or squid LC-2(B) to desensitized scallop myosin(approximately two mol of light chain per mol of myosin). It should be pointed out that calcium binding to resensitized myosin, like that to untreated myosin (see Fig. 35), was unaffected by the addition of 2.5 mM MgCl_2 .

It was found that scallop myosin and scallop resensitized myosin were capable of binding approximately two mol of calcium per mol of myosin even if MgCl_2 was absent. Moreover, EDTA-LC in an isolated form was found to have little calcium binding ability at around 10 μM calcium ions(see Fig. 34). Therefore, these results rather suggest that EDTA-LC was present in a form bound to myosin even if MgCl_2 was absent. This view was further supported by the recent finding of Ojima *et al*(personal communication) that the release of EDTA-LC from Akazara scallop myosin was prevented by the presence of small amounts of calcium even if MgCl_2 was absent.




Fig. 36. Calcium binding to desensitized and resensitized scallop myosins

Calcium binding was measured under the same conditions as in Fig. 35. Symbols are the same as in Fig. 35.

(A) Desensitized scallop myosin, (B) resensitized scallop myosin with squid LC-2, and (C) resensitized scallop myosin with scallop EDTA-LC.

2-(F) Calcium sensitivity of squid myosin alone

In the previous sections, Ca sensitivity of acto-squid myosin or acto-scallop myosin were studied. Recently, Asada et al.(39) showed that Mg-ATPase of clam foot myosin alone was sensitive to calcium ion. In this section, the Ca sensitivity of myosin alone from squid and scallop was therefore studied.

2-(F)-a) Ca requirement for activation of Mg-ATPase of squid myosin alone

As shown in Fig. 37, Mg-ATPase of squid myosin alone was activated by a very low concentration of calcium ions(1 - 10 μ M) and half-maximal activation was obtained at 3 μ M free calcium ion, which was identical to that obtained in the activation of Mg-ATPase of acto-squid myosin. These results indicated that squid myosin Mg-ATPase was sensitive to calcium ion as well as actomyosin Mg-ATPase. It should be pointed out that myosin Mg-ATPase activity was remarkably activated by F-actin in the presence of calcium. In addition, Ca-ATPase activity of squid myosin(activity in the absence of MgCl_2) are able to be distinguished from Mg-ATPase activity in the concentration of calcium required for half-maximal activation; 3 μ M calcium for Mg-ATPase and 150 μ M for Ca-ATPase.




Fig. 37. Calcium requirements for Mg-ATPase activity of squid myosin and of acto-squid myosin and for Ca-ATPase activity of squid myosin alone

Mg-ATPase of myosin alone(○) and of acto-myosin (●) were assayed in a medium containing 30 mM KCl, 20 mM Tris-maleate(pH 6.8), 2 mM $MgCl_2$ and 1 mM ATP at 25°C. Ca-ATPase of myosin alone(Δ) was assayed in the above medium without 2 mM $MgCl_2$. Free calcium concentration was set by using Ca-EGTA buffer system.




Fig. 38. Temperature dependence of Mg-ATPase activity of squid myosin

Mg-ATPase activity of squid myosin alone was measured in a medium containing 30 mM KCl, 20 mM Tris-maleate(pH 6.8), 2 mM MgCl_2 , 1 mM ATP and either 0.1 mM CaCl_2 (●) or 0.5 mM EGTA (○). Reaction temperature was changed from 5°C to 30°C.




Fig. 39. PH dependence of Mg-ATPase activity of squid myosin

Mg-ATPase of squid myosin alone was assayed in a medium containing 30 mM KCl, 2 mM MgCl_2 , 1 mM ATP and either 0.1 mM CaCl_2 (●) or 0.5 mM EGTA (○). PH for ATPase assay was changed from 6.0 to 9.5 using 20 mM Tris-maleate or Tris-HCl buffer system.

2-(F)-b) Effects of temperature and pH on Ca sensitivity
of squid myosin

As shown in Fig. 38, change in the temperature for the ATPase assay from 5°C to 30°C did not affect the Ca sensitivity of squid myosin, that is, Ca sensitivity was observed at all temperature tested.

Ca sensitivity of squid myosin was also detectable even if the pH for the ATPase assay was changed from 6.0 to 9.5(Fig. 39).

2-(F)-c) Effect of KCl concentration on Ca sensitivity
of squid myosin

The change in KCl concentration for the ATPase assay remarkably affected the Ca sensitivity of squid myosin alone. As shown in Fig. 40, Mg-ATPase of squid myosin alone showed maximal activity at around 0.3 M KCl in the presence of calcium. The activity in the absence of calcium was well inhibited in low-salt medium(lower than 0.2 M KCl), but it was gradually increased with increasing KCl concentration, reaching to the identical activity with that in the presence of calcium at 0.4 M KCl. In other word, the Ca sensitivity of Mg-ATPase activity of squid myosin alone was observed only when the KCl concentration was lower than 0.4 M. On the other hand, actin-activation of Mg-ATPase of myosin occurred when the KCl concentration was lower than 0.2 M. Figure 40B shows that




Fig. 40. KCl concentration dependence of Mg-ATPase and that of superprecipitation of acto-squid myosin.

(A) The ATPase activity of myosin(● ○) and that of acto-squid myosin(▲ △) were assayed at 25°C in a medium containing 20 mM Tris-maleate(pH 6.8), 2 mM MgCl_2 , 1 mM ATP, either 0.1 mM CaCl_2 (closed) or 0.5 mM EGTA(open), and various concentrations of KCl from 0.03 M to 0.5 M.

(B) The superprecipitation activity of acto-squid myosin was assayed under the conditions used for the ATPase activity(A), but in the presence of 0.1 mM CaCl_2 .




Fig. 41. Effect of KCl concentration on Ca sensitivity of squid myosin.

ATPase of myosin alone was assayed in the medium containing 30 mM KCl, 20 mM Tris-maleate(pH 6.8), 2 mM MgCl_2 , 1 mM ATP and either 0.1 mM CaCl_2 (●) or 0.5 mM EGTA (○). At the time indicated by the arrows, KCl concentration was changed from 0.03 M to 0.4 M without changing other medium compositions.




Fig. 42. Recovery of Ca sensitivity of squid myosin by dilution of KCl.

ATPase was assayed in 0.4 M KCl, 20 mM Tris-maleate (pH 6.8) 2 mM MgCl_2 , 1 mM ATP and either 0.1 mM CaCl_2 (●) or 0.5 mM EGTA (○). At the time indicated by the arrows, KCl concentration was reduced to 0.2 M without changing other medium compositions.

in accordance with the actin-activated Mg-ATPase in Fig. 40A, superprecipitation of acto-squid myosin occurred only when the KCl concentration was lower than 0.2 M. Therefore, it was able to be distinguished the effect of KCl on actin-activation from that on Ca sensitivity of myosin alone.

The effect of KCl on Ca sensitivity was further studied by KCl concentration jump method. Figure 41 showed the time course of Pi liberation by Mg-ATPase of squid myosin alone. When the KCl concentration was rapidly increased from 0.03 M to 0.4 M at the time indicated by the arrows, the immediate increase in Pi liberation in the absence of calcium occurred, and it was no longer sensitive to calcium. It should be noted that lag phase in the time course of Pi liberation was not found during this treatment. Then the recovery of Ca sensitivity by dilution of KCl was tested. As shown in Fig. 42, when the KCl concentration in assay medium was decreased rapidly from 0.4 M to 0.2 M, the Ca sensitivity was found to be recovered with no lag time.

2-(F)-d) ITP as substrate for squid myosin

In order to know whether ATP was essential for Ca sensitivity of myosin as substrate or not, ITP instead of ATP was used in the same experiment of Fig. 42. The KCl concentration dependence of Mg-ATPase of squid myosin




Fig. 43. KCl concentration dependence of Mg-ITPase of squid myosin

ITPase activity of squid myosin alone was assayed under the same conditions as in Fig. 40A, except that 0.5 mM ITP was used instead of 1 mM ATP. The same symbols were also used.

alone was shown in Fig. 43. Essentially the same KCl concentration dependence of ITPase activity was obtained as that of ATPase activity; Mg-ITPase showed Ca sensitivity in a low-salt medium, but lost it in high-salt medium. Therefore, ATP was found not to be essential for Ca sensitivity of myosin.

2-(F)-e) Effect of urea on the Ca sensitivity of
squid myosin

Ca sensitivity of Mg-ATPase of squid myosin alone was measured in the medium containing various concentrations of urea. As shown in Fig. 44, Mg-ATPase of myosin alone was Ca sensitive when the urea concentration was lower than 1 M and it was lost in the presence of 1.2 M urea. It should be pointed out that the Mg-ATPase activity of myosin alone in the presence of calcium was unaffected by urea. The effects of urea on Ca sensitivity of myosin alone and on actin-activation of Mg-ATPase were then compared. Figure 44 showed that actin-activation of myosin Mg-ATPase in the presence of calcium decreased as the urea concentration increased, though it still occurred even at 1 M urea. On the other hand, actin-activation in the absence of calcium slightly increased at concentration of urea higher than 1 M. The decrease in actin-activation of myosin Mg-ATPase in the presence of calcium was closely related to the irregular conformation of




Fig. 44. Effect of urea on Mg-ATPase activities of squid myosin and of acto-squid myosin

The ATPase activity of myosin alone(● ○) and that of actomyosin(▲ △) were assayed under the same conditions as in Fig. 38, except that various concentrations of urea from 0 M to 1.2 M were further added to the reaction mixture.




Fig. 45. Effect of urea on the Ca sensitivity of squid myosin Mg-ATPase

The ATPase activity of squid myosin was assayed in the same medium as in Fig. 38, and at the time indicated by the arrows, 1.1 M urea was added to the reaction mixture without changing other medium compositions. The same symbols as in Fig. 44 were also used.




Fig. 46. Recovery of Ca sensitivity of squid myosin Mg-ATPase by dilution of urea

The ATPase activity of squid myosin was assayed in 30 mM KCl, 20 mM Tris-maleate(pH 6.8), 2 mM MgCl_2 , 1 mM ATP, 1.1 M urea and either 0.1 mM CaCl_2 (●) or 0.5 mM EGTA (○). At the time indicated by the arrows, urea concentration in reaction mixture was decreased to 0.6 M without changing other medium compositions.

myosin thick-filaments induced by urea(40). It should be noted that at the same concentration of urea(1 M), both myosin Mg-ATPase and actin-activation of myosin ATPase increased in the absence of calcium.

Therefore, followings were suggested: when inhibitory effect of regulatory light chain on myosin ATPase was removed by urea, the inhibitory effect on interaction between actin and myosin was also removed, and resulted in increment of actin-activation in the absence of calcium.

Effect of urea on the Ca sensitivity of myosin Mg-ATPase was further studied by urea concentration jump method. As shown in Fig. 45, when the urea was added to reaction mixture to the final concentration of 1.1 M, at the time indicated by the arrow, Pi liberation in the absence of calcium immediately increased with no lag time. Then the reversibility of the effect of urea on Ca sensitivity of myosin was tested. The result was shown in Fig. 46. When the concentration of urea was diluted from 1.2 M to 0.6 M, Pi liberation in the absence of calcium immediately decreased and recovered Ca sensitivity.

It was, therefore, suggested that some conformational change in myosin molecule occurred by urea, which resulted in the loss of Ca sensitivity. However, this change may be small because Ca sensitivity lost was easily recovered by dilution of urea concentration.

2-(F)-f) Effect of tryptic digestion of squid myosin
on Ca sensitivity in its Mg-ATPase

The effect of tryptic digestion of squid myosin on its Ca sensitivity was studied. Squid myosin in 0.5 M KCl, 20 mM Tris-maleate(pH 7.5) was digested by trypsin in an amount of 1/500 for trypsin/myosin by weight at 10°C, and digestion was stopped by addition of 5 times weight of trypsin inhibitor. Change in Mg-ATPase activity was shown in Fig. 47, Mg-ATPase activity of myosin in the absence of calcium gradually increased with increasing the digestion time and resulted in loss of Ca sensitivity; Ca sensitivity of squid myosin was lost by the tryptic digestion for 60 - 90 min.

In addition, the Ca sensitivity of water-soluble fraction obtained by tryptic digestion was also measured. The digest of squid myosin was dialyzed against 0.05 M KCl, 20 mM Tris-HCl(pH 7.5) overnight and centrifuged at 100,000 x g for 60 min. The supernatant thus obtained was used for Mg-ATPase assay. As shown in Table III, water-soluble fraction obtained after 5 minutes digestion was still highly sensitive to calcium. However, when myosin was digested for longer time(30 - 45 min), Ca sensitivity of soluble fraction was decreased.

As shown in Fig. 48, it was also found that regulatory light chain(squid LC-2) was gradually digested away under this condition. The digestion of light chain seemed to




Fig. 47. Change in Mg-ATPase of squid myosin during tryptic digestion

Squid myosin was digested with trypsin at 10°C in an amount of 1/500 for trypsin/myosin by weight in the medium of 0.5 M KCl, 20 mM Tris-maleate(pH 7.5). The ATPase of squid myosin was assayed in the same medium as in Fig. 38, and the same symbols were used.




Fig. 48. Change in SDS gel electrophoretic patterns of squid myosin by tryptic digestion

Squid myosin was digested as described in Fig. 47, and the whole digest of myosin was applied on 10 % polyacrylamide gel containing 0.1 % SDS.

Table III. Ca sensitivity of water-soluble fractions obtained by tryptic digestion of myosin.

Water-soluble fraction of squid myosin digest was obtained by the following procedures; squid myosin digest obtained as described in Fig. 47 was dialyzed against 0.05 M KCl, 20 mM Tris-HCl(pH 7.5) and centrifuged at 100,000 x g for 60 min. The supernatant thus obtained was used as water-soluble fraction. ATPase was assayed as in Fig. 47, and Ca sensitivity was calculated as described in Fig. 31.

Digestion time (min)	Mg-ATPase ($\mu\text{mol Pi/min}\cdot\text{mg}$)		Ca sensitivity (%)
	+Ca	-Ca	
5	.189	.009	95
10	.201	.043	79
20	.211	.071	66
40	.214	.093	57




Fig. 49. Effect of intact squid LC-2 on the Ca sensitivity of Mg-ATPase of squid myosin digest

Squid myosin was digested by trypsin as described in Fig. 47, and approximately 10 % (w/w) of squid LC-2 was then added to myosin digest. The ATPase was assayed as in Fig. 38. (● ○), squid myosin digest alone, (▲ △), myosin digest with squid LC-2.

be proportional to loss of Ca sensitivity. These results also indicated that the regulatory light chain was essential for the development of Ca sensitivity of myosin alone.

Then, the effect of intact squid regulatory light chain(squid LC-2) on the Ca sensitivity of tryptic digest of squid myosin was studied. As shown in Fig. 49, Ca sensitivity of Mg-ATPase of squid myosin digest was unchanged even when the excess amount of squid LC-2 was added to it. These results strongly suggested that not only LC-2 molecule itself but also LC-2 binding site in squid myosin molecule was digested away by trypsin.

2-(F)-g) Ca sensitivity of scallop myosin alone and
loss of it by EDTA-treatment

It was studied to know that Mg-ATPase activity of scallop myosin alone was also sensitive to calcium. As shown in Fig. 50A, Mg-ATPase of scallop myosin as well as that of squid myosin were activated by the same concentrations(1 - 10 μ M) of free calcium as that of acto-scallop myosin were. However, it should be noted that scallop myosin showed a high ATPase activity than squid myosin did in the absence of calcium.

Then, using scallop desensitized myosin, the author studied the role of regulatory light chain in Ca sensitivity of myosin alone. Figure 50 shows that the Mg-ATPase activity of desensitized scallop myosin as well as that




Fig. 50. Calcium requirement for the activation of Mg-ATPases of scallop myosins and of rabbit acto-scallop myosins

ATPase was assayed under the conditions described in Fig. 37, and the same symbols as in Fig. 37 are used.

(A) Scallop untreated myosin, (B) scallop desensitized myosin, (C) scallop resensitized myosin with squid LC-2, and (D) scallop resensitized myosin with scallop EDTA-LC.

of acto-desensitized scallop myosin lost Ca sensitivity. The ATPase activity of scallop myosin in the presence of calcium(10 - 100 μ M free calcium) was unaffected by EDTA-treatment, whereas the activity in the absence of calcium(0.1 - 1 μ M free calcium) was greatly increased by the same treatment procedure, thus resulting in loss of Ca sensitivity. The Ca sensitivity of the ATPase activity of acto-scallop myosin was lost upon EDTA-treatment, but the ATPase activity in the presence of calcium was also reduced by the same treatment(B). Ca sensitivity of myosin ATPase and that of actomyosin ATPase were both recovered on addition of either squid LC-2(C) or scallop EDTA-LC(D). Moreover, the reduced ATPase activity of acto-desensitized myosin(B) was also recovered on addition of regulatory light chains.

Effects of regulatory light chain on the Ca sensitivity of myosin alone, and on that of actomyosin were then studied in detail. Various amounts of regulatory light chain were added to desensitized scallop myosin, and its ATPase activity was measured. As shown in Fig. 51, Mg-ATPase activity of desensitized scallop myosin in the presence of calcium was unaffected by addition of regulatory light chain, while it was strongly inhibited in the absence of calcium.

With respect to Mg-ATPase of actomyosin, ATPase activity of acto-desensitized myosin in the presence of




Fig. 51. Effect of squid LC-2 on the Mg-ATPase of desensitized scallop myosin and on that of actodesensitized myosin

The ATPase activity was assayed in a medium containing 30 mM KCl, 20 mM Tris-maleate(pH 6.8), 2 mM MgCl₂, 1 mM ATP and either 0.1 mM CaCl₂(closed) or 0.5 mM EGTA(open). (● ○), myosin ATPase, (▲ △), actomyosin ATPase.

calcium was elevated on addition of regulatory light chain, and ATPase activity in the absence of calcium was well inhibited by addition of regulatory light chain. The inhibition of ATPase activity of myosin in the absence of calcium and the elevation of actomyosin ATPase in the presence of calcium both reached maximum on addition of 7 % (w/w) of regulatory light chain to desensitized scallop myosin. This value corresponded to two mol of light chain per mol of desensitized myosin.

2-(F)-h) Effects of desensitization and resensitization
on superprecipitation ability of acto-
scallop myosin

Similar results were also obtained by measuring superprecipitation activity instead of ATPase activity. Figure 52A shows that acto-scallop myosin lost its superprecipitation ability as EDTA-treatment of scallop myosin progressed, even in the presence of calcium. Figure 52 (B and C) shows that the lost ability was recovered on addition of either squid LC-2 or scallop EDTA-LC. The two regulatory light chains were equally effective in the recovery of superprecipitation ability. The recovery was saturated when the amount of regulatory light chains added reached about 7 % of that of desensitized scallop myosin, which is approximately equivalent to a molar ratio of 1/1 for light chain/heavy chain. This value (7%)




Fig. 52. Effects of desensitization and resensitization on the superprecipitation of rabbit acto-scallop myosin

Superprecipitation was measured in a medium containing 30 mM KCl, 20 mM Tris-maleate(pH 6.8), 2 mM MgCl_2 , 0.1 mM CaCl_2 , and 0.5 mM ATP. The turbidity (absorbance) change of actomyosin suspension after addition of ATP was followed at 550 nm. (A) Acto-desensitized scallop myosin; numbers in the figure denote the numbers of repetitions of EDTA-treatment. (B and C) Acto-resensitized scallop myosin; numbers in the figure denote the amounts of squid LC-2(B) and of scallop EDTA-LC(C) added to desensitized scallop myosin(% w/w).

was also equal to that required for recovery of Ca sensitivity in the ATPase activity(Fig. 32).

2-(G) ATP concentration dependence of ATPase and
superprecipitation

2-(G)-a) ATP concentration dependence of superprecipitation and ATPase of squid and scallop
myosin Bs

In the previous section, Ca sensitivity of myosin alone and that of actomyosin were studied at a fixed ATP concentration of 1 mM. Then the author studied Ca sensitivity of them at various ATP concentrations.

Figure 53 shows the ATP concentration dependence of the superprecipitation activity of myosin B's from squid and scallop muscles. Superprecipitation occurred in the absence of calcium when the ATP concentration was very low(1 - 10 μ M ATP). Moreover, superprecipitation of both myosin B's was practically insensitive to calcium at 1 - 10 μ M ATP. The superprecipitation activity of both myosin B's, either in the absence or presence of calcium, responded in a biphasic manner to change in the ATP concentration, and the maximal activity was obtained at about 3 - 10 μ M ATP in the absence of calcium and at 100 - 300 μ M ATP in the presence of calcium. Figure 54 shows that the ATPase activities of scallop and squid myosin B's responded essentially in the same manner as their superprecipitation activities did(Fig. 53).




Fig. 53. ATP concentration dependence of the superprecipitation activities of myosin B's from squid and scallop muscles

Superprecipitation was measured in a medium containing 30 mM KCl, 20 mM Tris-maleate(pH 6.8), 2 mM $MgCl_2$, either 0.1 mM $CaCl_2$ (●), or 0.5 mM EGTA (○) and various concentrations of ATP (1 μ M - 1 mM). Pyruvate kinase (0.3 mg/ml) and phosphoenol pyruvate (1 mM) were added to maintain the ATP concentration. Superprecipitation activity was expressed in terms of $\Delta A \times 1/\tau_{1/2}$, where ΔA is the maximal increase in turbidity, and $\tau_{1/2}$ is the time required to reach the half-maximal increase. (A) Squid myosin B. (B) Scallop myosin B.




Fig. 54. ATP concentration dependence of Mg-ATPases of myosin B's from squid and scallop muscles

The ATPase activity of myosin B was assayed under the conditions described in Fig. 53, and the same symbols as in Fig. 53 are used. (A) Squid myosin B, and (B) scallop myosin B.

2-(G)-b) ATP concentration dependence of Mg-ATPase
of squid and scallop myosin

Figure 55 shows that Mg-ATPase activity of squid myosin and that of scallop myosin responded in a monophasic manner to change in the ATP concentration. It was also observed(see also Fig. 37, 50) that the Mg-ATPase activity of scallop myosin was higher than that of squid myosin in the absence of calcium at all ATP concentrations tested, and accordingly that Ca sensitivity at 1 μ M ATP was more obvious with squid myosin than with scallop myosin.

2-(G)-c) ATP concentration dependence of Mg-ATPase
activity of desensitized and resensitized
scallop myosin and of actomyosin

In order to study the effects of regulatory light chain on the ATP concentration dependence of myosin alone and on that of actomyosin, the author examined the ATP concentration dependence of desensitized scallop myosin and that of acto-desensitized scallop myosin as well as those of resensitized scallop myosin and of acto-resensitized myosin. As shown in Fig 56, the essential features of the ATP concentration dependence of both myosin ATPase and actomyosin ATPase were unaffected by desensitization and by resensitization: for example, as the ATP concentration increased, myosin ATPase increased in a monophasic manner whereas actomyosin ATPase did so in a biphasic manner.




Fig. 55. ATP concentration dependence of Mg-ATPases of myosins from squid and scallop muscles

The ATPase activity was assayed under the conditions described in Fig. 53, and the same symbols as in Fig. 53 are used. (A) Squid myosin, and (B) scallop myosin.




Fig. 56. Effects of desensitization and of resensitization on the ATP concentration dependence of Mg-ATPases of myosin and of actomyosin from scallop muscle

The ATPase activity was assayed under the conditions described in Fig. 53. (● ▲) 0.1 mM CaCl_2 , (○ Δ) 0.5 mM EGTA, and (● ○) myosin alone, (▲ Δ) actomyosin.

(A) Desensitized scallop myosin, (B) resensitized myosin with squid LC-2, and (C) resensitized myosin with scallop EDTA-LC.

It should be pointed out that resensitized myosin with squid LC-2 showed much lower ATPase activity than did that with scallop EDTA-LC, in the absence of calcium. These differences would arise not from differences in the properties of heavy chains but from differences in the properties of squid LC-2 and scallop EDTA-LC.

2-(H) Tension development of glycerinated muscle fiber
of scallop adductor muscle

2-(H)-a) Effect of EDTA-treatment of glycerinated
fibers on its Ca sensitivity in tension
development

In the previous study on the biochemical role of regulatory light chain, it was found that removal of two mol of EDTA-LC was required to desensitize the scallop myosin, and that removal of EDTA-LC remarkably reduced superprecipitation activity of acto-desensitized myosin.

Successively, the role of regulatory light chain was studied by measuring the physiological properties of tension development. The glycerinated muscle fiber from scallop adductor striated muscle was prepared by using the routine method for that from rabbit psoas muscle. The details were described in "Methods and materials" section.

In Fig. 57A, a typical tension development of scallop muscle fibers were shown. The tension developed




Fig. 57. ATP induced tension development in glycerinated fibers of scallop adductor muscle

A typical record of tension development in glycerinated muscle fibers of scallop was shown. Tension development in fibers of scallop was measured in a medium containing 0.1 M KCl, 20 mM imidazole-HCl (pH 6.8), 0.3 mM DTT, 5 mM MgCl_2 , 4 mM ATP, and either 2 mM EGTA(EGTA) or 0.2 mM CaCl_2 plus 0.1 mM EGTA(Ca) at 20°C. Untreated fibers(A) and EDTA-treated fibers for 20 min(B) were used.




Fig. 58. ATP induced tension development in glycerinated fibers of scallop adductor muscle in the presence and absence of calcium

Glycerinated fiber bundles (each approximately 300 μ thick and 5 mm long) were treated with 10 mM EDTA at 20°C for various periods. The extent of tension developed was measured at 20°C in a fiber bath containing 0.1 M KCl, 20 mM imidazole-HCl (pH 6.8), 0.3 mM DTT, 5 mM MgCl₂, 4 mM ATP, and 2 mM EGTA (○) or 0.2 mM CaCl₂ plus 0.1 mM EGTA (●).

only in the contracting solution, and the average tension developed in the presence of calcium was ca. 112 g/cm^2 . As shown in Fig. 57B, when the fibers were treated with EDTA solution for 15 min, as described in "Methods and materials", scallop muscle fibers became capable of developing tension in the absence of calcium. Moreover, it was observed that tension development in the presence of calcium was decreased by the same treatment.

In Fig. 58, tension development in the presence of calcium and that in the absence of calcium were both shown as a function of the duration of EDTA-treatment. When the fibers were treated with EDTA solution for 10 min, glycerinated fibers became capable of developing tension in the absence of calcium. The tension developed was nearly half that developed by untreated glycerinated fibers in the presence of calcium. However, it was also observed that tension developed in the presence of calcium was decreased to about three-fourth by the same treatment of fibers, and that the tension developed continued to decrease to as low as less than one-third that before the EDTA-treatment.

2-(H)-b) Relationship between EDTA-LC contents in
fibers and Ca sensitivity of tension
developed in the presence of calcium

In the polyacrylamide gel electrophoreses of glycerinated fibers in the presence of either urea or SDS,




Fig. 59. Effect of EDTA-treatment of glycerinated fibers of scallop on the EDTA-LC content, on the Ca sensitivity in tension development, and on the active (calcium) tension.

The EDTA-treatment was conducted at 20°C as described in Fig. 58. The regulatory light chain(EDTA-LC) content (○) was estimated by urea-gel electrophoresis. Ca sensitivity(△) and active(calcium) tension(□) were obtained from the data shown in Fig 58.

practically no change was detectable in the contents of heavy chain, actin, tropomyosin, paramyosin and SH-LC during EDTA-treatment. The only change detectable was in the EDTA-LC content. As shown in Fig. 59, the content of EDTA-LC decreased exponentially to about 15 % of the initial content, and the half-maximal decrease occurred at about 6 min of EDTA-treatment. Ca sensitivity was estimated from the data shown in Fig. 58, expressed in terms of $(1 - T_E/T_{Ca}) \times 100$, where T_E and T_{Ca} were tension developed in the absence and presence of calcium, respectively. The results indicated that Ca sensitivity of tension development decreased essentially in the same manner as the EDTA-LC content did, the half-maximal decrease occurring at approximately 6 min of the EDTA-treatment. Tension developed in the presence of calcium decreased exponentially, and the half-maximal decrease occurred at 15 min of EDTA-treatment.

2-(H)-c) Effect of regulatory light chain on Ca sensitivity and on tension development in the presence of calcium

Moreover, it was found that the lost ability of developing tension in the presence of calcium and the lost Ca sensitivity in tension development were both fully recovered. As shown in Fig. 60A and 60B, when the desensitized fibers treated with EDTA for 60 min were

soaked in a solution containing either scallop(A) or squid(B) regulatory light chain for recombination, fibers regained the ability to develop tension in the presence of calcium as much as about 70 - 90 % of that developed by untreated fibers, and they developed no tension in the absence of calcium. Therefore, recovery of active tension(tension in the presence of calcium) as well as that of Ca sensitivity were very good. In both cases, time required for the recombination of regulatory light chain to EDTA-treated muscle fibers was about 30 min.




Fig. 60. Resensitization of glycerinated fibers of scallop muscle by regulatory light chain from squid of scallop muscle

EDTA-treated fibers of scallop for 60 min at 20°C as described in Fig. 58, were soaked in a solution containing 0.1 M KCl, 20 mM imidazole(pH 6.8), 5 mM MgCl_2 , 4 mM ATP, 0.3 mM DTT, 2 mM EGTA and squid LC-2(A)(1.8 mg/ml) or scallop EDTA-LC(B)(1.8 mg/ml) for various durations. The tension developed in the presence(●) and absence (○) of calcium were measured as described in Fig. 57, and Ca sensitivity in tension developed were estimated as in Fig. 59.

IV DISCUSSION

Szent-Györgyi et al. made a great discovery that muscle contraction in scallop adductor muscle is regulated not by troponin-tropomyosin system but by myosin molecule. They concluded that all of the moluscan muscles are regulated only by myosin-linked Ca regulatory system(6). At present time, there are few report on the Ca regulation of molluscan muscle, except those of Szent-Györgyi and his group.

The author tried to know detailed mechanism of Ca regulation in molluscan muscle, especially in squid and scallop muscles, which are both categorized in a group containing only myosin-linked regulation by Lehman and Szent-Györgyi. However, applying the competitive test(6, 24) for squid myosin B, it is found that, being conflicted with the results of Lehman et al., myosin-linked and actin-linked Ca regulations are both suggested in squid myosin B. Since two Ca regulatory systems were assumed in squid muscle, the author tried to ascertain whether squid muscle truly contains two regulation systems or not.

At first, actin-linked Ca regulation in squid muscle was studied. Ca sensitivity in Mg-ATPase of actomyosin constituted from squid actin and rabbit myosin were measured. The author then found that actomyosin using purified squid actin is not Ca sensitive, while actomyosin constituted by using crude squid actin is sensitive to calcium. These results suggested that squid crude actin contains protein

components conferring Ca sensitivity to actomyosin. Accordingly, a trial for isolation of native tropomyosin was made and biologically active native tropomyosin and troponin are both obtained from squid muscle. Squid native tropomyosin, thus obtained, confers Ca sensitivity to rabbit actomyosin as rabbit native tropomyosin does. The functional identity of squid native tropomyosin with rabbit native tropomyosin is suggested, though squid native tropomyosin is less effective than that of rabbit native tropomyosin, i.e. inhibition of actomyosin Mg-ATPase activity by squid native tropomyosin in the absence of calcium is relatively small. However, squid native tropomyosin as well as rabbit one was found to be a complex of troponin and tropomyosin, which were both successfully separated and isolated. Squid troponin also contains three subunits(52,000, 28,000 and 24,000 daltons)(Fig. 8). The similar 52,000 daltons component has been widely observed in troponin preparations from various arthropod (lobster, crayfish and Limulus)(27, 31, 41, 42). Moreover, arthropod occupies a just upper position of mollusca on phylogenetic tree. These fact may relate to the speculation on the evolution of regulatory protein system as Lehman et al. have proposed(6). Squid troponin fraction containing 52,000 daltons component was found to be capable of binding with tropomyosin. Since 52,000 dalton component in lobster troponin is corresponded to be TN-T in rabbit

troponin, which is tropomyosin binding component, this 52,000 dalton component in squid troponin also corresponds to TN-T of mammalian muscle. It was found that squid troponin is present in a form of complex of three components under the physiological conditions, and the complex of 3 subunits is easily dissociated by low concentration of urea(0.1 - 0.2 M). These results suggested that there are weak bindings among 3 subunits of squid troponin. It was also found that squid native tropomyosin as well as squid troponin can bind only a small amount of calcium (0.2 mol of Ca/ 1×10^5 g protein). These two prospects may be included in the reason for the characteristic properties of squid native tropomyosin in conferring Ca sensitivity.

Myosin-linked Ca regulation, especially in scallop adductor muscle, has been studied in the laboratory of Szent-Györgyi. The author's present results was essentially the same as their results in the following respect that actomyosin reconstituted from squid myosin and rabbit purified actin is sensitive to calcium in ATPase and in superprecipitation. These results clearly show that squid muscle contains myosin-linked regulation system besides actin-linked regulation system as described above, and two regulatory systems are found to be functional, if tested separately.

There are a few reports suggesting the presence of

both myosin-linked and actin-linked regulations, e.g., in insect flight and leg muscles(6), in earthworm body wall(6), and in Physarum polycephalum(43). However, there have been no reports presenting the direct evidences, i.e. isolation and characterization of both systems. Therefore, the present report on two Ca regulation systems in squid mantle muscle may give the first evidence.

Naturally, the question may arise as to which of the two regulatory systems is predominantly operating in squid muscle. Fortunately, these two systems are distinguished by their Ca and Sr concentration requirements for the activation of Mg-ATPase activity, i.e. actin-linked regulation requires lower Ca and Sr concentrations for its activation of Mg-ATPase than myosin-linked regulation system does. Kendrick-Jones et al.(2) also reported that scallop myosin B Mg-ATPase(myosin-linked) is fully activated only at higher Ca concentration than rabbit skeletal myosin B Mg-ATPase is. The Ca requirement for activation of squid myosin B, which indeed contained both systems, was found to be identical to that for activation of myosin-linked system. It should also be noted that the Sr requirement of squid crude acto-rabbit myosin(actin-linked) is identical to that of rabbit myosin B. Therefore, it seems that myosin-linked system rather than actin-linked system is predominantly operating. It was further supported by the fact that addition of large amount of native tropo-

myosin from squid or rabbit does not shift Ca requirement of squid myosin B from that of myosin-linked system to that of actin-linked system.

A further study on the myosin-linked Ca regulation in squid muscle was developed. In scallop muscle, it is well known that regulatory subunit of myosin is one type of light chain named EDTA-LC(4), which is able to be isolated by EDTA-treatment of myosin. The author tried to isolate regulatory light chain from squid myosin. As already described, none of light chain is released from myosin by treating with EDTA, but they are released from myosin by treating with urea. A mixture of light chains released are successfully separated into each light chain by using DEAE-Sephadex column chromatography. One of light chain named LC-2 was suggested to be calcium binding subunit, judging from the results that calcium induces spectral change of LC-2 and calcium induces mobility change of LC-2 in disc gel electrophoresis. The calcium binding ability of squid LC-2 was measured by equilibrium dialysis method, and isolated LC-2 was found to bind small amount of calcium(0.4 - 0.5 mol of Ca/mol of LC-2), this value is somewhat smaller than that obtained with intact squid myosin(2 mol of Ca/mol of myosin, or 1 mol of Ca/mol of LC-2). These results suggest that when LC-2 is released from myosin, its conformation, probably in the calcium binding site, may be changed and result in decrease

in its calcium binding ability. It should be pointed out that squid LC-2 has better calcium binding ability than scallop EDTA-LC(0.15 mol of Ca/mol of EDTA-LC) has.

In order to test the biological function of squid regulatory light chain(LC-2), scallop myosin was used, because squid myosin could not be desensitized by EDTA-treatment or by any other procedures. In the process of desensitization of scallop myosin, the author found the conflicting results with that of Szent-Györgyi et al. For instance, they reported that among four mol of light chains(2 mol of SH-LC, 1 mol of EDTA-LC and 1 mol of DTNB-LC) in scallop myosin, removal of 1 mol of EDTA-LC is enough to desensitize scallop myosin. Based on these results, they have proposed the functional cooperativity between two heads of myosin or two mol of light chains in the molecular mechanism of Ca regulation. On the other hand, the present author found that removal of 2 mol of EDTA-LC is required to desensitize scallop myosin, i.e. Ca sensitivity is still remained even if 1 mol of EDTA-LC is removed from myosin. Moreover, it was found that two mol of EDTA-LC is easily removed by EDTA-treatment, and that DTNB-LC is not able to be distinguished from EDTA-LC in this experiment. These results also suggested the absence of functional cooperativity of light chains in the Ca regulation mechanism.

Desensitized scallop myosin was used for the functional

test of regulatory light chain of squid, in terms of Mg-ATPase activity of reconstituted actomyosin. Squid LC-2 is capable of binding to desensitized scallop myosin and confers Ca sensitivity to it, the binding of LC-2 and Ca sensitivity recovered are both maximum at 2 mol of squid LC-2 addition per mol of myosin. Therefore, squid LC-2 is equivalent to scallop EDTA-LC and is not distinguished from EDTA-LC in its function. Furthermore, the bound squid LC-2 to desensitized myosin is easily removed by EDTA-treatment of this myosin. Therefore, it seems likely that the difficulty to remove light chain from squid myosin by EDTA-treatment is not caused by the properties of myosin regulatory light chain but by that of myosin heavy chain.

Recently, Asada et al. reported that Mg-ATPase of clam foot muscle myosin alone is sensitive to calcium as well as actomyosin(39). The author studied whether this property of clam foot myosin was restricted to clam or general in all molluscan myosin. As a result, squid myosin Mg-ATPase was found to be clearly activated by the low concentration of calcium ions between 1 μ M and 10 μ M. Moreover, the Ca requirement for activation of myosin Mg-ATPase was identical to that required for activation of actomyosin Mg-ATPase. It is, therefore, concluded that Mg-ATPase of squid myosin alone is sensitive to calcium. Changes in temperature and pH for Mg-ATPase

assay unaffected Ca sensitivity of myosin alone. However, KCl and urea for Mg-ATPase assay affected Ca sensitivity of myosin alone. Ca sensitivity of myosin Mg-ATPase is lost at KCl concentration higher than 0.4 M. On the other hand, actin activation of Mg-ATPase of myosin in the presence of calcium is lost at 0.2 M KCl. Therefore, the effects of KCl on Ca sensitivity of myosin Mg-ATPase and on actin-activation of it is able to be distinguished.

In the presence of 1 - 1.2 M urea, myosin Mg-ATPase is converted to be insensitive to calcium, but actin-activation of it still occurs. At the same time, it was found that actin-activation of myosin Mg-ATPase in the absence of calcium increased, resulting in the loss of Ca sensitivity its actin-activation. These results suggested that Ca sensitivity of myosin alone may be always coupled with actin-activation. It appears, therefore, that a primary role of regulatory light chain is to regulate (inhibit) myosin Mg-ATPase, and that if inhibition of myosin Mg-ATPase is removed, binding of myosin to actin occurs.

Scallop myosin Mg-ATPase is sensitive to calcium as squid myosin Mg-ATPase is. Ca sensitivity in its Mg-ATPase is lost by removal of all of EDTA-LC from scallop myosin. Moreover, Ca sensitivity of actomyosin measured by Mg-ATPase and by superprecipitation are both lost by removal of EDTA-LC from its myosin. It was also

found that actin-activation of desensitized scallop myosin Mg-ATPase and superprecipitation of acto-desensitized scallop myosin are remarkably reduced even in the presence of calcium and they are both recovered on addition of either squid or scallop regulatory light chain. Therefore, it is suggested that another function of regulatory light chain is to maintain regular binding between myosin and actin in the presence of calcium. This idea is further confirmed by measuring the change in tension development of glycerinated muscle fibers of scallop during EDTA-treatment. Ca sensitivity in tension development of glycerinated fibers are lost by removal of EDTA-LC from them, while the tension development in the presence of calcium decreased with decreasing the contents of EDTA-LC. Moreover, Ca sensitivity and tension development in the presence of calcium are both recovered on addition of either squid or scallop regulatory light chain.

From these results, the author speculates that when light chain is removed from myosin, Ca sensitivity of myosin as well as actomyosin are concomitantly lost, and that removal of light chain may induce conformational change in myosin molecule, and presumably also in thick-filament formation. This change may lead a decrease in the tension development and in the actin-activation of myosin ATPase.

V SUMMARY

Squid mantle myosin B was Ca sensitive, and its Ca sensitivity was unaffected by addition of large amount of rabbit skeletal myosin or rabbit skeletal F-actin. Therefore, it was suggested that both myosin-linked and actin-linked Ca regulatory systems were present in squid mantle muscle.

1 Actin-linked regulatory system

1-(a) Squid mantle actin was prepared by the method of Spudich and Watt. Hybrid actomyosin reconstituted by using the purified squid actin with skeletal myosin showed no Ca sensitivity in Mg-ATPase assay, whereas that reconstituted using crude squid actin showed marked Ca sensitivity. The crude squid actin contained four protein components which were capable of associating with F-actin in 0.1 M KCl, 1 mM MgCl_2 and 20 mM Tris-maleate(pH 7.5).

1-(b) Native tropomyosin was prepared from squid mantle muscle, and it conferred Ca sensitivity to skeletal actomyosin as well as to hybrid actomyosin reconstituted from squid actin and skeletal myosin, but it is less effective than skeletal native tropomyosin in conferring Ca sensitivity.

1-(c) Squid native tropomyosin was separated into

troponin and tropomyosin fractions by placing it in 0.4 M LiCl at pH 4.7. The troponin fraction was further purified by DEAE-cellulose chromatography. Squid troponin thus obtained was different in mobility in SDS gel electrophoresis from rabbit skeletal troponin; three bands of squid troponin corresponded to molecular weights of 52,000, 28,000 and 24,000 daltons. It could confer Ca sensitivity in the presence of tropomyosin on skeletal actomyosin as well as on hybrid actomyosin reconstituted from squid actin and skeletal myosin. It was also found that troponin-tropomyosin complex using squid troponin was less effective than that using skeletal troponin.

Squid troponin could not bind to F-actin, whereas it could bind to F-actin in the presence of tropomyosin from squid or skeletal muscle.

2 Myosin-linked regulatory system

2-(A) Ca sensitive myosin in squid mantle muscle

2-(A)-(a) Squid myosin was prepared from the mantle muscle, which showed a heavy chain component and two light chain components in SDS gel electrophoretic pattern: the molecular weights of the latter two were 17,000 and 15,000 daltons. Actomyosin reconstituted from squid myosin and skeletal(or squid) actin showed Ca sensitivity in super-precipitation and in Mg-ATPase assay. EDTA-treatment had no effect on Ca sensitivity of squid myosin.

2-(A)-(b) Squid myosin B and two hybrid actomyosins were compared as regards Ca and Sr requirements in their Mg-ATPase activities. The result predicted that myosin-linked regulatory system was predominantly operating in squid myosin B. Squid myosin B required higher Ca or Sr concentration for Mg-ATPase activity: half-maximal activation of Mg-ATPase was obtained at 0.8 μ M calcium ions and at 28 μ M strontium ions with skeletal myosin B, and at 2.5 μ M calcium ions and at 140 μ M strontium ions with squid myosin B.

2-(B) Role of regulatory light chain

2-(B)-(a) The author was able to separate two types of light chain by five steps of procedure, yielding LC-1(17,000 daltons) and LC-2(15,000 daltons). It was found that squid mantle LC-1 and LC-2 function exactly like SH-LC and EDTA-LC of scallop adductor myosin, respectively.

2-(B)-(b) Calcium or strontium ion was found to induce changes in the UV absorption spectrum of scallop adductor EDTA-LC, although the apparent binding constants estimated from the difference spectrum were too low to account for the Ca sensitivity of scallop actomyosin Mg-ATPase. The divalent cations also induced changes in the UV absorption spectrum of squid LC-2, and the apparent binding constants estimated from the difference spectrum

were sufficiently high($1.5 \times 10^5/\text{M}$ for Ca binding, and $1.6 \times 10^3/\text{M}$ for Sr binding) to account for the Ca and Sr sensitivities of squid myosin B ATPase.

2-(B)-(c) In contrast with the results of Kendrick-Jones et al., the author showed that removal of two mol of EDTA-LC from scallop myosin was required to desensitize actomyosin Mg-ATPase. Desensitized scallop myosin was capable of regaining its full content of EDTA-LC (two mol) as well as its full recovery of Ca sensitivity of actomyosin Mg-ATPase. It was also found that as regards combining with, and conferring Ca sensitivity on the desensitized scallop myosin, squid LC-2 could effectively replace scallop EDTA-LC.

2-(B)-(d) Calcium binding to the regulatory light chain was studied by an equilibrium dialysis method. It was found that calcium binding ability of regulatory light chain was decreased from 1 mol of Ca/mol of bound light chain to myosin to 0.15 - 0.4 mol of Ca/mol of isolated light chain. The calcium binding of isolated light chain was reduced by addition of magnesium ion. Moreover, calcium binding ability of regulatory light chain was fully recovered by recombination with myosin molecule.

2-(B)-(e) The Mg-ATPase activity of squid myosin alone was sensitive to calcium. Changes in temperature(5°C - 30°C) and pH(6.0 - 9.5) for Mg-ATPase assay unaffected Ca sensitivity of squid myosin. The Ca sensitivity of myosin alone was not observed under the conditions, where the KCl concentration was higher than 0.4 M or 1.2 M urea was present in the reaction medium. Moreover, actin-activation of myosin Mg-ATPase in the presence of calcium was lost at the KCl concentration higher than 0.2 M. Therefore, effects of KCl on Ca sensitivity of myosin alone and on actin-activation were distinguishable. Tryptic digestion of squid myosin also decreased Ca sensitivity with digestion of LC-2.

2-(B)-(f) Scallop myosin was also sensitive to calcium, and its Ca sensitivity was lost by removal of EDTA-LC. On addition of squid LC-2 or scallop EDTA-LC, Ca sensitivity of desensitized scallop myosin was recovered. Furthermore, removal of EDTA-LC from scallop myosin resulted in reducing the superprecipitation and Mg-ATPase activities of actomyosin, but they were recovered on addition of regulatory light chains from squid or scallop.

2-(B)-(g) Removal of all of the regulatory light chain(EDTA-LC) of myosin in glycerinated fibers of scallop adductor striated muscle resulted in an immediate

loss of Ca sensitivity in tension development and a concomitant decrease in tension development in the presence of calcium. These two properties of glycerinated muscle fibers were recovered by soaking the fibers in a solution containing regulatory light chain from squid or scallop.

From the results described above, the author revealed that squid mantle muscle contained not only myosin-linked but also actin-linked regulation system, although the myosin-linked regulation rather than actin-linked regulation is effectively operating in squid myosin B. It was further speculated that a primary role of regulatory light chain of squid or scallop myosin is to inhibit myosin Mg-ATPase, leading inhibition of actin-myosin binding in the absence of calcium. Moreover, the regulatory light chain was found to be essential for the regular binding of myosin to actin in the presence of calcium, which was considered to be basic reaction in muscle contraction mechanism.

VI ACKNOWLEDGEMENTS

I express my sincere thanks to Professor S. Watanabe and to Dr. Arai for their helpfull advices and cencouragements throughout the period of this investigation.

I am grateful to Professors M. Akiba and T. Yasui and to Dr. Shibata for their critical reading of the manuscript.

To Mrs. K. Ishida and B. Yoshitomi, I wish to express gratitude for their technical assistances.

Finally my deepest thanks go to my wife Yoshiko for her enduring encouragement.

VII REFERENCES

- 1) S. Ebashi & M. Endo; Prog. Biophys. Mol. Biol.,
18, 123 - 131, (1968)
- 2) J. Kendrick-Jones, W. Lehman & A.G. Szent-Györgyi;
J. Mol. Biol., 54, 313 - 326, (1970)
- 3) S. Ebashi & F. Ebashi; J. Biochem., 55, 604 - 613,
(1964)
- 4) A.G. Szent-Györgyi, E.M. Szent-Kiralyi & J. Kendrick-
Jones, J. Mol. Biol., 74, 179 - 203, (1973)
- 5) J. Kendrick-Jones; Nature, 249, 631 - 634, (1974)
- 6) W. Lehman & A.G. Szent-Györgyi; J. Gen. Physiol.,
66, 1 - 30, (1975)
- 7) S. Ebashi; J. Biochem., 79, 229 - 231, (1975)
- 8) J.A. Spudich & S. Watt; J. Biol. Chem., 246, 4866 -
4871, (1971)
- 9) S. Ebashi, A. Kodama & F. Ebashi; J. Biochem.,
64, 465 - 477, (1968)
- 10) K. Konno; J. Biochem., 83, 1431 - 1440, (1978)
- 11) J.C. Holt & S. Lowey; Biochemistry, 14, 4600 - 4609,
(1975)
- 12) W.T. Perrie, L.B. Smillie & S.V. Perry; Biochem. J.,
135, 151 - 164, (1973)
- 13) J.J. Connell; Biochem. J., 58, 360 - 367, (1954)
- 14) A. Szent-Györgyi; Biol. Bull., 90, 140 - 149, (1949)
- 15) C.H. Fiske & Y. Subbarow; J. Biol. Chem., 66,
375 - 400, (1925)

- 16) A.M. Reynard, L.F. Haas, D.D. Jacobsen, & P.D. Boyer;
J. Biol. Chem., 236, 2277 - 2291, (1961)
- 17) Y. Ogawa; J. Biochem., 64, 255 - 260, (1968)
- 18) K. Weber & M. Osborn; J. Biol. Chem., 244, 4406 -
4412, (1969)
- 19) B.J. Davis; Ann. N.Y. Acad. Sci., 121, 404 - 427,
(1964)
- 20) J.D. Potter & J. Gergely; Biochemistry, 13, 2697 -
2703, (1974)
- 21) H. Kuwayama & K. Yagi; J. Biochem., 82, 25 - 33,
(1977)
- 22) A.G. Gornall, C.T. Bardwill & M.M. David; J. Biol.
Chem., 177, 751 - 766, (1949)
- 23) R.M. Simmons & A.G. Szent-Györgyi; Nature, 273,
62 - 64, (1978)
- 24) W. Lehman, J. Kendrick-Jones & A.G. Szent-Györgyi;
Cold Spring Harbor Symp. Quant. Biol., 37, 319 -
330, (1972)
- 25) T. Tsuchiya, T. Kaneko & J.J. Matsumoto; J. Biochem.,
83, 1191 - 1193, (1978)
- 26) A. Goldberg & W. Lehman; Biochem. J., 171, 413 - 418,
(1978)
- 27) K. Konno, K. Arai & S. Watanabe; J. Biochem., 82,
931 - 938, (1977)
- 28) S. Ebashi, T. Wakabayashi, F. Ebashi; J. Biochem.,
69, 441 - 445, (1971)

- 29) M.L. Greaser & J. Gergely; J. Biol. Chem., 246,
4226 - 4234,(1971)
- 30) J.M. Wilkinson, S.V. Perry, H.A. Cole & I.P. Trayer;
Biochem. J., 127, 215 - 221,(1972)
- 31) J.M. Regenstien & A.G. Szent-Györgyi; Biochemistry,
14, 917 - 926,(1975)
- 32) J. Kendrick-Jones, E.M. Szent-Kiralyi & A.G. Szent-
Györgyi; J. Mol. Biol., 104, 747 - 775(1976)
- 33) T. Tsuchiya, N. Yamada & J.J. Matsumoto; Bull. Japan
Soc. Sci. Fish., 44, 175 - 179,(1978)
- 34) T. Tsuchiya, H. Suzuki & J.J. Matsumoto; Bull. Japan
Soc. Sci. Fish., 43, 877 - 884,(1977)
- 35) M. Ikebe, H. Onishi & S. Watanabe; J. Biochem., 82,
299 - 302,(1977)
- 36) J. Gazith, S. Himmelfarb & W.F. Harrington; J. Biol.
Chem., 245, 15 - 22,(1970)
- 37) J.F. Head, R.A. Weeks & S.V. Perry; Biochem. J.,
161, 465 - 471,(1977)
- 38) M. Yazawa & K. Yagi; J. Biochem., 82, 287 - 289,(1977)
- 39) T. Asada, G. Ashiba & S. Watanabe; J. Biochem.,
85, 1543 - 1546,(1979)
- 40) I. Kimura, K. Arai, K. Takahashi & S. Watanabe;
J. Biochem., 88, 1703 - 1713,(1980)
- 41) G. Benzonana, L. Kohler & E.A. Stein; Biochem. Biophys.
Acta, 368, 247 - 258,(1974)
- 42) W. Lehman, J.M. Regenstien & A.L. Ranson; Biochem.
Biophys. Acta., 434, 215 - 228,(1976)

- 43) V.T. Nachimias & A. Asch; Biochemistry, 15, 4273 -
4278, (1976)