




HOKKAIDO UNIVERSITY

Title	Plant symbiotic microorganisms in acid sulfate soil: significance in the growth of pioneer plants
Author(s)	Maki, Takanori; Nomachi, Miya; Yoshida, Shigekata et al.
Citation	Plant and Soil, 310(1-2), 55-65 https://doi.org/10.1007/s11104-008-9628-y
Issue Date	2008-09
Doc URL	https://hdl.handle.net/2115/38543
Rights	The original publication is available at www.springerlink.com
Type	journal article
File Information	PLS0maki_etal.pdf



Title: Plant symbiotic microorganisms in acid sulfate soil: significance in the growth of pioneer plants

Authors: Takanori Maki¹, Miya Nomachi¹, Shigekata Yoshida¹ and Tatsuhiko Ezawa^{2*}

Address: ¹Graduate School of Bioagricultural Science, Nagoya University, Togo-cho, Aichi 470-0151 Japan and ²Graduate School of Agriculture, Hokkaido University, Sapporo, Hokkaido 060-8589 Japan

***Author for correspondence:**

Tatsuhiko Ezawa

Graduate School of Agriculture, Hokkaido University, Sapporo, Hokkaido 060-8589 Japan

(tel +81-11-706-3845; fax +81-11-706-3845; e-mail tatsu@res.agr.hokudai.ac.jp).

No. of tables: 2

No. of figures: 6

No. of pages: 2

Key words

Acid sulfate soil, arbuscular mycorrhizal fungi, nodule bacteria, pioneer plants.

Abstract

Acid sulfate soil is generated by chemical and microbial oxidization of sulfide-rich minerals/sediments. Although revegetation of the soil is difficult due to low-pH and poor nutrient availability, pioneer plants may adapt to such an extreme environment via associating with mycorrhizal fungi and/or N-fixing bacteria for acquisition of mineral nutrients. In this study, an abandoned quarry in which acid sulfate soil was found was chosen to investigate the influence of soil acidity on the levels of colonization by the microsymbionts, the identities of the microsymbionts that associated with pioneer plants and the dependency of pioneer plants on the microsymbionts. The levels of arbuscular mycorrhizal (AM) colonization in pioneer grass, forbs and legume shrubs grown in the field were assessed, and no significant decline in the levels with an increase in soil acidity was observed. Most of the legume shrubs formed root nodules. Several AM fungi and bradyrhizobia were cultured from the rhizosphere soils of pioneer plants grown in the quarry and identified based on the sequences of the small subunit ribosomal RNA genes. Pot experiments revealed that the microsymbionts isolated from the field significantly promoted the growths of pioneer grasses and legume shrubs in acid sulfate soil at pH 3.4.

21 These results suggest that plant-microbial symbiotic associations play significant roles in
22 the growth of pioneer plants in acid sulfate soil.

24 **Introduction**

26 Acid sulfate soil is widely distributed in volcanic and coastal areas in the world. The soil is
27 formed by chemical and microbial oxidization of sulfide-rich minerals/sediments, in most
28 cases pyrite (FeS_2) that are exposed by large-scale land development, reclamation and
29 quarrying. As a result of oxidation, a large amount of sulfuric acid is generated, thus the soil
30 shows extremely low pH. Recently, the environmental impacts of acid sulfate soil have
31 become a major concern. Drainage from acid sulfate soil causes serious damage to marine
32 ecosystems via acidification, metal contamination, deoxygenation and iron precipitation
33 (Powell and Martens 2005). Acidification and metal contamination of freshwater caused by
34 the soil threaten not only terrestrial ecosystems but also human health (Appleyard et al.
35 2004). Revegetation of acid sulfate soil is one way of reducing the environmental impacts
36 of the soil but is difficult due to strong acidity, high concentrations of toxic elements e.g. Al
37 and Mn and deficiencies in the essential nutrients. In particular, P-deficiency often limits
38 plant growth in the soil (Do et al. 2004).

39 It is considered that pioneer plants that are characterized as primary vegetation in
40 disturbed areas may have developed diverse strategies to adapt to such extreme

environments, although their adaptation mechanisms are largely unknown. Nutrient acquisition via symbiotic associations with soil microorganisms is likely to be one of the strategies of pioneer plants. Arbuscular mycorrhizal (AM) fungi form mutualistic associations with about 80% of land plants and improve phosphorus (P) nutrition of the host plants via enhanced P-uptake through extensive hyphal networks in soil (Smith and Read 1997). It is, therefore, expected that AM associations may play significant roles in the primary vegetation of acid sulfate soil. On the other hand, Allen (1991) suggested that the first plant colonizers after disturbance would be non-mycotrophic or at least facultative mycotrophic. This idea, however, contradicts the fact that ectomycorrhizal formation on *Salix reinii*, a pioneer tree species, was essential in facilitating seedling establishment of successional plant species in the early successional volcanic desert in Mt. Fuji (Nara and Hogetsu 2004). Given the fact that P-deficiency is a major limiting factor for plant growth in acid sulfate soil, it is worth evaluating the dependency of pioneer plants on AM fungi in the soil for further understanding of the roles of mycorrhiza in disturbed ecosystems.

Leguminous plants benefit not only from AM symbiosis but also from the symbiotic nitrogen (N)-fixing associations with rhizobia. Extensive studies have demonstrated that the efficiency of N-fixation in legume-rhizobial associations is promoted by AM fungal colonization (Barea et al. 1992), and this might be mainly due to the high-requirement of P in the N-fixing process (Bohloul et al. 1992). The effectiveness of plant-AM fungi-rhizobia tripartite associations has been demonstrated in the management of

desertified ecosystems in the Mediterranean semi-arid region (Herrera et al. 1993; Requena et al. 1997; Requena et al. 2001). According to these observations, it is likely that the tripartite associations are of significance in the colonization of legume pioneers in acid sulfate soil.

It is likely that soil acidity has negative impacts on the formation and effectiveness of the symbiotic associations. Commercial *Bradyrhizobium* sp. that showed slower growth at lower pH *in vitro* could form nodules in acidic soil but did not fix appreciable N (Cline and Senwo 1993). The colonization, spore germination and the growth of germ tubes of an AM fungus that occurred in non-acidic soil were significantly reduced in acidic soil (Porter et al. 1987). The colonization levels of indigenous AM fungi on *Plantago lanceolata* in an acidic soil were significantly lower than those in limestone (non-acidic) soil, and the hyphal biomass of the fungi in the acidic soil was much smaller than that in the limestone soil (van Aarle et al. 2003). Based on these observations, it is hypothesized that the colonization of the symbiotic microorganisms on the host plants may be dramatically suppressed in extremely acidic soil. However, the ecology/physiology or even the presence of the symbiotic microorganisms in extremely acidic soil has not yet been reported.

In this study, an abandoned quarry located in the middle of Honshu Isl., Japan was chosen as a model field for studying plant-microbial associations in the pioneer vegetation of acid sulfate soil. The objectives of the present study were to investigate: i) whether the colonization of the microsymbionts on pioneer plants is suppressed with an increase in soil

acidity, ii) the origin and identity of the microorganisms that associate with pioneer plants under extremely acidic conditions and iii) whether the indigenous microsymbionts contribute to the growth of pioneer plants in acid sulfate soil. For these purposes, the levels of colonization in naturally occurring pioneer plants in the field were first assessed with respect to rhizosphere soil pH. Then AM fungi and rhizobia were isolated from the rhizosphere of the pioneer plants grown in the field and surrounding forest and identified, and the effectiveness on the growth of pioneer plants were evaluated by pot experiments.

Materials and methods

Sampling site

The experimental field is located in hilly area of Hazu-cho, Aichi prefecture, Japan (150-200 m altitude, 34°47'N, 137°07'E) where the annual mean temperature and rainfall are 16°C and 1,600 mm, respectively, and is surrounded by evergreen broad-leaved forest dominated by *Quercus glauca* Thunberg, *Q. myrsinaefolia* Blume and *Castanopsis cuspidata* Schottky. The understory of the forest is dominated by *Sasa nipponica* Makino et Shibata and *Dryopteris hondoensis* Koidz. The field had been quarried for a long time so that the bottom of the field (ca. 10 ha) is 20-100 m below the top of the surrounding hills and is flat except for the two ponds (2-4 m depth). The field has not been severely disturbed

since 1999. A thin silt/sand layer (5-15 cm in depth) is formed on sulfur-rich gravel/base rock, and the most of plant roots grow within this layer. *Alnus firma* Sieb. et Zucc. (Betulaceae) (0.1 to 3 m height) is observed around the two ponds and along the border between the quarry and surrounding forest (Fig. 1). The rest of the area is not fully covered with vegetation, but scattered distributions of herbaceous plants, *Miscunthus sinensis* Andresson (Gramineae), *Solidago altissima* L. (Asteraceae) and *Artemisia princeps* L. (Asteraceae), and leguminous shrub, *Lespedeza cyrtobotrya* L. (Leguminosae) are observed. *M. sinensis* and *L. cyrtobotrya* dominate in the site, although the former is observed more frequently than the latter.

Root and soil sampling

The roots and rhizosphere soils were collected both from the quarry and surrounding forest yearly from *M. sinensis*, *L. cyrtobotrya*, *S. altissima* and *A. princeps* in June from 2000 to 2002 as shown in Fig. 1. The roots were collected by two core samplers (100 mL in vol) from each plant, and then 2 - 3 kg rhizosphere soils were collected underneath the plants (top 5 - 10 cm depth, 30 - 40 cm in diam). Small parts of the soil samples were air-dried, crushed and passed through a 2 mm sieve for chemical analyses. The rest of the soils were stored at 7°C for the isolation of symbiotic microorganisms. Non-rhizospheric acid sulfate soil was collected as a culture medium for pot experiments from a sand pile in the quarry on

which no vegetation was observed (Fig. 1).

The root samples collected by the two core samplers were combined, washed with tap water, cut into 1 cm pieces, cleared with 10% KOH (w/v) and stained with 0.05% Trypan blue in lactoglycerol. The percentage of mycorrhizal colonization was estimated by the grid-line intersect method (Giovannetti and Mosse 1980). Soil pH (H₂O) was measured at 1:2.5 soil:water ratio, and pH (H₂O₂) was measured after oxidation by boiling in neutralized 30% H₂O₂. Total carbon (C) and nitrogen (N) contents were analyzed by a CHN analyzer (Perkin-Elmer, Yokohama, Japan). Available (Truog)-P was measured by the vanado-molybdate method after extraction with 0.001 M H₂SO₄ at ratio of 1:200 (w/v) (Truog 1930).

Isolation of microsymbionts

AM fungi in the rhizosphere of the pioneer plants were first proliferated by the soil trap culture (Brundrett et al. 1999). A half liter of the rhizosphere soils collected in July 2000 and 2002 was layered between autoclaved river sand in a 4 L plastic pot, and *M. sinensis*, *Arachis hypogaea* L. and *Sorghum bicolor* L. were sown together in the same pot. The plants were fertilized with 0.25 g L⁻¹ 20-5-30 PETERS liquid fertilizer (W. R. Grace & Co., Forgelsville, PA) in sufficient amounts until the solution flowed out from the drain twice or three times a week and were grown in a greenhouse (20 - 35°C). After four months, the

141 plants and media were dried in the greenhouse, and the roots and media were stored in
142 plastic bags at room temperature. The cultures were maintained yearly by the sand culture
143 system using the same host plants. The AM fungal spores were extracted by wet sieving,
144 and the dominant spores were collected under a dissecting microscope. Ten to 20 spores
145 were mounted on a glass slide to examine spore size and color under a light microscope.

146 Nodule bacteria were isolated from the rhizosphere soils or the root nodules of *L.*
147 *cryptobotrya* collected from the quarry and surrounding forest. The nodules were surface
148 sterilized with 70% ethanol for 30 s and sodium hypochlorite (0.5% available chlorine) for
149 5 min, washed with sterilized deionized water and crushed in 100 μ L sterilized deionized
150 water. Then the suspension was spread on the yeast extract-mannitol agar (YMA) medium
151 and cultured at 28°C. To isolate from the stored rhizosphere soils, 30 g of the soil was
152 layered between autoclaved river sand 1cm below surface in a 9 cm plastic pot (400 mL in
153 vol), and *L. bicolor* L. was grown in the greenhouse. We used *L. bicolor* instead of *L.*
154 *cryptobotrya* as a host plant, because *L. cryptobotrya* seed was not commercially available.
155 The plants were watered by a capillary sheet through the drain hole and grown without
156 fertilizer. The nodules formed on the roots were harvested after 2 months, and nodule
157 bacteria were isolated and maintained on the YMA medium. Single strains isolated from
158 each nodule or soil sample were randomly chosen and re-inoculated to *L. bicolor* grown on
159 the non-rhizospheric acid sulfate soil under the same conditions to select acid-tolerant
160 strains. The chemical properties of the soil are shown in Table 1, and the absence of AM

fungi and rhizobia in the soil was confirmed by cultivation of *L. bicolor* in a greenhouse prior to the experiments. Acid-tolerant nodule bacteria were isolated from the nodules formed on the roots after 2 months and maintained on the YMA medium.

Molecular identification of microsymbionts

Ten to 50 spores of the AM fungi were crushed in 10 μ L buffer of InstaGene matrix (BioRad Laboratories) under a microscope and mixed with 20 - 200 μ L of the matrix in a 1.5 mL tube. DNA was purified according to the manufacturer's instructions and stored at -30°C . A part (1.1 kbp) of the small subunit ribosomal RNA gene (18S rDNA) was amplified with the Expand High-Fidelity PCR System (Roche Diagnostics) with the NS1 and NS4 primer set (Table 2) using 1 μ L of the template DNA. The PTC-150 MiniCycler or PTC-225 DNA Engine Tetrad (MJ Research) was employed for the amplification, and the program was as follows: initial denaturation at 94°C for 2 min, followed by 30 cycles of denaturation at 94°C for 30 s, annealing at 52°C for 60 s, polymerization at 72°C for 80 s and final elongation at 72°C for 10 min.

DNA of the nodule bacteria was extracted from the colonies formed on the YMA medium by the InstaGene matrix according to the manufacturer's instructions and stored at -30°C . A part (1.3 kbp) of the small subunit ribosomal RNA gene (16S rDNA) was amplified under the same conditions described for AM fungi except for the primer set and

annealing temperature as follows. The 52f and 1320r primers were designed on the consensus regions of 16S rDNAs of the root nodule-forming bacteria across the four genera: *Mesorhizobium loti* (GenBank accession no. D14514), *Rhizobium leguminosarum* (D14513), *Shinorhizobium meliloti* (D14509), *Bradyrhizobium japonicum* (AF417550), *B. elkanii* (U35000) and *B. yuanmingense* (AF193818). The annealing temperature for this primer set was programmed at 60°C.

The PCR products were cloned into the pT7Blue T-vector (Novagen), and the nucleotide sequence of randomly chosen four clones (AM fungi) or a single clone (nodule bacteria) was determined by the dideoxy-sequencing method using the BigDye Terminator v3.0 or v3.1 Cycle Sequencing Kit with the ABI PRISM 3100 Genetic Analyzer (Applied Biosystems). The DNA sequences were subjected to phylogenetic analysis with published data by the CLUSTAL W program.

Effects of microsymbionts on the growth of pioneer plants

Non-rhizospheric acid sulfate soil collected from the quarry (Table 1) was used as a culture medium for the experiments. A preliminary screening of effective nodule bacteria was carried out using *L. bicolor* as a host, and *Bradyrhizobium* sp. QC1 and F3b (described later) were chosen for this experiment. The bacteria were cultured in 10 mL yeast extract-mannitol broth for 4 days at 28°C, centrifuged at 1,500×g for 15 min, washed twice with

deionized water then suspended in 50 mL deionized water. The mixed population of AM fungal inoculum was prepared by wet sieving from the soil trap culture of *L. cyrtobotrya* rhizosphere soil taken from the R2 plot (Fig. 1). *L. bicolor* or *M. sinensis* was sown on the acid sulfate soil in 400 mL plastic pots and inoculated with the AM fungi at 1,000 spores pot⁻¹. In the case of *L. bicolor*, 2 mL of the bacterial suspension was applied to the medium in addition to the AM fungi. The non-mycorrhizal plants received the same amount of the filtrate of spore suspension. The seedlings were thinned to 5 (*L. bicolor*) or 10 (*M. sinensis*) plants pot⁻¹ two weeks after sowing and grown in the greenhouse without fertilization. After 3 months, the shoots were dried at 80°C and weighed. The roots were cut into 1 cm segments, randomized completely in water, cleared and stained with Trypan blue to assess the levels of AM colonization.

Statistical analysis

All statistical analyses were performed with the StatView software (SAS Institute Inc., Cary, NC). For calculating the average pH values of rhizosphere soils, the data were transformed to real numbers. For the analysis of the pH preference of pioneer plants, the *t*-test was employed. In this analysis, the transformed pH values (real numbers) were to be retransformed to logarithmic values for normalization. Analysis of variance (ANOVA) and the Fisher's protected least significant difference test as a post hoc test ($P < 0.05$) were

applied for the analyses of the effect of the microsymbionts on the growth of pioneer plants.

Results

Occurrence of pioneer plant-microbial associations with respect to soil acidity

The three herbaceous plants, *M. sinensis*, *S. altissima* and *A. princeps*, and the leguminous shrub, *L. cyrtobotrya*, in the quarry formed arbuscular mycorrhizas without exception, although the extent of colonization was highly variable among the individuals. The percentage of AM colonization of *M. sinensis* was 3 - 46% ($n = 15$), and those of *L. cyrtobotrya* and *S. altissima* were 3 - 40% ($n = 13$) and 2 - 57% ($n = 7$), respectively. *A. princeps* was found only in 2002, and the levels of AM colonization were 10 - 27% ($n = 3$). Most of the *L. cyrtobotrya* found in the quarry formed root nodules. The precise assessment of the presence of nodules, however, was difficult because the whole root system could not be collected from the gravel-rich soil. Soil pH (H₂O) in the rhizosphere of the pioneer plants varied from 4.2 to 6.7, but most values were within a range of 4.2 - 5.5. Total C levels of the rhizosphere soils ranged from 2.1 to 20.4 g kg⁻¹ with an average value of 6.2 g kg⁻¹. Total N levels of the soils ranged from 0.38 to 1.23 g kg⁻¹ with an average value of 0.49 g kg⁻¹. Truog-P levels of the soils ranged from 20.5 to 71.3 mg kg⁻¹ with an average value of 41.6 mg kg⁻¹.

The levels of AM colonization and rhizosphere soil pH of the two dominant pioneer plants *M. sinensis* and *L. cyrtobotrya* were plotted, but the levels were not significantly decreased with an increase in soil acidity (Fig. 2a). The levels of AM colonization also did not correlate with the total C, N and Truog-P levels (data not shown). It is noteworthy that *L. cyrtobotrya* occurred preferentially in less acidic soils than for *M. sinensis* (Fig. 2b): the average soil pH of *L. cyrtobotrya* (4.96) was significantly higher than that of *M. sinensis* (4.64) ($P < 0.01$). The average values of the other soil parameters did not differ significantly between the two pioneer species (data not shown).

Isolation and identification of microsymbionts

AM fungi were successfully cultured from the rhizosphere soils of R2 (*L. cyrtobotrya*), R5 (*M. sinensis*), R8 (*S. altissima*), R9 (*L. cyrtobotrya*), QR2 (*M. sinensis*) and QR3 (*M. sinensis*) plots (Fig. 1). HR1 of which spore diameter and wall thickness were 243.3 ± 46.5 μm and 16.0 ± 4.5 μm , respectively, dominated in the R2, R8, R9 and QR2 soils (Fig. 3a). The 1.1 kbp sequences of 18S rDNA of HR1 fell into the cluster of *Gl. manihotis*, *Gl. clarum* and *Gl. intraradices* to which HR1 sequences showed 97.8-99.1% similarity (Fig. 4), and thus HR1 was classified in the genus *Glomus*. HB1 was a dominant in the R5 soil, and its spore diameter and wall thickness were 135.5 ± 21.0 μm and 8.9 ± 3.0 μm , respectively (Fig. 3b). The spores occasionally retained a hyphal terminus that was a typical structure for

the genus *Entrophospora*. The 18S rDNA sequences of HB1 were almost identical to that of *Entrophospora colombiana*. Based on the morphological and molecular data, HB1 was classified in the genus *Entrophospora*. WL1 was found in the QR2 and QR3 soils, and the spore diameter and wall thickness were $154.5 \pm 30.4 \mu\text{m}$ and $8.9 \pm 1.0 \mu\text{m}$, respectively (Fig. 3c). The 18S rDNA sequences of WL1 showed highest similarity to that of *Archaeospora leptoticha*, and thus WL1 was classified in the genus *Archaeospora*. HRF1 was the only fungus that was cultured successfully from the surrounding forest (FR13 plot), and the spore morphology was similar to that of *Glomus* sp. HR1 (Fig. 3d). The 18S rDNA sequences of HRF1 and *Glomus* sp. HR1 showed highest similarity to each other (99-100% similarity), indicating that HRF1 was also classified in the genus *Glomus*.

All acid-tolerant nodule bacteria isolated from the quarry (QC1, QC2, QC3 and QC7) and the surrounding forest (F3b and F4b) were classified into the genus *Bradyrhizobium* (Fig. 5). The 1.3 kbp sequence of 16S rDNA of QC1 was identical to that of *Bradyrhizobium* sp. MAFF210318. The sequence of QC3 was identical to that of *B. japonicum* USDA62. All of the QC2, QC7, F3b and F4b sequences were identical and showed highest similarity to that of *B. elkanii* USDA76.

Effectiveness of indigenous microsymbionts in acid sulfate soil

The single inoculation of the AM fungi (*Glomus* sp. HR1 was a dominant) or the nodule

bacterium, *Bradyrhizobium* sp. QC1 or F3b, did not improve the growth of *L. bicolor* significantly (Fig. 6a). Dual inoculation of the fungi and bacterium, however, promoted the growth of *L. bicolor* significantly compared with that of the non-mycorrhizal/non-bacterial control. The effectiveness of nodule bacterium on the growth did not differ between the two strains. The percentages of AM colonization of *L. bicolor* in the presence or absence of the nodule bacterium were $19.3 \pm 3.5\%$ (no bacterium), $11.6 \pm 7.2\%$ (QC1) and $24.5 \pm 3.5\%$ (F3b), respectively. No colonization was observed in the uninoculated plants. Non-mycorrhizal *M. sinensis* showed chlorosis and quite poor growth, whereas the growth of the mycorrhizal plants was more than 10-fold higher than that of the non-mycorrhizal plants (Fig. 6b). The levels of AM colonization were $39.0 \pm 5.6\%$ and 0% in the mycorrhizal and non-mycorrhizal plants, respectively.

Discussion

Significance of pioneer plant-microbial associations in acid sulfate soil

The present study clearly demonstrated that the symbiotic associations play significant roles in the growth of pioneer plants in acid sulfate soil. Without exception, all of the pioneer herbs and shrubs formed AM associations, and the grass and legume pioneers showed significant growth responses to the indigenous microsymbionts. Pioneer grass species are

generally regarded as facultative mycotrophic (Allen 1991; Oba et al. 2004). Wilson and Hartnett (1998) found, however, a strong relationship between the phenology of grass species and mycorrhizal responsiveness in a tallgrass prairie ecosystem: perennial warm-season C₄ grasses were highly mycorrhiza-responsive but perennial cool-season C₃ grasses were not. Given the fact that *M. sinensis* is a perennial C₄ grass, our observations fit well with their results. *Lespedeza* spp. are known to be responsive to AM fungi (Wilson 1988; Wilson and Hartnett 1998), and Johnson et al. (2005) pointed out that the relatively coarse root system of *Lespedeza* spp. might be a key characteristic to receive benefit from AM fungi. On the other hand, it has been suggested that the establishment of *Lespedeza* spp. may be limited by the density of rhizobia in soil, based on the observations in N-poor grasslands (Larson and Siemann 1998). Considering that *M. sinensis* (Kayama 2001) and *Lespedeza* spp. (Cline and Silvernail 1997) are intrinsically acid-tolerant, it is likely that nutrient deficiency is a major limiting factor for these pioneers rather than soil acidity during early primary succession of acid sulfate soil. In addition, our observations also suggest that the establishment of symbiotic associations with the microorganisms is crucial for the initial growth and subsequent survival of the pioneer plants in this ecosystem. Further investigations, e.g. field inoculation experiments or pot experiments with a longer period are required to examine this hypothesis.

AM fungi in acid sulfate soil

321

322 The present study provides the first information about AM fungi in extremely acidic soil. It
323 is considered that the fungi associated with the pioneer plants in the site are highly acid-
324 tolerant due to the following reasons: i) no significant decline in the levels of colonization
325 with an increase in soil acidity was observed, and ii) the fungi isolated from the field
326 promoted the growth of the pioneer plants significantly even at pH 3.4. *Gl. clarum*, *E.*
327 *colombiana* and *Gl. leptotichum* (a synanamorph of *Ar. leptoticha*) as close relatives of
328 *Glomus* sp. HR1, *Entrophospora* sp. HB1 and *Archaeospora* sp. WL1, respectively, have
329 been isolated from acidic soils with pH less than 4
330 (<http://invam.caf.wvu.edu/cultures/accessions.htm>). Recently, the community compositions
331 of AM fungi that associate with *M. sinensis* grown in acid sulfate soil have been
332 investigated (An et al. 2008). Interestingly, *Glomus* sp. HR1 (corresponding to GLO3
333 phylotype in An et al. (2008)) was widely distributed in the rhizosphere from subarctic to
334 subtropical Japan and occurred in a wide range of soil pH. Combining their observations
335 with our finding that *Glomus* sp. HR1 was isolated from the rhizosphere of several pioneer
336 species, it is suggested that this fungus is the ‘pioneer AM fungus’ that is widely distributed
337 and plays major roles in pioneer vegetation of acid sulfate soil. In contrast to *Glomus* sp.
338 HR1, *Entrophospora* sp. HB1 was cultured from only one soil sample in the present study.
339 Oba et al. (2004) investigated the AM fungal communities in the pioneer vegetation of the
340 lahar (volcanic mud flow) area of Mt. Pinatubo, Philippines and found a preferential

association of *E. colombiana* with a leguminous plant *Calpogonium mucunoides* that was regarded as a member of secondary vegetation after the establishment of grass pioneers. According to An et al. (2008), *Entrophospora* spp. could not be detected from *M. sinensis* roots grown in acid sulfate soil even from Hazu site (the same field as the present study). These observations suggest that *Entrophospora* sp. HB1 may preferentially occur in a late phase of early succession. In the case of *Archaeospora* sp. WL1, information about ecosystems in which *Ar. leptoticha* occurs is much less available. Further investigations are required to evaluate the significance of this species in pioneer vegetation.

Rhizobia in acid sulfate soil

All of the nodule bacteria isolated in the present study were classified into the genus *Bradyrhizobium*. It has been reported that *Lespedeza* spp. associates with a wide range of nodule bacteria, e.g. *Sinorhizobium* spp., *Rhizobium legumae* and *B. elkanii* formed nodules with *L. bicolor* (Yao et al. 2002). In acidic soil, however, it seems likely that *Bradyrhizobium* spp. are the major symbiotic partners (Cline and Senwo 1994). The acid-tolerant bradyrhizobia associated with *Lespedeza* spp. were able to grow on acidic agar media, but the lower limit of the *in vitro* growth of these strains was pH 4.5 (Cline and Senwo 1994), reflecting the acid-tolerance of the host plant that could not grow below pH 4.5 (Cline and Senwo 1993). Our observation that *L. cyrtobotrya* occurred in soils with pH

more than 4.5 is in good agreement with that of Cline and Senwo (1993). Although the acid-tolerance of the bacterial symbionts is likely to be a significant factor for the effectiveness of N-fixation under acidic conditions, Richardson and Simpson (1989) found negative correlations between the acid tolerance of *R. triforii* isolates *in vitro* and their effectiveness in the field, indicating that the more acid-tolerant strains were less capable of infecting the host plants. In the present study, bradyrhizobia that promoted the growth of the host plants under extremely acidic conditions were successfully isolated by using acid sulfate soil as a selection medium, suggesting that choosing an appropriate medium/culture system for the screening is particularly important to obtain effective rhizobia. The P-status of the culture medium, in addition, should be taken into account for evaluating the effectiveness of selected bacteria under acidic conditions in which P-availability is generally low, because extremely low-P conditions suppress N-fixing activity. In this case, P-fertilization or the inoculation of suitable (acid-tolerant) AM fungi is to be considered.

Origin of microsymbionts in acid sulfate soil

It is likely that the symbiotic microorganisms observed in the quarry originated from the surrounding forest, because the phylogenetic analyses revealed that the AM fungi and nodule bacteria isolated from the quarry were closely related to those isolated from the surrounding forest. The idea that the forest soil is a potential habitat for acid-tolerant

microsymbionts may be further supported by the fact that soil pH in the surrounding forest was 4.5 - 5.0 (data not shown), as is typical of temperate forest. It has been known that AM fungal propagules are dispersed into disturbed areas by wind, animal activities and erosion (Allen 1987; Warner et al. 1987). In the case of our field, soil erosion from the surrounding forest by rainfall might be a major vector of the fungal propagules and bradyrhizobia, because the ground level of the forest is 20-100 m higher than that of the quarry.

Conclusion

Isolation of acid-tolerant symbiotic microorganisms from pioneer plants will be a potential strategy for the application of the microorganisms to the revegetation program of acid sulfate soil. Further investigations are required, however, to clarify whether the inoculation of multiple strains is more effective than that of a single strain. For example, Jansa et al. (2008) demonstrated that the inoculation of multiple AM fungal species resulted in the greater P-uptake in medic and leek than that of the single species. Given the large heterogeneities in soil microenvironments and water regime, it is expected that the colonization with the multiple strains that differ in environmental preference may improve the adaptability of the host plants via maintaining the overall efficiency of acquisition of mineral nutrients under more diverse environments.

Acknowledgements

We are grateful to Drs. I. Nioh, K. Saito, M. Satio, M. Abe and Y. Hashimoto for invaluable suggestions, to M. Maesaka, S. Mizuno and Y. Tahara in Nagoya University for technical assistance and to Aichi prefecture for allowing us to collect the samples from the field. This study was supported by Tokai Gakujutsu Shoreikai and the Japan Society for the Promotion of Science (TE).

References

- Allen MF 1987 Re-establishment of mycorrhizas on Mount St Helens Washington USA migration vectors. *Trans Brit Mycol Soc* 88:413-417
- Allen MF 1991 The ecology of mycorrhizae. Cambridge University Press, Cambridge, New York, USA. pp. 127-140
- An G-H, Miyakawa S, Kawahara A, Osaki M and Ezawa T 2008 Community structures of arbuscular mycorrhizal fungi associated with pioneer grass species *Miscanthus sinensis* in acid sulfate soils: habitat segregation along pH gradients. *Soil Sci Plant Nutri* (in press)
- Appleyard S, Wong S, Willis-Jones B, Angeloni J and Watkins R 2004 Groundwater acidification caused by urban development in Perth, Western Australia: source, distribution, and implications for management. *Aust J Soil Res* 42:579-585
- Barea JM, Azcon R and Azconaguilar C 1992 Vesicular-arbuscular mycorrhizal fungi in nitrogen-fixing systems. *Meth Microbiol* 24:391-416

424 Bohlool BB, Ladha JK, Garrity DP and George T 1992 Biological nitrogen-fixation for
 425 sustainable agriculture - a perspective. *Plant Soil* 141:1-11

426 Brundrett MC, Abbott LK and Jasper DA 1999 Glomalean mycorrhizal fungi from tropical
 427 Australia I. Comparison of the effectiveness and specificity of different isolation
 428 procedures. *Mycorrhiza* 8:305-314

429 Cline GR and Senwo ZN 1993 Inhibitory effects of acidic minesoil on the *Sericea*
 430 *lespedeza/Bradyrhizobium* symbiotic relationship. *J Plant Nutri* 16:1867-1880

431 Cline GR and Senwo ZN 1994 Tolerance of *lespedeza Bradyrhizobium* to acidity, aluminum,
 432 and manganese in culture media containing glutamate or ammonium. *Soil Biol*
 433 *Biochem* 26:1067-1072

434 Cline GR and Silvernail AF 1997 Effects of soil acidity on the growth of *sericea lespedeza*.
 435 *J Plant Nutri* 20:1657-1666

436 Do TTR, Tinh TK, Nguyen TNM and Linh TB 2004 Applying mixed manure and inorganic
 437 phosphorus fertiliser to improve rice yield on acid sulfate soil (Hydraquentic
 438 Sulfaquept). *Aust J Soil Res* 42:693-698

439 Giovannetti M and Mosse B 1980 An evaluation of techniques for measuring vesicular-
 440 arbuscular mycorrhizal infection in roots. *New Phytol* 84:489-500

441 Herrera MA, Salamanca CP and Barea JM 1993 Inoculation of woody legumes with
 442 selected arbuscular mycorrhizal fungi and rhizobia to recover desertified
 443 Mediterranean ecosystems. *Appl Environ Microbiol* 59:129-133

444 Jansa J, Smith FA and Smith SE 2008 Are there benefits of simultaneous root colonization
 445 by different arbuscular mycorrhizal fungi? *New Phytol* 177:779-789

446 Johnson NC, Wolf J, Reyes MA, Panter A, Koch GW and Redman A 2005 Species of plants
 447 and associated arbuscular mycorrhizal fungi mediate mycorrhizal responses to CO₂
 448 enrichment. *Global Change Biol* 11:1156-1166

449 Kayama M 2001 Comparison of the aluminum tolerance of *Miscanthus sinensis* Anderss.
 450 and *Miscanthus sacchariflorus* Bentham in hydroculture. Int J Plant Sci 162:1025-
 451 1031

452 Larson JL and Siemann E 1998 Legumes may be symbiont-limited during old-field
 453 succession. American Midland Naturalist 140:90-95

454 Nara K and Hogetsu T 2004 Ectomycorrhizal fungi on established shrubs facilitate
 455 subsequent seedling establishment of successional plant species. Ecology 85:1700-
 456 1707

457 Oba H, Shinozaki N, Oyaizu H, Tawaraya K, Wagatsuma T, Barraquio WL and Saito M
 458 2004 Arbuscular mycorrhizal fungal communities associated with some pioneer
 459 plants in the Lahar area of Mt. Pinatubo, Philippines. Soil Sci Plant Nutri 50:1195-
 460 1203

461 Porter WM, Robson AD and Abbott LK 1987 Factors controlling the distribution of
 462 vesicular-arbuscular mycorrhizal fungi in relation to soil pH. J Appl Ecol 24:663-
 463 672

464 Powell B and Martens M 2005 A review of acid sulfate soil impacts, actions and policies
 465 that impact on water quality in Great Barrier Reef catchments, including a case
 466 study on remediation at East Trinity. Mar Poll Bull 51:149-164

467 Requena N, Jimenez I, Toro M and Barea JM 1997 Interactions between plant-growth-
 468 promoting rhizobacteria (PGPR), arbuscular mycorrhizal fungi and *Rhizobium* spp.
 469 in the rhizosphere of *Anthyllis cytisoides*, a model legume for revegetation in
 470 mediterranean semi-arid ecosystems. New Phytol 136:667-677

471 Requena N, Perez-Solis E, Azcon-Aguilar C, Jeffries P and Barea J-M 2001 Management of
 472 indigenous plant-microbe symbioses aids restoration of desertified ecosystems. Appl
 473 Environ Microbiol 67:495-498

474 Richardson AE and Simpson RJ 1989 Acid-Tolerance and symbiotic effectiveness of
 475 rhizobium-trifolii associated with a *Trifolium-Subterraneum* L-based pasture
 476 growing in an acid soil. Soil Biol Biochem 21:87-95
 477 Smith SE and Read DJ 1997 Mycorrhizal symbiosis. Academic Press, San Diego, California.
 478 pp. 605
 479 Truog E 1930 The determination of the readily available phosphorus of soils. J Am Soc
 480 Agron 22:874-882
 481 van Aarle I M, Soderstrom B and Olsson PA 2003 Growth and interactions of arbuscular
 482 mycorrhizal fungi in soils from limestone and acid rock habitats. Soil Biol Biochem
 483 35:1557-1564
 484 Warner NJ, Allen MF and Macmahon JA 1987 Dispersal agents of vesicular-arbuscular
 485 mycorrhizal fungi in a disturbed arid ecosystem. Mycologia 79:721-730
 486 Wilson D 1988 Differential plant response to inoculation with two VA mycorrhizal fungi
 487 isolated from a low-pH soil. Plant Soil 110:69-75
 488 White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal
 489 ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ,
 490 White TJ (eds) PCR protocols: a guide to methods and applications. Academic Press,
 491 San Diego, pp 315-322
 492 Wilson GWT and Hartnett DC 1998 Interspecific variation in plant responses to
 493 mycorrhizal colonization in tallgrass prairie. Am J Bot 85:1732-1738
 494 Yao ZY, Kan FL, Wang ET, Wei GH and Chen WX 2002 Characterization of rhizobia that
 495 nodulate legume species of the genus *Lespedeza* and description of *Bradyrhizobium*
 496 *yuanmingense* sp. nov. Int J Syst Evol Microbiol 52:2219-2230




Fig. 1 Sampling location of *Miscanthus sinensis*, *Lespedeza cyrtobotrya*, *Solidago altissima* and *Artemisia princeps* in the quarry in Hazu-cho, Aichi. The sample ID from which arbuscular mycorrhizal fungi were successfully proliferated by the trap culture is indicated. Asterisk shows the sampling point of the soil for the pot experiments.




Fig. 2 Rhizosphere soil pH and percentage mycorrhizal colonization of *Miscanthus sinensis* ($n = 15$) and *Lespedeza cyrtobotrya* ($n = 13$) roots grown in the quarry in Hazu-cho, Aichi. **a**, Scatter plot of the percentage colonization and soil pH. **b**, Frequency distributions of the soil pH.





Fig. 3 Spore morphology of arbuscular mycorrhizal fungi cultured from the rhizosphere soils of pioneer plants. **a**, HR1 isolated from the R2, R8, R9, R10 and QR2 soils from the quarry. **b**, HB1 isolated from the R2 and R5 soils from the quarry. ht, hyphal terminus. **c**, WL1 isolated from the QR3 soil from the quarry. **d**, HRF1 isolated from the FR13 soil from the surrounding forest. Bars are 100 μm .



0.01

Fig. 4 Phylogenetic analysis of arbuscular mycorrhizal fungi isolated from the rhizosphere of pioneer plants grown in acid sulfate soil (HR1, HB1 and WL1) or in the surrounding forest (HRF1) based on the 1.1 kbp sequences of small subunit ribosomal RNA gene. Bootstrap values more than 70% (1,000 replications) are indicated. The numbers in the parentheses are GenBank accession numbers.





Fig. 5 Phylogenetic analysis of nodule bacteria isolated from the rhizosphere soil or the nodules of *Lespedeza cyrtobotrya* grown in the quarry (QC1, QC2, QC3 and QC7) or in the surrounding forest (F3b and F4b) based on the 1.3 kbp sequences of small subunit ribosomal RNA gene. Bootstrap values more than 70% (1,000 replications) are indicated. The numbers in the parentheses are GenBank accession numbers.

a) *L. bicolor*



b) *M. sinensis*




Fig. 6 Effectiveness of symbiotic microorganisms on the growth of pioneer plants in acid sulfate soil at pH 3.4. **a**, The effect of dual inoculation of bradyrhizobium (QC1 or F3b) and arbuscular mycorrhizal fungi (AMF) on the growth of *Lespedeza bicolor*. **b**, The effect of AMF on the growth of *Miscanthus sinensis*. *Bradyrhizobium* sp. QC1 and F3b were isolated from the quarry and the surrounding forest, respectively. NM, non-mycorrhizal control; AMF, inoculated with the mixed population of AMF proliferated from the R2 plot soil (*Glomus* sp. HR1 was a dominant). Different letters show significant differences (Fisher's LSD test, $P < 0.05$).

Table 1 Chemical properties of the potting medium collected from the quarry.

pH (H ₂ O)	3.4	
pH (H ₂ O ₂)	2.3	
EC	88.0	mS m ⁻¹
Total C	1.9	g kg ⁻¹
Total N	0.3	g kg ⁻¹
C/N	6.3	
Truog-P	7.6	mg P kg ⁻¹
CEC	83.2	mmol _c kg ⁻¹
Na	1.7	mmol _c kg ⁻¹
Mg	3.4	mmol _c kg ⁻¹
K	1.4	mmol _c kg ⁻¹
Ca	27.1	mmol _c kg ⁻¹
Fe	0.2	mmol _c kg ⁻¹
Al	11.4	mmol _c kg ⁻¹
Mn	Trace	

Table 2 PCR primers used in this study.

Name	Sequence (5' > 3')	Notes
NS1	GTAGTCATATGCTTGTCTC	18S rDNA, universal (White et al., 1990)
NS4	CTTCCGTCAATTCCTTTAAG	18S rDNA, universal (White et al., 1990)
52f	ACGCTGGCGGCAGGCTTAAC	16S rDNA, rhizobia universal (this study)
1320r	CAACTCCCATGGTGTGACG	16S rDNA, rhizobia universal (this study)