HOKKAIDO UNIVERSITY

Title	Stroboscopic Picosecond Pulse Radiolysis Studies of the Short Lived Cation Produced in Butylhalides
Author(s)	Sumiyoshi, Takashi; Yamada, Tatsuya; Ohtaka, Akira et al.
Citation	北海道大學工學部研究報告, 139, 149-152
Issue Date	1988-02-20
Doc URL	https://hdl.handle.net/2115/42076
Type	departmental bulletin paper
File Information	139_149-152.pdf

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

北海道大学工学部研究報告
第 139 号（昭和 63 年）

Bulletin of the Faculty of Engineering,
Hokkaido University. No. 139 (1988)

Stroboscopic Picosecond Pulse Radiolysis Studies of the Short Lived Cation Produced in Butylhalides

Takashi SUMIYOSHI, Tatsuya YAMADA, Akira OHTAKA,
Kazunori TSUGARU, and Meiseki KATAYAMA
(Received Spetember 30, 1987)

Abstract

Transient species produced in neat butylchloride, butylbromide and butyliodide were observed by means of picosecond pulse radiolysis using 11 ns pulses of 45 MeV electrons.

The visible band which was produced during electron pulse and then decayed with a half lifetime of 2 ns was assigned to a vibrationally excited cation.

Introduction

Butylchloride has been used as one of the suitable solvents to produce solute cations by charge transfer for its high ionization potential (10.67 eV).[1,2] Pulse radiolysis studies at low temperature in the μs time range[3] and at room temperature in the ns time range[2] have been carried out. However, the nature of the solvent cations has not been well understood so far. Because of the short lifetime of cations in the liquid phase, the picosecond pulse radiolysis technique should provide important information on the behavior of cations in the very early stage after the irradiation. We studied the kinetic behavior of cations in the ps and ns time regions and found fast decaying cations which have a much shorter lifetime than that observed in the ns pulse radiolysis.

Experimental

Butylchloride, butylbromide and butyliodide used were guaranteed reagents of Tokyo Kasei and were purified by distillation. The stroboscopic picosecond pulse radiolysis system[4] and the conventional ns one[5] were described earlier. Samples were circulated through the 2-cm optical cell by a pump. The absorbed dose was measured by the KCNS dosimeter[6] for ns macro pulse and by the aqueous electron dosimeter for picosecond fine structure electron pulse using $G(e^-_{aq})$ at 30 ps as 4. 8.[7]

Results

Butylchloride

Figure 1 shows typical picosecond kinetic traces observed in neat butylchloride at

Department of Atomic Science and Nuclear Engineering, Hokkaido University, Sapporo 060

indicated wavelengths. Since the decay of the transient species is not completed between fine structure electron pulses (350 ps), the absorption signal produced by each fine structure pulse indicated as h is superimposed on the buildup absorption indicated as H. Dashed lines are computer simulated kinetic traces. The simulation was carried out to obtain the best fit for h, H and the slope of traces. All kinetic traces could be reproduced by a single exponential decay with a half lifetime of 1.7±0.3ns, and estimated $G\varepsilon$ values were 11,000, 14,000, and 9,000 at 450, 520, and 630 nm, respectively.

Figure 2 shows the buildup spectrum (H) and step signal (h) produced by fine structure electron pulses with a half width of 30 ps. Both spectra with peaks at 500 nm are quite similar to those reported already[1-3] and could be assigned to parent cations. On the contrary, the present ns pulse radiolysis experiment shows that the transient absorption with a maximum at 500 nm decays with a half lifetime of 50 ns which is good agreement with that



Fig. 1 Picosecond kinetic traces observed 450, 520, and 630 nm in neat butylchloride at room temperature. Dashed lines are simulated traces



Fig. 2 The buildup spectrum (○) and the spectrum produced by fine structure (■) in butylchloride



Fig. 3 Buidup spectra of butylbromide. Dashed lines are estimated two components



Fig. 4 Kinetic traces observed at 390, 450, and 600 nm in butylbromide at room temperature. Dashed lines are simulated traces with parameters given in Table 2.

Table 1 $G\varepsilon$ and Half Lifetimes Observed by Nanosecond Pulse Radiolysis of Butylbromide

λ/nm	$G\varepsilon$	$T_{1/2}$
380	10,000	4μs
450	4,500	2.3μs
590	2,500	100ns

Table 2 $G\varepsilon$ Values Observed by Picosecond Pulse Radiolysis of Butylbromide

λ/nm	$T_{1/2}$=2ns	$T_{1/2}$=4 μs
390	5,000	10,000
450	9,000	4,000
600	17,000	0



Fig. 5 Buildup spectra of butyliodide



Fig. 6 Kinetic traces observed at 400, 600 and 700 nm in butyliodide at room temperature. Dashed lines are simulated traces with parameters given in the text

observed by Mehnert et al.[2] Further, the $G\varepsilon$ value at 500 nm in ns pulse radiolysis was about 2,000$M^{-1}cm^{-1}$.

Butylbromide

Figure 3 shows buidup spectra of butylbromide indicating two absorption peaks at 390 and 600 nm which are in good agreement with the spectrum observed by the ns pulse radiolysis. $G\varepsilon$ values and half lifetimes obtained in the ns pulse radiolysis are given in Table 1. Figure 4 shows ps kinetic traces of butylbromide. The kinetic trace at 600 nm was reproduced by a single exponential decay with a half lifetime of 2 ± 0.2 ns. On the other hand, traces at 390 and 450 nm could be simulated by assuming two components which are given in Table 2. Dashed lines in Fig. 3 were estimated buildup spectra of two components. The band at 380 nm with the half lifetime of 4 μs can be assigned to Br_2^-[8]. And the visible band at 600 nm with a half lifetime of 2 ns can be assigned to parent cations.[3,8]

Butyliodide

Figure 5 shows the buildup spectrum of irradiated butyliodide indicating three absorption maxima at 400, 500 and>800 nm. A similar spectrum is observed by the ns pulse radiolysis.

Figure 6 shows picosecond kinetic traces observed in butyliodide at indicated wave-

lengths. The kinetic traces at 400 and 700 nm were reproduced by a single exponential decay with a half lifetime of 5 ns and 2.5 ns, and estimated $G\varepsilon$ values were 15,000 and 15,500 M^{-1} cm^{-1}, respectively.

Discussion

The present experimental results of three butylhalides can be better explained by the existence of vibrationally excited butylhalide cations as suggested in the low temperature pulse radiolysis experiment.[3] The observation of rapid formation of visible bands (<30 ps), excludes the contribution of an ion-pair intermediate such as ($BuCl^+ \ldots {}^-Cl$) which was suggested by an analogy of radiolysis of other alkyl halides.[2] The low $G\varepsilon$ value and the relatively long half lifetime of cations observed in ns pulse radiolysis suggested that a part of vibrationally excited cations might be deactivated to the ground state. The existence of vibrationally excited cations has been also suggested in a series of picosecond pulse radiolysis experiments of alkyl halides and alkanes.[9-11]

References

1) T. Shida and W. H. Hamill, J. Chem. Phys., **44**, 4372 (1966).
2) R. Mehnert, O. Brede, and W. Naumann, Ber. Bunsenges. Phys. Chem., **86**, 525 (1982).
3) S. Arai, A. Kira, and M. Imamura, J. Phys. Chem., **80**, 1968 (1976).
4) T. Sumiyoshi, S. Sawamura, Y. Koshikawa, and M. Katayama, Bull. Chem. Soc. Jpn., **55**, 2341 (1982).
5) Y. Kondo, M. Aikawa, T. Sumiyoshi, M. Katayama, and J Kroh, J. Phys. Chem., **84**, 2544 (1980).
6) E. M. Fielden and N. W. Holm, "Manual on Radiation Dosimetry," ed. by N. M. Holm and R. J. Berry, Marcel Dekker, New York (1970), p. 261.
7) T. Sumiyoshi and M. Katayama, Chem. Lett., **1982**, 1887.
T. Sumiyoshi, K. Tsugaru, T. Yamada, and M. Katayama, Bull. Chem. Soc. Jpn., **58**, 3073 (1985).
8) R. F. C. Claridge and J. E. Willard, J. Am. Chem. Soc., 88, 2404 (1966).
9) T. Sumiyoshi, S. Sawamura, Y. Koshikawa, and M. Katayama, Proc. 1982 Tihany Symp. Radiat. Chem., **1983**, 453.
10) T. Sumiyoshi, K. Tsugaru, and M. Katayama, Chem. Lett., **1982**, 1431.
11) T. Sumiyoshi, K. Tsugaru, and M. Katayama, Chem. Lett., **1982**, 1883.