



HOKKAIDO UNIVERSITY

Title	HIV-1 Vpr: A Novel Role in Regulating RNA Splicing
Author(s)	Zhang, Xianfeng; Aida, Yoko
Citation	Current HIV Research, 7(2), 163-168
Issue Date	2009-03
Doc URL	https://hdl.handle.net/2115/42849
Type	journal article
File Information	7-2_p163-168.pdf



HIV-1 Vpr: a novel role in regulating RNA splicing

Xianfeng Zhang^{a, b} and Yoko Aida^{b*}

^a Institute for Genetic Medicine, Hokkaido University, Kita-ku, Sapporo 060-0815,
Japan

^b *Viral Infectious Diseases Unit, RIKEN, Wako, Saitama 351-0198, Japan*

Running Title: RNA splicing regulation by HIV-1 Vpr

Keywords: HIV-1 Vpr, splicing inhibition, cellular pre-mRNA, HIV-1 genome,
spliceosome

***Corresponding author.** Tel.: +81-48-462-4408; fax: +81-48-462-4399; e-mail
address: aida@riken.jp; postal address: Viral Infectious Diseases Unit, RIKEN, 2-1
Hirosawa, Wako, Saitama 351-0198, Japan.

ABSTRACT

Pre-mRNA splicing is a critical step in gene expression for metazoans. Several viral proteins regulate the splicing of pre-mRNAs through its complex interactions between the virus and the host cell RNA splicing machinery. Here, we focus on a novel function of HIV-1 Vpr, that selectively inhibit cellular and viral pre-mRNA splicing, via interactions with components of functional spliceosomal complexes. This review discusses our current knowledge of how RNA splicing regulation is accomplished by Vpr-host factors interactions.

INTRODUCTION

The relatively small genome of HIV-1 encodes structural and enzymatic proteins, such as Gag, Pol and Env, several regulatory proteins, such as Tat, Rev and accessory proteins such as Vif, Vpr, Vpu and Nef, through alternative splicing and alternate codon usage. In recent years, considerable attention has been paid to these accessory genes and great deals of evidences had been found to discover the great contributions of these proteins to HIV-1 replication. Vif, for example, was found recently to overcome the sabotage effect of APOBEC3G against HIV reverse transcription (Sheehy *et al*, 2003). Vpu enhances HIV particle assembly or release in a variety of human cells by antagonize a host factor, tetherin's antiviral activity (Neil *et al* 2008). Nef manipulate the expression of host cell surface molecules to benefit the viral propagation.

Vpr, a virion-associated viral protein 96 amino-acids in length, has multiple biological functions including its nuclear localization activity (Ijima *et al*, 2004; Jenkins *et al*, 1998; Kamata and Aida, 2000; Kamata *et al*, 2005; Popov *et al*, 1998), arrest at the G2/M phase of the cell cycle (Sherman *et al*, 2000; Poon *et al*, 1998), increasing the activity of the HIV-1 long terminal repeat (LTR) (Goh *et al*, 1998), positive and negative regulation of apoptosis (Ayyavoo *et al*, 1997; Azuma *et al*, 2006; Jacotot *et al*, 2000; Nishizawa *et al*, 1999, 2000a, 2000b; Stewart *et al*, 2000) and induction of cellular nuclear envelope herniation (de Noronha *et al*, 2001). Recently, we discovered a novel role for Vpr in the selective inhibition of cellular pre-mRNA splicing both *in vivo* and *in vitro* (Kuramitsu *et al*, 2005). The role, if any, of Vpr in splicing of the HIV-1 genome is unknown. In this review, we will describe our

current understanding of the novel role of Vpr in regulating RNA splicing via Vpr-host factors interactions.

WHAT WE KNOW ABOUT SPLICING REACTION

Protein-coding sequences (exons) in primary transcripts are interrupted by intervening sequences (introns) that are eliminated from the pre-mRNA by splicing in the nucleus (Krämer 1996). The splicing reaction is carried out by the spliceosome, which comprises the small nuclear ribonucleoproteins (snRNPs) U1, U2, U4/U6 and U5, and over 150 proteins of non-snRNP proteins termed splicing factors. During splicing, successive recognition of splice sites occurs through the stepwise assembly of spliceosomal complexes E, A, B, and C on pre-mRNA (Hastings and Krainer, 2001). In the commitment complex E, U1 snRNP is bound to the 5' splice site and the U2 snRNP auxiliary factor (U2AF) and splicing factor-1 (SF1) are attached to the polypyrimidine tract upstream of the 3' splice site and the branchpoint sequence (BPS), respectively. As U2 snRNP binds to the branch site and complex E is converted to presplicing complex A, a short helix forms between a single-stranded region in U2 snRNA and the intron branch site. The next step is binding of U4/U5/U6 tri-snRNP at the 5' splice site to form the B complex. Just prior to splicing, conformational changes within complex B destabilize the association between U4 and U2/U5/U6 and transform complex B into complex C. The U2 snRNP, one of the key factors in the spliceosome assembly pathway, is classified to two complexes, SF3a and SF3b, SF3a consists of three subunits, SAP60, 62 and 114, and SF3b consists of at least four subunits, SAP49, 130, 145 and 155 an essential component of the SF3b subunit, can bind to pre-mRNA (Staknis *et al.*, 1994; Hastings and Krainer, 2001; Will *et al.*, 2001) and is implicated in the tethering of U2 snRNA to the branch point

site (Champion-Arnaud and Reed, 1994). The large catalytic spliceosome performs two transesterification reactions on the splice sites and subsequently result in the ligation of the exons and the excision of the intron which formed as lariat RNA with its 5' phosphate joined to the 2' hydroxyl at the branch point.

COMPLICATE SPLICING PATTERN OF HIV-1

A complex pattern of splicing generates more than 30 mRNA species from the single primary mRNA transcript of the HIV genome (Purcell *et al*, 1993, Schwartz *et al*, 1990; Tang *et al*, 1999). This complex pattern of splicing is required to generate mRNAs for viral structural, regulatory and accessory proteins, and undoubtedly plays a critical role in the regulation of HIV-1 gene expression. Viral mRNA species are classified into three main groups. The 2kb-class of mRNAs, which are fully spliced and constitutively exported to the cytosol early in the HIV-1 replication cycle, encode the regulatory proteins Tat, Rev and Nef. The singly spliced 4kb-class of mRNAs are bicistronic, and encode Env, Vpr, Vif and Vpu. The unspliced 9kb-class of mRNA encodes Gag and Gag-Pol polyproteins, and also serves as the source of HIV genomic RNA for packaging into virions. Nuclear export and translation of the 4kb- and 9kb-class of mRNAs is dependent on Rev, which binds to the Rev responsive element (RRE) present in these mRNAs. Correct splicing to produce HIV-1 viral mRNAs is dependent on alternative utilization of four 5'-donor sites, termed D1 to D4, and eight 3'-accept sites, A1, A2, A3, A4a, A4b, A4c, A5 and A7 (Purcell *et al*, 1993). An additional 3'-splice site, A6, has been identified in only one HIV-1 strain to date (Benko *et al*, 1990). Donor sites D1, D2 and D3 can couple to any of the A1 to A5 acceptor sites, whereas donor site D4 is exclusively coupled to site A7, and to site A6 when this site is present.

Well-regulated expression of all viral genes is critical for efficient viral replication and successful infection. The complex process of alternative splicing of HIV-1 mRNA is regulated primarily by the efficiencies at which the alternative splice sites of HIV-1 are utilized. Several features of suboptimal HIV-1 acceptor sites have been described, such as short and interrupted polypyrimidine tracts (O'Reilly *et al*, 1995; Si *et al*, 1997; Staffa *et al*, 1994), and irregular branch point sequences (Damier *et al*, 1997; Dyhr-Mikkelsen and Kjems, 1995). In addition, there is abundant evidence for the existence of several positive and negative *cis*-acting elements within the viral genome, which have been termed exonic splicing silencer (ESS), intronic splicing silencer (ISS), and exonic splicing enhancer (ESE) elements (Amendt *et al*, 1995; Caputi *et al*, 1999; Pongsoki *et al*, 2002; Tange *et al*, 2001). Members of the heterogeneous nuclear ribonucleo-protein (hnRNP) family such as hnRNP A/B or H selectively bind ESS and ISS elements and suppress the removal of adjacent introns (Bilodeau *et al*, 2001; Jacquenet *et al*, 2001a, 2001b). Members of the SR family of spliceosome factors have been shown to selectively bind to ESE elements and recruit additional cellular factors, and ultimately suppress nearby splicing silencers (Faustino and Cooper, 2003; Manley and Tacke, 1996). However, there have also been some recent reports that SR proteins are involved in negative regulation of viral pre-mRNA splicing, while the hnRNPs exert a positive effect on splicing (Graveley *et al*, 1999; Krecic and Swanson, 1999). Recently, it was shown that sequential removal of intronic sequences from the 5' end of the HIV-1 genome was critical for efficient viral gene expression (Bohne *et al*, 2005). However, the mechanism for this remains to be elucidated

VPR BINDS NUMEROUS PROTEIN PARTENERS IN HOST CELLS

Being a multi-functional protein, Vpr, is well known to modulate the host intracellular environment: A large body of research had revealed the ability of Vpr to delay or arrest cells in the G2 phase and induce the apoptosis of host cells. Numerous Vpr binding partners have subsequently been found in host cells. The glucocorticoid receptor (GR) (Mahalingam et al, 1998) and uracil DNA glycosylase (UNG) (Gragerov et al 1998, Withers-ward et al, 1997) had been reported to associate with Vpr and induce the G2 arrest. Another protein, Hsp70, has been proposed to inhibit Vpr induced G2 arrest, in contrast to other Vpr-binding partners (Iordanskiy, 2004). A plausible protein family, named DDB1- and Cullin 4A-associated factor (DCAF)-1 has been discovered. According to several laboratories reports, Vpr associate with a cul4A-containing E3 ligase complex via its interaction with DCAF-1 or DDB1. (Dehart et 2007, Le Rouzic et al, 2007, Schrofelbauer et al, 2007, Belzile et al, 2007). Moreover, it was suggested that Vpr induces checkpoint activation and G2 arrest by binding to the SAP145, a subunit of SF3b (Terada and Yasuda, 2006).

VPR IS REGULATOR OF CELLULAR pre-mRNA SPLICING

As mentioned above, SAP145 is an essential component of SF3b, a key factor in the spliceosome assembly. Even, Terada and Yasuda (2006) revealed that inhibition of SAP145 had no effect on the splicing of Cyclin B1, Ddk1 or cdc25, that are essential for cell cycle progression. Aida and colleagues do found that SAP145 is required for Vpr-induced inhibition of splicing of pre-mRNA. In their recent report by Hashizume they found that Vpr play a novel role as a regulator of pre-mRNA splicing both *in vivo* and *in vitro* (Kuramitsu *et al.*, 2005; Hshizume *et al.*, 2007) as follows. Given SAP145, an essential component of the SF3b factor in U2 snRNP, can bind to pre-mRNA and tethers U2 snRNP to the branch point site required for

complex A assembly (Champion-Arnaud and Reed, 1994), it has therefore been predicted that in addition physical interaction Vpr and SAP145 a splicing functional relationship exists. Likewise, it has been demonstrated that Vpr binds to ribonucleic acid via a process that requires the carboxy-terminal basic domain of the protein (de Rocquigny, 2002), which suggests the possibility of a functional association with pre-mRNA. Does Vpr control a splicing? As expected, our subsequent analyses has shown that Vpr inhibits the splicing of certain pre-mRNAs, such as *immunoglobulin M* pre-mRNA and α , β -*globin* pre-mRNA and caused the accumulation of incompletely spliced forms. In our view, because these experiments are performed during the time when Vpr is not able to induce significant G2 arrest, these novel effects of Vpr should be considered to occur via a pathway that is distinct from arrest of the cell cycle at G2. As is already our evidence, Vpr expressed from an HIV-1 provirus, in fact, was sufficient to inhibit splicing of α -*globin 2* pre-mRNA in infected cells.

VPR INHIBITS SELECTIVELY SPLICING OF THE HIV-1 GENOME

Interestingly, in Kuramitsu's research, it was proposed that the inhibition effect of Vpr on splicing is intron specific: the splicing of intron 1 of α -*globin 2* pre-mRNA was modestly inhibited by Vpr but the splicing of intron 2 was unaffected. Similarly, it was found Vpr regulated Env expression by interfering with *env* pre-mRNA splicing. Vpr selectively inhibit the removal of 3' intron of HIV-1 but not 5' intron, and subsequently increase the accumulation of *env* mRNA, which belongs to the singly spliced 4-kb class of mRNA. As a result the expression of Env protein is enhanced providing more antigenic Env to incorporated into virion and subsequently maintaining the high infectivity of the virus. (Zhang et al submitted).

PROPOSED MECHANISM FOR RNA PROCESSING REGULATED BY VPR

An extending investigation by Hashizume revealed that Vpr interacts with SAP145, resulting in the inhibition of binding between SAP145 and the splicing factor SAP49 which has two RNA recognition motifs that located on the surface of SF3b and interacts directly and highly specifically with both SAP145 and pre-mRNA within the essential splicing factor SF3b. This finding is in consistence with Terada and Yasuda's report. However, the interfering of Vpr to association between SAP145 and SAP49 may have more contribution to RNA processing rather than leading to G2 arrest, since, a Vpr mutant, named R80A, could bind to SAP145 and inhibit pre-mRNA processing but failed to induce G2 arrest (Hashizume, 2007).

It is conceivable Vpr may exert its splicing inhibition effect by interfering with the assembly of the spliceosome. Actually, in our recent research (Zhang et al., submitted), Vpr bound to U1-70K (a component of the U1 snRNP), U2AF65 (associated with the polypyrimidine tract), U2B'' (a component of the U2 snRNP), U1 and U2 snRNAs, as well as SAP145, indicating that Vpr associates with functional spliceosomes to regulate splicing. Here we proposed a possible model for Vpr mediated splicing regulation (Fig 1). SAP49 tethers U2 snRNP to the branch site. The U1 and U2 snRNPs are critical for spliceosome assembly, in which U1 snRNP is tightly associated with the 5' splice site, and U2AF65 directly contacts the 3' acceptor site of the pre-mRNA during Complex E assembly. Therefore, the interaction of Vpr with a component of the snRNP may impair the tethering of the U2 snRNP to the branch point site upstream of the acceptor site (Fig1). The Herpes Simplex Virus type 1 (HSV-1) ICP27 protein, an essential HSV-1 regulatory protein, has also been

reported to inhibit splicing prior to the first catalytic step through its association with SAP145 (Smith *et al.*, 2005, Bryant *et al.*, 2001), while it is likely that ICP27 was also inhibits pre-mRNA splicing from its interaction with SRPK1, a member of a family of conserved kinases that are highly specific for arginine/serine dipeptides, resulting in hypophosphorylation of Serine-arginine rich (SR) proteins, which impairs their role in spliceosome assembly (Smith *et al.*, 2005). NS1, an influenza virus non-structural protein, inhibits the splicing of the major class of mammalian pre-mRNAs (GU-AU introns) by binding to a specific stem-bulge in U6 snRNA, thereby blocking the formation of U4/U6 and U2/U6 complexes (Lu *et al.*, 1994; Qiu *et al.*, 1995). Therefore, these observations raise the possibility that Vpr may block spliceosome assembly by interfering with the function of the SAP145 and SAP49 complex in host cells.

Another question should be solved is that how the inhibition of splicing by Vpr is intron specific. Splicing of cellular *α -globin 2 pre-mRNA* removes 2 introns. Interestingly, only removal of intron 1 was modestly inhibited by Vpr. Also in our recent data, Vpr selectively increased the level of *env* mRNA by inhibiting HIV-1 *env* pre-mRNA splicing, that is, removal the 3' intron of HIV-1 primary transcripts. A splicing site can be regulated by consensus sequence up or downstream of the splice site, and splicing factors. HIV-1 A7 site is regulated by ESS3a, ESS3b, ISS and ESE3 (20), and many cellular splicing factors, such as the hnRNP family and SR family proteins, bind to these sites (21). It is still not clear whether or not the specificity of inhibition of splicing by Vpr is related to these elements or host factors. Vpr also displays high affinity for nucleic acids but no specific DNA sequence targeted by Vpr has been yet identified. Vpr might bind specific DNA sequence once associated with cellular partners to subsequently drive expression of both host cell and viral genes.

Further investigation to define whether there exist synergistic interactions between Vpr and consensus sequence(s) and splicing factors will help to better define the mechanism of pre-mRNA splicing inhibition by Vpr.

STRUCTURE OF VPR AND ITS ESSENTIAL DOMAINS FOR SPLICING REGULATION

An NMR structural analysis showed that full-length Vpr forms three amphipathic alpha helices folded around a hydrophobic core (Morellet et al, 2003). The N terminal domain is flexible and negatively charged, flanking to the helices. The C-terminal domain is also flexible, with positive charge and rich in arginine residues (Morellet, 2003). Two domains of the arginine-rich C-terminal domain from aa 77 to aa 96 and the third α -helical domain α H3, extending from aa 56 to aa 77 out of these domains, have been identified to be critical for inhibition of pre-mRNA splicing by Vpr (Hashizume, 2007). For example, Leu residue at position 67 in α H3 of Vpr, might be critical for Vpr to interact with SAP145 and exert its effect on splicing regulation, since replacement of Leu residue by Pro at position 67, which weakly retained its ability to induce G2 arrest and apoptosis (Nishizawa *et al.*, 2000) but lost its ability to target to the perinuclear region (Kamata and Aida., 2000; Kamata *et al.*, 2005), remarkably impaired the splicing inhibition, the interaction with SAP145 and the inhibition of SAP49 and SAP145 complex formation (Hashizume 2007). By contrast, although the arginine-rich C-terminal domain appeared to be involved in splicing inhibition, two Vpr mutants with a single mutation in this domain, R80A and R88A, are defective for G2 arrest but could bind SAP145 and inhibit mRNA splicing, indicating that Arg residues at positions 80 and 88 are dispensable for the splicing inhibition via an interaction between Vpr and SAP145. Thus, the association of Vpr to

SAP145 to inhibit pre-mRNA processing seemed to be independent of Vpr's G2 arrest function.

On the other hand, The CUS1 domain on SAP145 has been identified to be required to associate with Vpr. (Terada and Yasuda, 2006). On the other hand, the CUS1 domain on SAP145 has been identified to be required to associate with Vpr. (Terada and Yasuda, 2006). This CUS1 domain is highly conserved (Nurse P, 1990). Further investigation on the structure of binding between SAP145 and Vpr may provide us great information to understand the mechanism how Vpr regulate pre-mRNA processing.

A PROSPECTIVE ROLE IN FUTURE ANTIVIRAL THERAPY

Additional work is needed for the complete understanding of the novel role of Vpr. For instance, how the inhibitory effect of Vpr on RNA splicing can be intron specific; whether regulation of cellular mRNA processing benefit the virus replication and whether it closely related with G2 arrest or apoptosis of host cells. It is now well agreed that Vpr might be critical for pathogenesis of HIV-1 infection in vivo.

Vpr could potentially be an attractive target for the development of novel therapeutic strategies. Multiple host factors have been reported to interact with Vpr. The structural information of binding of Vpr with its partners may greatly facilitate the design of antiviral therapy. Development of compound that are potentially block the function of Vpr to induce G2 arrest or nuclear import are in process in several laboratories. The information of Vpr are involved in RNA splicing provide us a novel choice to screen antiviral drugs. Actually, a certain small-molecule chemical has been found to have anti-viral effect via inhibition of HIV pre-mRNA splicing. (Bakkour N

et al, 2007). The understanding of the novel role of Vpr may make great contribution to design a novel antiviral therapy.

Acknowledgements

Work in the author's Lab was supported in part by a Health Sciences Research Grant from the Ministry of Health, Labor and Welfare of Japan (Research on HIV/AIDS); and by the program for Promotion of Fundamental Studies in Health Sciences of the National Institute of Biomedical Innovation (NIBIO) of Japan. and by a President's Special Research Grant from RIKEN.

Amendt BA., Si ZH. and Stoltzfus CM. (1995) Presence of exon splicing silencers within human immunodeficiency virus type 1 tat exon 2 and tat-rev exon 3: evidence for inhibition mediated by cellular factors. *Mol Cell Biol.* **15**:6480.

Ayyavoo V, Mahboubi A, Mahalingam S, Ramalingam R, Kudchodkar S. Williams, WV, Green DR, Weiner DB. (1997) HIV-1 Vpr suppresses immune activation and apoptosis through regulation of nuclear factor kappa B. *Nat Med.* **3**:1117-1123.

Azuma A, Matsuo A, Suzuki T, Kurosawa T, Zhang X, Aida Y. (2006) Human immunodeficiency virus type 1 Vpr induces cell cycle arrest at the G1 phase and apoptosis via disruption of mitochondrial function in rodent cells. *Microbes Infect.* **8**:670-679.

Bakkour N, Lin YL, Maire S, Ayadi L, Mahuteau-Betzer F, Nguyen CH, Mettling C, Portales P, Grierson D, Chabot B, Jeanteur P, Branlant C, Corbeau P, Tazi J. (2007). Small-molecule inhibition of HIV pre-mRNA splicing as a novel antiretroviral therapy to overcome drug resistance. *PLoS Pathog.* **3**:1530-9

- Bryant HE, Wadd SE, Lamond AI, Silverstein SJ, Clements JB. (2001) Herpes simplex virus IE63 (ICP27) protein interacts with spliceosome-associated protein 145 and inhibits splicing prior to the first catalytic step. *J Virol.* 75:4376-4385.
- Belzile JP, Duisit G, Rougeau N, Mercier J, Finzi A, Cohen EA (2007) HIV-1 Vpr-mediated G2 arrest involves the DDB1-CUL4AVPRBP E3 ubiquitin ligase. *PLoS Pathog.* 3:0882-0893.
- Benko DM, Schwartz S, Pavlakis GN, Felber BK. (1990) A novel human immunodeficiency virus type 1 protein, tev, shares sequences with tat, env, and rev proteins. *J Virol*, 64:2505-2518.
- Bennett EM, Lever AM, Allen, J.F. (2004) Human immunodeficiency virus type 2 Gag interacts specifically with PRP4, a serine-threonine kinase, and inhibits phosphorylation of splicing factor SF2. *J Virol*, **78**:11303-11312.
- Berro R, Kehn K, de la Fuente C, Pumfery A, Adair R, Wade J, Colberg-Poley AM, Hiscott J, Kashanchi F. (2006) Acetylated Tat regulates human immunodeficiency virus type 1 splicing through its interaction with the splicing regulator p32. *J Virol*, **80**:3189-3204.
- Bilodeau PS, Domsic JK, Mayeda A, Krainer AR, Stoltzfus CM. (2001) RNA splicing at human immunodeficiency virus type 1 3' splice site A2 is regulated by binding of hnRNP A/B proteins to an exonic splicing silencer element. *J Virol*, **75**:8487-8497.
- Bohne J, Wodrich H, Krausslich HG. (2005) Splicing of human immunodeficiency virus RNA is position-dependent suggesting sequential removal of introns from the 5' end. *Nucleic Acids Res.* **33**:825-837.
- Bryant HE, Wadd SE, Lamond AI, Silverstein SJ, Clements JB. (2001) Herpes simplex virus IE63 (ICP27) protein interacts with spliceosome-associated

- protein 145 and inhibits splicing prior to the first catalytic step. *J Virol*, **75**:4376-4385.
- Caputi M, Mayeda A, Krainer AR, Zahler AM. (1999) hnRNP A/B proteins are required for inhibition of HIV-1 pre-mRNA splicing. *EMBO J*, **18**:4060-4067.
- Champion-Arnaud P, Reed R. (1994) The prespliceosome components SAP 49 and SAP 145 interact in a complex implicated in tethering U2 snRNP to the branch site. *Genes Dev*. **8**:1974-83.
- Damier, L., Domenjoud, L. and Branlant, C. (1997) The D1-A2 and D2-A2 pairs of splice sites from human immunodeficiency virus type 1 are highly efficient in vitro, in spite of an unusual branch site. *Biochem Biophys Res Commun*, **237**:182-187.
- DeHart JL, Zimmerman ES, Ardon O, Monteiro-Filho CM, Argañaraz ER, Planelles V (2007) HIV-1 Vpr activates the G2 checkpoint through manipulation of the ubiquitin proteasome system. *Virol J*. **4**:57
- de Noronha CM, Sherman MP, Lin HW, Cavrois MV, Moir RD, Goldman RD, Greene WC. (2001) Dynamic disruptions in nuclear envelope architecture and integrity induced by HIV-1 Vpr. *Science*. **294**:1105-1108.
- de Rocquigny H, Caneparo A, Delaunay T, Bischerour J, Mouscadet JF, Roques BP. (2000). Interactions of the C-terminus of viral protein R with nucleic acids are modulated by its N-terminus. *Eur J Biochem* **267**:3654-3660.
- Dyhr-Mikkelsen H, Kjems J. (1995) Inefficient spliceosome assembly and abnormal branch site selection in splicing of an HIV-1 transcript in vitro. *J Biol Chem*, **270**:24060-24066.
- Faustino NA, Cooper TA. (2003) Pre-mRNA splicing and human disease. *Genes Dev*, **17**:419-437.

- Gaynor ES, Chen IS (2001) Analysis of apoptosis induced by HIV-1 Vpr and examination of the possible role of the hHR23A protein. *Exp cell Res.* 267:243-57.
- Goh WC, Rogel ME, Kinsey CM, Michael SF, Fultz PN, Nowak MA, Hahn BH, Emerman M. (1998) HIV-1 Vpr increases viral expression by manipulation of the cell cycle: a mechanism for selection of Vpr in vivo. *Nat Med*, 4:65-71.
- Golas MM, Sander B, Will CL, Luhrmann R, Stark H. (2003) Molecular architecture of the multiprotein splicing factor SF3b. *Science.* 300:980-984.
- Gragerov A, Kino T, Ilyina-Gragerova G, Chrousos GP, Pavlakis GN (1998) HHR23A, the human homologue of the yeast repair protein RAD23, interacts specifically with Vpr protein and prevents cell cycle arrest but not the transcriptional effects of Vpr. *Virology.* 245:323-330.
- Graveley BR, Hertel KJ, Maniatis T. (1999) SR proteins are 'locators' of the RNA splicing machinery. *Curr Biol*, 9:R6-R7.
- Hashizume C, Kuramitsu M, Zhang X, Kurosawa T, Kamata M, Aida Y. (2007) Human Immunodeficiency Virus Type1 Vpr interacts with spliceosomal protein SAP145 to mediate cellular pre-mRNA splicing inhibition. *Microbes Infect.* 9: 490-497.
- Hastings ML, Krainer AR. (2001) Pre-mRNA splicing in the new millennium. *Curr Opin Cell Biol.* 13:302-309.
- Iijima S, Nitahara-Kasahara Y, Kimata K, Zhong Zhuang W, Kamata M, Isogai M, Miwa M, Tsunetsugu-Yokota Y, Aida Y. (2004) Nuclear localization of Vpr is crucial for the efficient replication of HIV-1 in primary CD4+ T cells. *Virology*, 327:249-261.
- Iordanskiy S, Zhao Y, Dubrovsky L, Iordanskaya T, Chen M, Liang D, Bukrinsky M

- (2004) Heat shock protein 70 protects cells from cell cycle arrest and apoptosis induced by human immunodeficiency virus type 1 viral protein R. *J Virol*. 78:9697-704
- Jacotot E, Ravagnan L, Loeffler M, Ferri KF, Vieira HL, Zamzami N, Costantini P, Druillennec S, Hoebeke J, Briand JP, Irinopoulou T, Daugas E, Susin SA, Cointe D, Xie ZH, Reed JC, Roques BP, Kroemer G. (2000) The HIV-1 viral protein R induces apoptosis via a direct effect on the mitochondrial permeability transition pore. *J Exp Med*. **191**:33-46.
- Jacquet S, Mereau A, Bilodeau PS, Damier L, Stoltzfus CM, Branlant C. (2001a) A second exon splicing silencer within human immunodeficiency virus type 1 tat exon 2 represses splicing of Tat mRNA and binds protein hnRNP H. *J Biol Chem*, **276**:40464-40475.
- Jacquet S., Ropers D, Bilodeau PS, Damier L, Mouglin A, Stoltzfus CM, Branlant C. (2001b) Conserved stem-loop structures in the HIV-1 RNA region containing the A3 3' splice site and its cis-regulatory element: possible involvement in RNA splicing. *Nucleic Acids Res*, **29**:464-478.
- Jenkins Y, McEntee M, Weis K, Greene WC. (1998) Characterization of HIV-1 vp r nuclear import: analysis of signals and pathways. *J Cell Biol*, **143**:875-885.
- Kamata M, Aida Y. (2000) Two putative alpha-helical domains of human immunodeficiency virus type 1 Vpr mediate nuclear localization by at least two mechanisms. *J Virol*, **74**:7179-7186.
- Kamata M, Nitahara-Kasahara Y, Miyamoto Y, Yoneda Y, Aida, Y. (2005) Importin-alpha promotes passage through the nuclear pore complex of human immunodeficiency virus type 1 Vpr. *J Virol*, **79**:3557-3564.
- Krämer A. (1996). The structure and function of proteins involved in mammalian pre-

- mRNA splicing. *Annu Rev Biochem.* 65:367-409.
- Krecic AM, Swanson MS. (1999) hnRNP complexes: composition, structure, and function. *Curr Opin Cell Biol.* 11:363-371.
- Kuramitsu M, Hashizume C, Yamamoto N, Azuma A, Kamata M, Tanaka Y, Aida Y. (2005) A novel role for Vpr of human immunodeficiency virus type 1 as a regulator of the splicing of cellular pre-mRNA. *Microbes Infect.* 7:1150-1160.
- Le Rouzic E, Belaïdouni N, Estrabaud E, Morel M, Rain JC, Transy C, Margottin-Goguet F (2007) HIV1 Vpr arrests the cell cycle by recruiting DCAF1/VprBP, a receptor of the Cul4-DDB1 ubiquitin ligase. *Cell Cycle.* 6:182-188.
- Le Rouzic E, Morel M, Ayinde D, Belaïdouni N, Letienne J, Transy C, Margottin-Goguet F. (2008). Assembly with the Cul4A-DDB1DCAF1 ubiquitin ligase protects HIV-1 Vpr from proteasomal degradation. *J Biol Chem.* 283:21686-21692
- Lu Y, Qian XY, Krug RM. (1994). The influenza virus NS1 protein: a novel inhibitor of pre-mRNA splicing, *Genes Dev.* 8:1817-1828.
- Lum JJ, Cohen OJ, Nie Z, Weaver JG, Gomez TS, Yao XJ, Lynch D, Pilon AA, Hawley N, Kim JE, Chen Z, Montpetit M, Sanchez-Dardon J, Cohen EA, Badley AD. (2003). Vpr R77Q is associated with long-term nonprogressive HIV infection and impaired induction of apoptosis. *J Clin Invest.* 111:1547-54
- Mahalingam S, Ayyavoo V, Patel M, Kieber-Emmons T, Kao GD, Muschel RJ, Weiner DB. (1998). HIV-1 Vpr interacts with a human 34-kDa mov34 homologue, a cellular factor linked to the G2/M phase transition of the mammalian cell cycle. *Proc Natl Acad Sci USA.* 95:3419-3424.
- Manley JL, Tacke R. (1996). SR proteins and splicing control. *Genes Dev,* 10:1569-1579.

- Morellet N, Bouaziz S, Petitjean P, Roques BP. (2003) NMR structure of the HIV-1 regulatory protein VPR. *J Mol Biol.* 327:215-27.
- Neil SJ, Zang T, Bieniasz PD. (2008). Tetherin inhibits retrovirus release and is antagonized by HIV-1 Vpu. *Nature.* 451:425-30.
- Nie Z, Bergeron D, Subbramanian RA, Yao XJ, Checroune F, Rougeau N, Cohen EA.. (1998). The putative alpha helix 2 of human immunodeficiency virus type 1 Vpr contains a determinant which is responsible for the nuclear translocation of proviral DNA in growth-arrested cells. *J Virol.* 72:4104-15.
- Nishizawa M, Kamata M, Katsumata R, Aida Y. (2000a). A carboxy-terminally truncated form of the human immunodeficiency virus type 1 Vpr protein induces apoptosis via G1 cell cycle arrest. *J Virol.* **74**:6058-6067.
- Nishizawa M, Kamata M, Mojin T, Nakai Y, Aida Y. (2000b). Induction of apoptosis by the Vpr protein of human immunodeficiency virus type 1 occurs independently of G2 arrest of the cell cycle. *Virology.* **276**:16-26.
- Nishizawa M, Myojin T, Nishino Y, Nakai Y, Kamata M, Aida Y. (1999). A carboxy-terminally truncated form of the Vpr protein of human immunodeficiency virus type 1 retards cell proliferation independently of G2 arrest of the cell cycle. *Virology.* **263**:313-322.
- Nitahara-Kasahara Y, Kamata M, Yamamoto T, Zhang X, Miyamoto Y, Muneta K, Iijima S, Yoneda Y, Tsunetsugu-Yokota Y, Aida Y. (2007). Novel nuclear import of Vpr promoted by importin alpha is crucial for human immunodeficiency virus type 1 replication in macrophages. *J Virol.* 81:5284-5293.
- Nurse P. (1990). Universal control mechanism regulating onset of M-phase. *Nature.* 344:503-508.

- O'Reilly MM, McNally MT, Beemon KL. (1995). Two strong 5' splice sites and competing, suboptimal 3' splice sites involved in alternative splicing of human immunodeficiency virus type 1 RNA. *Virology*, **213**:373-385.
- Pongoski J, Asai K, Cochrane A. (2002). Positive and negative modulation of human immunodeficiency virus type 1 Rev function by cis and trans regulators of viral RNA splicing. *J Virol*, **76**:5108-5120.
- Poon B, Grovit-Ferbas K, Stewart SA, Chen IS. (1998) Cell cycle arrest by Vpr in HIV-1 virions and insensitivity to antiretroviral agents. *Science*, **281**:266-269.
- Popov S, Rexach M, Zybarth G, Reiling N, Lee MA, Ratner L, Lane CM, Moore MS, Blobel G, Bukrinsky M. (1998) .Viral protein R regulates nuclear import of the HIV-1 pre-integration complex. *EMBO J*. 17: 909-917.
- Purcell DF, Martin MA. (1993). Alternative splicing of human immunodeficiency virus type 1 mRNA modulates viral protein expression, replication, and infectivity. *J Virol*. **67**:6365-6378.
- Qiu Y, Nemeroff M, Krug RM. (1995). The influenza virus NS1 protein binds to a specific region in human U6 snRNA and inhibits U6-U2 and U6-U4 snRNA interactions during splicing. *RNA*. 1:304-316.
- Schröfelbauer B, Hakata Y, Landau NR. (2007). HIV-1 Vpr function is mediated by interaction with the damage-specific DNA-binding protein DDB1. *Proc Natl Acad Sci USA*. 104:4130-5
- Schwartz S, Felber BK, Benko DM, Fenyo EM, Pavlakis GN. (1990). Cloning and functional analysis of multiply spliced mRNA species of human immunodeficiency virus type 1. *J Virol*, **64**:2519-2529.
- Sheehy AM, Gaddis NC, Malim MH. (2003). The antiretroviral enzyme APOBEC3G is degraded by the proteasome in response to HIV-1 Vif. *Nat Med*. 9:1404-

1407.

- Sherman MP, de Noronha, CM, Pearce D, Greene WC. (2000) Human immunodeficiency virus type 1 Vpr contains two leucine-rich helices that mediate glucocorticoid receptor coactivation independently of its effects on G2 cell cycle arrest. *J Virol*, **74**:8159-8165
- Si, Z, Amendt BA, Stoltzfus CM. (1997) Splicing efficiency of human immunodeficiency virus type 1 tat RNA is determined by both a suboptimal 3' splice site and a 10 nucleotide exon splicing silencer element located within tat exon 2. *Nucleic Acids Res*, **25**:861-867.
- Smith RWP, Malik P, Clements JB. (2005). The herpes simplex virus ICP27 protein: a multifunctional post-transcriptional regulator of gene expression, *Biochem. Soc. Trans.* 33:499-501.
- Staffa A, Cochrane A. (1994) The tat/rev intron of human immunodeficiency virus type 1 is inefficiently spliced because of suboptimal signals in the 3' splice site. *J Virol.* **68**:3071-3079.
- Staknis D, Reed R. (1994) Direct interactions between pre-mRNA and six U2 small nuclear ribonucleoproteins during spliceosome assembly. *Mol Cell Biol.* 14:2994-3005.
- Stewart SA., Poon B, Song JY, Chen IS. (2000). Human immunodeficiency virus type 1 vpr induces apoptosis through caspase activation. *J Virol.***74**:3105-3111.
- Tang H, Kuhen KL, Wong-Staal F. (1999). Lentivirus replication and regulation. *Annu Rev Genet*, **33**:133-170.
- Tange TO, Damgaard CK, Guth S, Valcarcel J, Kjems J. (2001). The hnRNP A1 protein regulates HIV-1 tat splicing via a novel intron silencer element. *EMBO J*, **20**:5748-5758.

- Terada Y and Yasuda Y. (2006). Human immunodeficiency virus type 1 Vpr induces G2 checkpoint activation by interacting with the splicing factor SAP145. *Mol Cell Biol.* 26(21):8149-8158.
- Will CL, Schneider C, MacMillan AM, Katopodis NF, Neubauer G, Wilm M, Luhrmann R, Query CC. (2001). A novel U2 and U11/U12 snRNP protein that associates with the pre-mRNA branch site. *EMBO J.* 20:4536-4546.
- Withers-Ward ES, Jowett JB, Stewart SA, Xie YM, Garfinkel A, Shibagaki Y, Chow SA, Shah N, Hanaoka F, Sawitz DG, Armstrong RW, Souza LM, Chen IS. (1997). Human immunodeficiency virus type 1 Vpr interacts with HHR23A, a cellular protein implicated in nucleotide excision DNA repair. *J Virol.* 71:9732-42.
- Zhang X, Hashizume C, Kuramitsu M, Azuma A, Isogai M, Kimata K, Matsuda G, Suzuki T, Yamamoto N, Tanaka Y, Yamamoto N, Aida Y. The Vpr protein enhances envelope expression and viral infectivity via selective inhibition of HIV-1 splicing, submitted.
- Zhu J, Mayeda A and Krainer AR (2001) Exon identity established through differential antagonism between exonic splicing silencer-bound hnRNP A1 and enhancer-bound SR proteins. *Mol Cell* 8:1351-1361.

Figure legend:

Vpr inhibit RNA splicing via interaction with SAP145. Normally when initiate the assembly of the spliceosome at the early step of RNA splicing, the 5' splice site of an intron is recognized by the U1 small nuclear ribonucleoprotein (snRNP), followed by the binding of U2 auxiliary factor (U2AF) to 3' splice site and its upstream polypyrimidine tract and form a complex E, then U2AF recruit the u2snRNP, base pair the U2RNA with branch point to complete the formation of complex A, among recruited U2RNP, SAP145 bind SAP49 to form the subunit of SF3b. Subsequent, binding of the U4-U5-U6 snRNP and form a large spliceosome, which perform a reaction resulting in the ligation of the exons and excision of the intron. (left), however, in the presence of Vpr, the binding of Vpr with SAP145 block the recruitment of SAP49 and result in failure of catalytic spliceosome formation. Subsequently inhibit the splicing reaction (right).

