



HOKKAIDO UNIVERSITY

Title	The effect of climate change on the growth of Japanese chum salmon (<i>Oncorhynchus keta</i>) using a bioenergetics model coupled with a three-dimensional lower trophic ecosystem model (NEMURO)
Author(s)	Kishi, Michio J.; Kaeriyama, Masahide; Ueno, Hiromichi et al.
Citation	Deep Sea Research Part II: Topical Studies in Oceanography, 57(13-14), 1257-1265 https://doi.org/10.1016/j.dsr2.2009.12.013
Issue Date	2010-07
Doc URL	https://hdl.handle.net/2115/43182
Type	journal article
File Information	Kishi.pdf



The effect of climate change on the growth of Japanese chum salmon
(*Oncorhynchus keta*) using a bioenergetics model coupled with a
three-dimensional lower trophic ecosystem model (NEMURO)

Michio J. Kishi^{1,2}, Masahide Kaeriyama¹
Hiromichi Ueno¹, Yasuko Kamezawa^{3,4} ^{O A}

1: Faculty of Fisheries Sciences, Hokkaido University

2: Frontier Research System of Global Change, JAMSTEC

3: Graduate School of Environmental Science, Hokkaido University

4: AES, Co., Ltd

Abstract

From the 1970s to 1990s, a reduction in the body size of Japanese chum salmon (*Oncorhynchus keta*) was observed. To investigate this body size reduction in the North Pacific, we developed a bioenergetics model for chum salmon coupled with the results from a lower trophic ecosystem model embedded into a three-dimensional global model. In the bioenergetics model, respiration and consumption terms are assumed to be functions of water temperature and prey zooplankton density, which are the determining factors of the reduction of body size. The model reproduced the body size of the 1972 and 1991 year classes of chum salmon. The reproduced body size of the 1972 year class was larger than that of 1991 year class, and this result agrees with observations from the Bering Sea. Our model also reproduced the body size trend from 1970 to 2000. The prey density, especially in the eastern North Pacific, had a greater influence on the change of body size than did the SST. This suggests that the size reduction of Japanese chum salmon in the 1990s was partly affected by changes in prey zooplankton density. In the context of the global warming scenario, we discuss changes in the migration route of chum salmon and predict that the population of Japanese chum salmon experience significant declines over this century.

Keywords: bioenergetics model, Japanese chum salmon, size reduction, NEMURO, global warming

1. Introduction

Chum salmon (*Oncorhynchus keta*) are distributed widely in the Northern Pacific and can be divided into North American and Asian groups. The latter can be divided further into Russian and Japanese groups, which show different migration routes (Neave et al., 1980). The stock of the Japanese group is maintained by artificial release (Kaeriyama, 1999, Hiroi, 1998). Hiroi (1998) and Ishida et al. (1993) pointed out the decrease of body size of Japanese chum salmon in 1970s and 1980s. According to Azumaya and Ishida (2004) and Urawa (2000), in spring the released chum salmon in Hokkaido spend three months at the coast of the Okhotsk Sea, stay in the Okhotsk Sea until late fall, then migrate toward the western North Pacific (WNP), and by the next summer migrate into the Bering Sea (BS). In the BS, they prey mainly on zooplankton and increase their weight. In the fall they move into the Gulf of Alaska in the eastern North Pacific (ENP) to spend the winter. They repeat the BS/ENP migration 3-4 times, meaning for 3-4 years, and then when they are ready to mature in the summer of their fourth/fifth year they return to their home rivers in Hokkaido, Japan.

Kaeriyama (1998), Kaeriyama et al. (2004), and Azumaya and Ishida (2000) pointed out that mature salmon are mainly 4 years old (i.e., an ocean age of 3 years), and differences in growth (wet weight and growth rate) can be observed in fish older than 2 years in the Bering Sea. They suggested that the differences in body weight might arise from variability in prey density, the density effect, and water temperature. Morita et al. (2001) analyzed the correlation between salmon growth and the physical environment and described the positive correlation between the water temperature anomaly and the body weight anomaly.

The back-calculated growth rates of juvenile Hokkaido chum salmon obtained from their scales showed that their growth in the Okhotsk Sea increased in the 1990s (Fig. 1). These growth anomalies were strongly positively correlated with sea surface temperature (SST) in the Okhotsk Sea during summer and fall, but SST was negatively correlated with the area covered by sea ice during winter (Kaeriyama et al., 2007). Zooplankton biomass in the Okhotsk Sea has been decreasing since the 1980s (Shuntov and Dulepova, 1996). Therefore, the increase in growth of Hokkaido chum salmon during the 1990s appears to have been affected by the increase in SST and not directly by the decrease in zooplankton biomass in the Okhotsk Sea (Kaeriyama et al., 2007). Sea ice concentrations also have been decreasing during the last 100 years as air temperatures along the Okhotsk Sea coast of Hokkaido have increased (Aota, 1999). Aota (1999) suggested that this phenomenon could be a symptom of global warming. Therefore, increases in the growth of Hokkaido chum salmon in the Okhotsk Sea since

the 1990s might also be related to effects of global warming.

Return rates of chum salmon released from south Korea and Iwate Prefecture, which is located at the northern part of Honshu island, Japan, have decreased since the 1990s, whereas those of Hokkaido chum salmon have increased since the 1976 regime shift (Fig. 2a). In Hokkaido, the chum salmon populations from Nemuro City and the Okhotsk Sea coasts have had return rates that were markedly higher than those of other populations (Fig. 2b). Korean and Iwate chum salmon are distributed in the southern waters, which are influenced by the Tsushima Warm Current during the spring offshore-migration period. Declines in their return rates did not coincide with years of climate regime shifts (1976, 1988, and 1998). These results suggest that global warming can be expected to have positive effects on Hokkaido chum salmon populations in the Okhotsk Sea, but negative effects on Korean and Iwate populations in the Tsushima Warm Current.

In this study, we examined the causes of the observed growth difference of adult chum salmon using a bioenergetics model coupled with the output from the 3-D NEMURO model (NEMURO coupled with three dimensional physical global model, Aita et al., 2007). We also discuss the effect of water temperature increase on chum salmon survival in the near future.

2. Method

The bioenergetics fish model we used follows Rudstam (1988), Ware (1978), Beauchamp et al. (1989) and Trudel et al. (2004). Kamezawa et al. (2007) modified the bioenergetics model for chum salmon. The model includes temperature-dependent and food concentration-dependent terms. Aita et al. (2007) calculated time-dependent features of water temperature and lower trophic compartments using an ecological-physical coupled global 3-D model, and they used NEMURO (Kishi et al., 2007) as an ecological model. Our bioenergetics model used output data from Aita et al. (2007) that was generated by 3-D NEMURO.

In our discussion of the global warming scenario, we used Kawamiya et al.'s (2005) SST (Sea Surface Temperature) data, which predicted physical fields under the SRES A2 scenario. As the predicted SST has bias due to the technical reason, the predicted anomaly of SST was added to the present SST (<http://goos.kishou.go.jp>).

2.1 Migration

Figure 3 shows a schematic view of the three oceanic spatial domains and the numbers of life stages based on Urawa (2000). The stages are shown in Table 1 after

Kamezawa et al., (2007). In our model, chum salmon migrate among the boxes shown in Fig.3 according to stage transitions shown in Table 1 instantaneously.

2.2 Bioenergetics model

The following equation describes individual fish growth of chum salmon:

$$\frac{dW}{dt} = [C - (R + S + F + E)] \cdot \frac{CAL_z}{CAL_f} \cdot W \quad (1)$$

where C is consumption ($\text{g prey} \cdot \text{g fish}^{-1} \cdot \text{d}^{-1}$), E is excretion or losses of nitrogenous excretory wastes ($\text{g prey} \cdot \text{g fish}^{-1} \cdot \text{d}^{-1}$), F is egestion or losses due to feces ($\text{g prey} \cdot \text{g fish}^{-1} \cdot \text{d}^{-1}$), R is respiration or losses through metabolism ($\text{g prey} \cdot \text{g fish}^{-1} \cdot \text{d}^{-1}$), S is specific dynamic action or losses due to energy costs of digesting food ($\text{g prey} \cdot \text{g fish}^{-1} \cdot \text{d}^{-1}$), and CAL_z / CAL_f is the conversion rate between prey and fish. The individual processes are described below. Consumption and respiration are nonlinear functions of fish weight and water temperature. In addition to the physiological parameters, the model requires information about the caloric content of herring (this is seasonal) and prey, diet composition, typical prey densities, and water temperature.

Consumption (C)

Consumption is estimated as the proportion of maximum daily ration for chum salmon at a particular mass and temperature. The maximum daily consumption rate (g of prey per g body mass of salmon per day) is estimated using an allometric function of mass from *ad libitum* feeding experiments conducted at the optimum temperature.

The basic form of the consumption function is:

$$C = C_{MAX} \cdot \rho \cdot f_c(T) \quad (2)$$

$$C_{MAX} = ac \cdot W^{bc} \quad (3)$$

$$\rho = \frac{v_i}{K_i + PD \cdot v_i} \quad (4)$$

where C_{max} is the maximum specific consumption rate ($\text{g prey} \cdot \text{g fish}^{-1} \cdot \text{d}^{-1}$), $f_c(T)$ is a temperature-dependence function for consumption, T is water temperature ($^{\circ}\text{C}$), W is salmon mass (g wet weight), and C_{MAX} is the consumption rate ($\text{g} \cdot \text{g}^{-1} \cdot \text{d}^{-1}$) of an individual predator i , the values of which were determined following Beauchamp et al. (1989) (Table 2). PD is the density of prey (g weight m^{-3}) and v_i is the vulnerability at

stage i , which was determined to be 1 for all stages. K_i is the half saturation constant (g prey m^{-3}), which was determined to be 0.55 referring that of Pacific saury by Ito et al. (2004) (Table 2). a_c is the intercept of the mass-dependence function for a 1 g fish at the optimum water temperature and b_c is the coefficient of the mass dependence.

Thornton and Lessem's (1978) description of temperature dependence is essentially the product of two sigmoid curves – one fit to the increasing portion of the temperature-dependence function ($gcta$) and the other to the decreasing portion ($gctb$).

$$f_c(T) = gcta \cdot gctb \quad (5)$$

where

$$tt5 = \frac{1}{(te2 - te1)} \quad (6)$$

$$t5 = tt5 \cdot a \log \left[0.98 \cdot \frac{(1.0 - xk1)}{(0.02 \cdot xk1)} \right] \quad (7)$$

$$t4 = e^{[t5(T - te1)]} \quad (8)$$

$$tt7 = \frac{1}{(te4 - te3)} \quad (9)$$

$$t7 = tt7 \cdot a \log \left[0.98 \cdot \frac{(1.0 - xk4)}{(0.02 \cdot xk4)} \right] \quad (10)$$

$$t6 = e^{[t7(te4 - T)]} \quad (11)$$

$$gcta = \frac{(xk1 \cdot t4)}{(1.0 + xk1(t4 - 1.0))} \quad (12)$$

$$gctb = \frac{(xk4 \cdot t6)}{(1.0 + xk4(t6 - 1.0))} \quad (13),$$

and where $te1$ is the lower temperature at which the temperature dependence is a small fraction, $xk1$, of the maximum rate and $te2$ is the water temperature corresponding to $xk2$ of the maximum consumption rate in increasing portion of the curve. For the decreasing portion of the curve, $te3$ is the water temperature ($\geq te2$) at which dependence is $xk3$ of the maximum, and $te4$ is the temperature at which dependence is some reduced fraction $xk4$ of the maximum rate. Welch et al. (1995) showed that salmon in the ENP occur where SST is below 10.4°C in spring and between 8°C and 12°C in summer. Nagasawa (2000) reported that salmon are distributed between 3.9°C and 5.0°C in the WNP and between 5.2°C and 6.7°C in the central Pacific. Consequently, we chose the parameter values shown in Table 2 to use in our model.

Respiration and metabolism (R)

The respiration rate is the amount of energy used for routine metabolism and it depends on body weight, ambient temperature, and activity (swimming speed) (Trudel et al., 2004; Ware, 1978). The equations below illustrate how R is calculated.

$$R = (R_s + R_a) \cdot 5.258 \quad (14)$$

$$R_s = a_{rs} \cdot W^{br} \cdot \exp (cr \cdot T) \quad (15)$$

$$R_a = a_{ra} \cdot W^{dr} \cdot U^{er} \quad (16)$$

$$U = a_u \cdot W^{bu} \quad (17)$$

$$\left(\frac{13560 \text{ joules}}{gO_2} \cdot \frac{1cal}{4.18 \text{ joules}} \right) \div \left(\frac{2580 \text{ joules}}{g \text{ zoop}} \cdot \frac{1cal}{4.18 \text{ joules}} \right) = 5.258 \text{ g zoop} / gO_2$$

where

R_s : standard metabolism

a_{rs} : Oxygen consumption rate at 0°C per 1 g wet weight which depends on temperature

R_a : metabolism when the fish moves

a_{ra} : Oxygen consumption rate at 0°C per 1 g wet weight which depends on activity

U : swimming speed

5.258 : conversion rate from (g O_2 g fish⁻¹ day⁻¹) to (g prey g fish⁻¹ day⁻¹) (Megrey et al., 2002)

zoop :zooplankton biomass

br, cr, dr, er, au, bu: constants defined in Table 2.

Specific dynamic action (SDA) = losses due to energy costs of digesting food, egestion (F), and excretion(E)

These three terms can be calculated as follows,

$$SDA = S \cdot (C - F) \quad (19)$$

$$F = a_F \cdot C \quad (20)$$

$$E = a_E \cdot (C - F) \quad (21)$$

SDA is the proportion of assimilated energy lost to specific dynamic action, C is the specific consumption rate (g·g⁻¹·d⁻¹), and F is the specific egestion rate (g·g⁻¹·d⁻¹). Egestion (F , fecal waste) and excretion (E , nitrogenous waste) can be computed as a constant proportion of consumption.

S, a_F, a_E : constant

These values used in our model were taken from Ito et al. (2004), in which they were determined for Pacific saury. Although the ecology and physiology of chum salmon differ from those of saury, these values are supported to be useful to the other species in the Northern Pacific (Megrey et al., 2007a,b; Rose et al., 2007), thus we used the same values.

2.3 NEMURO ecosystem model

The prey density and water temperature values in our bioenergetics model were obtained from Aita et al. (2007), who calculated a 3-D physical-ecological coupled model (3-D NEMURO) for the northern Pacific from 1948 to 2002. We used the daily averaged output for the upper 20 m of the water column and also averaged spatially within each box. In the Gulf of Alaska, chum salmon prefer jelly fish rather than copepods (Kaeriyama et al., 2004), and we assumed predatory zooplankton (PZ) to be prey: in NEMURO, ZP includes jerry fish, salps, and/or krill (Kishi et al., 2007). NEMURO describes zooplankton density as nitrogen density (mol N l^{-1}) and we converted it into wet weight following Megrey et al., 2002)

$$\frac{14\mu\text{ gN}}{\mu\text{ moleN}} \cdot \frac{1.0\text{e}^{-6}\text{ g}}{\mu\text{ g}} \cdot \frac{1\text{gdry weight}}{0.07\text{Ndry weight}} \cdot \frac{1\text{g wet weight}}{0.2\text{dry weight}} \cdot \frac{1.0\text{e}^3\text{ liters}}{\text{m}^3} \quad (22)$$

However, the converted value is smaller than the observed value (Kamezawa et al. 2007). Kamezawa et al. (2007) converted ZP from NEMURO based on WOA (World Ocean Atlas) 2003 and NPAFC (North Pacific Anadromous Fish Commission) 2004 as follows:

$$\frac{\text{observational zooplankton value}}{\text{ZP value Aita et al. 2007}} = 1.937 \quad (23)$$

$$\text{ZP} = \text{ZP} \times 1.937 \quad (24)$$

In our global warming scenario, we used predicted water temperature taken from Kawamiya et al. (2005), who simulated physical fields according to the IPCC IS92a global warming scenario. We used the outputs from Kawamiya et al. (2005) to simulating responses of Japanese chum salmon to global warming.

2.4 Time steps for the bioenergetics model

Urawa (2000) suggested that the inter-annual variations in wet weight of chum salmon can be observed after they migrate into the WNP from the Okhotsk Sea. In our analysis, we excluded the juvenile period spent in the Okhotsk Sea and begin the

simulation at age 404 days when the fish are in the WNP. The time step used was 3600 sec and zooplankton density and water temperature were taken from Aita et al. (2007).

2.5 Global warming scenario

Welch et al. (1998) suggested that the geographic distribution of sockeye salmon in the BS will shrink if the atmospheric CO₂ base doubles based on the present relationship between the CPUE (Catch Per Unit Effort) and SST. We introduce a similar procedure for chum salmon. The survival of chum salmon released into the Hokkaido River is well correlated with the water temperature of the Okhotsk Sea while the fish remain there (Kaeriyama et al., 2007). In addition, the optimal temperature for chum salmon in the BS can be estimated based on the relationship between CPUE and SST (Ueno and Ishida, 1996), and Fukuwaka et al. (2007) estimated the optimal temperature for the salmon's time spent in the NWP during winter. Based on the above literature, we propose the following criteria for the survival of chum salmon of Hokkaido stock;

Growth, Feeding : 8–12 : 1st year in the Okhotsk Sea (July–October)

Wintering: 4–6 : 1st year in WNP (November–June)

Growth, Feeding: 8–12 : 2nd year in the Bering Sea (July–October)

Wintering: 4–6 : 2nd year in the Gulf of Alaska (November–June)

3. Results and discussion

Figure 4 shows the time series of the simulated body weight of chum salmon for the 1972 year class (a) and the 1991 year class (b) together with observed body weight (Kamezawa et al., 2007). Figure 4-(c) shows the difference in wet weight between the 1972 and 1991 year classes for the control case (thin line), for the constant prey with time-dependent temperature case (bold line), and for the constant temperature with time-dependent prey case (dashed line). The time derivative of growth in summer (April to November) is larger than that in winter (December to March) in each case, because the growth rate becomes high due to high temperature in this area. And the difference in growth between these two year classes was caused by differences in prey density in the ENP during winter to spring (from February to July) of ocean age 3 and 4 fish. Figure 5 shows the time-dependent features of body weight of chum salmon in the BS in summer from 1971 to 1999, a) **observed** and b) **simulated** results. Simulated results describe the observed features well: 1) Ocean age 1 does not exhibit interannual variation; 2) from ocean ages 2 to 4, interannual variation of body weight is evident; and 3) the data show a log trend of decrease from 1970s through the 1990s and an increase in the late 1990s. The correlation coefficients between the observed and the

simulated results are -0.19, 0.45, 0.43 and 0.65 for ocean age of 1, 2, 3 and 4, respectively (significant). The inter-annual variations in body weight were caused mainly by differences in prey density, which were brought about by interannual variation in climate change, like PDO (Pacific Decadal Oscillation). According to Aita et al., (2007), PDO brought about decadal zooplankton/phytoplankton density anomaly. The present model does not include feedback from feeding pressure by salmon to the lower trophic ecosystem, however, we should include top-down effects in future. Consequently we can conclude that inter-annual climate change brought about the change of prey density in the Gulf of Alaska, and the inter-annual body weight anomaly of chum salmon was caused. This means that without a competition among species, we can explain the time dependent change of body size of chum salmon to some extent.

We pointed out that the prey density plays important role in the wet weight of chum salmon described above, however, as pointed in section 2.5, sea surface temperature must also play an important role in the distribution area. The Intergovernmental Panel on Climate Change (IPCC) concluded in its Fourth Assessment Report of 2007 that global warming caused by human activity is indeed taking place. What impact will global warming have on salmon? We predicted the situation for chum salmon in 2050 and 2095 based on the IPCC SRES-A1B scenario using their optimal temperature. The hatched portions of Figures 6 and 7 represent the areas in which chum salmon can survive based on the estimated SST increase. The survival conditions are described in section 2.5. Our model predicts that (1) in the North Pacific, the habitat in eastern waters (i.e., the Gulf of Alaska) decreases significantly, and chum salmon distribution area moves northward and increases in parts of the Arctic Ocean during summer; the carrying capacity of chum salmon would decrease specifically in the subarctic area. (2) Hokkaido chum salmon could lose their migration route to the Sea of Okhotsk by 2050, and the population experience significant declines over this century. (3) In the BS, the carrying capacity of chum salmon would significantly decrease by 2050, it encounters strict density-dependence effects, and the population would lose its distribution area by 2100.

4. Future works

In the bioenergetics model, we assumed the date of migration to be determined by calendar date, but this assumption does not consider the ecological demands of the fish. We should include in the model the relationship between growth and migration date. We did not include the density effect of salmon either. Tadokoro et al. (1996) pointed out that there is bimodal competition for prey between chum salmon and pink

salmon, and this effect also must be included in the next step. In our global warming scenario, we used averaged water temperature in the top 20 m of the water column, but salmon might opt for deeper depths when swimming and migrating and only use shallower depths when feeding. This kind of change of ecological behavior is not considered in our discussions. Moreover, chum salmon might adapt to warmer temperatures. Consequently, this paper analyzed environmental effects on chum salmon growth and only suggests possible outcomes for the future.

Acknowledgement

We would like to express our sincere thanks to Dr. Kawamiya and Ms. Aita of Frontier Research System for Global Change for providing their numerical data.

References

- Aita, M. N., Yamanaka, Y. and Kishi, M.J., 2007. Interdecadal variation of the lower trophic ecosystem in the Northern Pacific between 1948 and 2002, in a 3-D implementation of the NEMURO model. *Ecol. Modell.*, **202**, 81–94.
- Aota, M., 1999. Long-term tendencies of sea ice concentration and air temperature in the Okhotsk Sea coast of Hokkaido. *PICES Sci. Rep.*, **12**, 1–2.
- Azumaya, T. and Ishida, Y., 2000. Density interactions between pink salmon (*Oncorhynchus gorbuscha*) and chum salmon (*O. keta*) and their possible effects on distribution and growth in the North Pacific Ocean and Bering Sea. *N. Pac. Anadr. Fish Comm. Bull.*, **2**, 165–174.
- Azumaya, T and Ishida, Y., 2004. An evaluation of the potential influence of SST and currents on the oceanic migration of juvenile and immature chum salmon (*Oncorhynchus keta*) by a simulation model. *Fish. Oceanogr.*, **13**, 10–23.
- Beauchamp, D.A., Stewart D.J. and Thomas G.L., 1989. Corroboration of abioenergetics model for sockeye salmon. *Trans. Am. Fish. Soc.*, **118**, 597–607.
- Fukuwaka, M., Sato, S. Takahashi, S., Onuma, T., Sakai, O., Tanimata, N., Makino, K., Davis, N.D., Volkov, A.F., Seong, K.B. and Moss, J.H., 2007. Winter distribution of chum salmon related to environmental variables in the North Pacific. *N. Pac. Anadr. Fish Comm.*, Technical Report No. 7, 29–30
- Hiroi, O., 1998. Historical trends of salmon fisheries and stock conditions in Japan. *N. Pac. Anadr. Fish Comm. Bull.*, **1**, 23–27.
- Ishida, Y, Ito, S. Kaeriyama, M. McKinnel, S. Nagasawa, K., 1993. Recent changes in age and size of chum salmon (*Oncorhynchus keta*) in the North Pacific Ocean and possible causes. *Can. J. Fish. Aquat. Sci.*, **50**, 290-295.
- Ito, S., Kishi, M.J., Kurita, Y., Oozeki, Y., Yamanaka, Y., Megrey, B.A. and Werner, F.E., 2004. Initial design for a fish bioenergetics model of Pacific saury coupled to a lower trophic ecosystem model. *Fish. Oceanogr.*, **13**, 111–124.

Kaeriyama, M., 1998. Dynamics of chum salmon , *Oncorhynchus keta* , populations released from Hokkaido, Japan. N. Pac. Anadr. Fish Comm. Bull., **1**, 90–102.

Kaeriyama, M., 1999. Hatchery programmes and stock management of Pacific salmon. In Stock Enhancement and Sea Ranching, ed. B.R. Howell, E. Moksness and T. Svasand. Blackwell, Oxford., 153–167.

Kaeriyama, M., Nakamura, M., Edpalina, R., Bower, J.R., Yamaguchi, H., Walker, R.V. and Myers, K.W., 2004. Change in feeding ecology and trophic dynamics of Pacific salmon (*Oncorhynchus* spp.) in the central Gulf of Alaska in relation to climate events. Fish. Oceanogr., **13**, 197–207.

Kaeriyama, M., Yatsu, A., Noto, M. and Saitoh, S., 2007. Spatial and temporal changes in the growth patterns and survival of Hokkaido chum salmon populations in 1970–2001. N. Pac. Anadr. Fish Comm. Bull., **4**, 251–256.

Kamezawa, Y., Azumaya, T., Nagasawa, T. and Kishi, M.J., 2007. A fish bioenergetics model of Japanese chum salmon (*Oncorhynchus keta*) for studying the influence of environmental factor changes. Bull. Japan. Soc. Fish. Oceanogr., **71**, 87–96.

Kawamiya, M., Yoshikawa, C., Kato, T., Sato, H., Suda, K., Watanabe, S. and Matsuno, T., 2005. Development of an integrated earth system model on the earth simulator. J. Earth Sim., **4**, 18–30.

Kishi, M.J., Kashiwai, M., Ware, D.M., Megrey, B.A., Eslinger, D.L., Werner, F.E. and 22 authors, 2007. NEMURO—A lower trophic level model for the North Pacific marine ecosystem. Ecol. Modell., **202**, 12–25.

Megrey, B.A., Rose, K.A., Ito, S., Hay, D.E., Werner, F.E., Yamanaka, Y. and Noguchi Aita, M., 2007a. North Pacific basin-scale differences in lower and higher trophic level marine ecosystem responses to climate impacts using a nutrient-phytoplankton–zooplankton model coupled to a fish bioenergetics model, Ecol. Modell., **202**, 196–210.

Megrey, B.A., Rose, K.A., Klumb, R.A., Hay, D.E., Werner, F.E., Eslinger, D.L. and Smith, L.S., 2007b. Abioenergetics-based population dynamics model of Pacific herring (*Clupea harengus pallasii*) coupled to a lower trophic level nutrient-phytoplankton-zooplankton model: description, calibration, and sensitivity analysis. Ecol. Modell., **202**, 144–164.

- Megrey, B.A., Rose, K.A., Werner, F.E., Klumb, R.A. and Hay, D., 2002. A generalized fish bioenergetics/biomass model with an application to Pacific herring. *PICES Sci. Rep.*, **20**, 4–12.
- Morita, S.H., Morita, K. and Sakano, H., 2001. Growth of chum salmon (*Oncorhynchus keta*) correlated with sea-surface salinity in the North Pacific. *ICES Journal of Marine Science*, **58** 1335–1339.
- Nagasawa, K., 2000. Winter zooplankton biomass in the subarctic North Pacific, with discussion on the overwintering survival strategy of Pacific salmon (*Oncorhynchus* spp.). *N. Pac. Anadr. Fish Comm. Bull.*, **2**, 21–32.
- Neave, F., Yonemori, Y. and Bakkala, R.G., 1980. Distribution and origin of chum salmon in offshore waters of the North Pacific Ocean. *INPFC. Bull.*, **35**, 79 pp.
- NPAFC Doc.769 Rev. Annual Report of the Bering-Aleutian Salmon International Survey(BASIS), 2003
- Rose, K.A., Werner, F.E., Megrey, B.A., Aita, M.N., Yamanaka, Y., Hay, D.E., Schweigert, J.F. and Foster, M.B., 2007. Simulated herring growth responses in the Northeastern Pacific to historic temperature and zooplankton conditions generated by the 3-dimensional NEMURO nutrient-phytoplankton-zooplankton model. *Ecol. Modell.*, **202**, 184–195.
- Rudstam, L.G., 1988. Exploring the dynamics of herring consumption in the Baltic: Applications of an energetic model of fish growth. *Kieler Meeresforsch., Sonderh.*, **6**, 312–322.
- Shuntov, V.P. and Dulepova, Y.P., 1996. Biota of the Okhotsk Sea: structure of communities, the interannual dynamics and current status. *PICES Sci. Rep.*, **6**, 263–271.
- Tadokoro, K., Ishida, Y., Davis, N.D., Ueyanagi, S. and Sugimoto, T., 1996. Change in chum salmon (*Oncorhynchus keta*) stomach contents associated with fluctuation of pink salmon (*O. gorbuscha*) abundance in the central subarctic Pacific and Bering Sea. *Fish. Oceanogr.*, **5**, 89–99.
- Thornton, K.W. and Lessem, A.S., 1978. A temperature algorithm for modifying biological rates. *Trans. Amer. Fish. Soc.*, **107**, 284–287.

Trudel, M., Geist D.R. and Welch D.W., 2004. Modeling the oxygen consumption rates in Pacific salmon and steelhead: An assessment of current models and practices. *Trans. Am. Fish. Soc.*, **133**,326–348.

Ueno, Y. and Ishida, Y., 1996. Summer distribution and migration routes of juvenile chum salmon (*Oncorhynchus keta*) originating from rivers in Japan. *Bull. Nat. Res. Far Fish.*, **33**, 139–147.

Urawa, S., 2000. Migration route and future works of Japanese salmon. In National salmon resources center news, No.5, 3–9. (in Japanese)

Ware, D.W., 1978. Bioenergetics of pelagic fish: theoretical change in swimming speed and ration with body size. *J. Fish. Res. Board Can.*, **34**,220–228.

Welch, D.W., Chigirinsky, A.I. and Ishida Y., 1995. Upper thermal limits on the oceanic distribution of Pacific salmon (*Oncorhynchus spp.*) in the spring. *Can. J. Fish. Aquat. Sci.*, **52**,489–503.

Welch, D.W., Ishida, Y. and Nagasawa, K., 1998. Thermal limits and ocean migrations of sockeye salmon (*Oncorhynchus nerka*): Long-term consequences of global warming. *Can. J. Fish. Aquat. Sci.*, **55**, 937–948.

Figure captions

Fig. 1. Annual change in growth at the first year of Ishikari River chum salmon in the Okhotsk Sea (Kaeriyama et al., 2007).

Fig. 2. Temporal changes in return rate of a) Hokkaido, Iwate, and Korean chum salmon populations and of b) local populations in Hokkaido.

Fig. 3. Schematic view of the three oceanic spatial domains and the numbers of life stages simulated in the model. The three oceanographic domains correspond to the western North Pacific (WNP), the Bering Sea (BS), and the eastern North Pacific (ENP).

Fig. 4. Simulated wet weight (line) and observed wet weight (squares) for a) the 1972 year class and b) the 1991 year class. c) The difference in wet weight between the 1972 and 1991 year classes for the control case (thin line), for the constant prey with time-dependent temperature case (bold line), and for the constant temperature with time-dependent prey case (dashed line).

Fig. 5. Time-dependent feature of body size in the Bering Sea in summer from 1971 to 1999. a) observed (based on data base of FRA, Japan) and b) simulated results.

Fig. 6. The impact of global warming on the ocean distribution of chum salmon based on the ICPP SRES-A1B scenario. Hatched portion represents the area where chum salmon can survive.

Fig. 7. The impact of global warming on the ocean distribution of chum salmon based on the ICPP SRES-A1B scenario near Hokkaido, Japan.

Table captions

Table 1. Life stages of the chum salmon bioenergetics model.

Table 2. Summary of parameter values used in the chum salmon bioenergetics model.




Fig.1




Fig.2




Fig.3




Fig.4




Fig.5




Fig.6




Fig.7

Stage	Ages(day)	Period	Region
1	404-586	12/1-5/31	W.North Pacific (WNP)
2	587-769	6/1-11/30	Bering Sea (BS)
3	770-951	12/1-5/31	E.North Pasific (ENP)
4	952-1134	6/1-11/30	Bering Sea (BS)
5	1135-1316	12/1-5/31	E.North Pasific (ENP)
6	1317-1499	6/1-11/30	Bering Sea (BS)
7	1500-1681	12/1-5/31	E.North Pasific (ENP)
8	1682-1834	6/1-10/31	Bering Sea (BS)

Table 1

Symbol	Parameter description	Value
Consumption, C_{max}		
ac	Intercept for C _{max} parameter	0.303
bc	Coefficient for C _{max} parameter versus weight	-0.275
xk1	Proportion of C _{max} for te1	0.1
xk2	Proportion of C _{max} for te2	0.98
xk3	Proportion of C _{max} for te3	0.98
xk4	Proportion of C _{max} for te4	0.5
te1	Temperature for xk1	3.0
te2	Temperature for xk2	3.9 ^a 5.0 ^b
te3	Temperature for xk3	5.0 ^c 10.0 ^b 8.0 ^d
te4	Temperature for xk4	10.4 ^a 12.0 ^b
K _i	Half saturation constant	0.55 ^e
Metabolism, R		
ars	Intercept for standard metabolic rate of 1-g fish at 0°C	0.0799
br	Coefficient standard metabolism versus body weight	0.8
cr	Coefficient standard metabolism versus temperature	0.069
ara	Intercept for swimming cost	eqn(11)
dr	Coefficient for swimming cost versus body weight	0.44
er	Coefficient for swimming cost versus swimming speed	2.42
Swimming Speed, U		
au	Intercept, for optimal foraging speed	20.3
bu	Coefficient versus weight	0.132
Egestion and Excretion, F, E and SDA		
af	Proportion of consumed food egested	0.16
ae	Proportion of consumed food excreted	0.1
ss	Coefficient for Specific Dynamic Action	0.175
Cal		
CALf	Caloric equivalent of fish	1323.68
CALz	Caloric equivalent of zooplankton	617.22
^a value for stage1,3,5,7 chum salmon ^b value for stage2,4,6,8 chum salmon ^c value for stage1 chum salmon ^d value for stage3,5,7 chum salmon ^e value for stage1-8 chum salmon		

Table 2