



Title	Kinetics of mixed venous CO <sub>2</sub> pressure in incremental-load exercise
Author(s)	Yano, Tokuo
Citation	Applied human science : journal of physiological anthropology, 16(1), 15-18
Issue Date	1997
Doc URL	<a href="https://hdl.handle.net/2115/44220">https://hdl.handle.net/2115/44220</a>
Rights	日本生理人類学会. 本文データは学協会の許諾に基づきCiNiiから複製したものである.
Type	journal article
File Information	yano_ahs16.pdf



## Kinetics of Mixed Venous CO<sub>2</sub> Pressure in Incremental-Load Exercise

Tokuo Yano

*Laboratory of Human Movement Science, Faculty of Education, Hokkaido University*

**Abstract.** The purpose of the present study was to examine the kinetics of mixed venous CO<sub>2</sub> pressure ( $\bar{P}\dot{V}CO_2$ ) in incremental-load exercise.  $\bar{P}\dot{V}CO_2$  seemed to indicate a linear increase after a somewhat slow increase. CO<sub>2</sub> store phase appearing in expiration of CO<sub>2</sub> (stored CO<sub>2</sub>) was calculated from the data of O<sub>2</sub> uptake ( $\dot{V}O_2$ ), CO<sub>2</sub> output ( $\dot{V}CO_2$ ) and work rate. The stored CO<sub>2</sub> indicated a linear increase after a time delay. The stored CO<sub>2</sub> also significantly related to  $\bar{P}\dot{V}CO_2$ . When kinetics of  $\bar{P}\dot{V}CO_2$  at light exercise intensity was supposed from stored CO<sub>2</sub>,  $\bar{P}\dot{V}CO_2$  seemed to indicate a linear increase with a time delay. Excessive expired CO<sub>2</sub> (excess CO<sub>2</sub>) was calculated from the data of  $\dot{V}O_2$  and work rate. The excess CO<sub>2</sub> significantly related to the increase of blood lactate. The blood lactate seemed to start to increase at around 1080 kpm/min. In spite of this change, a linear increase of  $\bar{P}\dot{V}CO_2$  was unchanged. This result was considered to be related to the excessive CO<sub>2</sub> expiration corresponding to the increase of blood lactate. Thus, within the present results and assumptions, it seemed that  $\bar{P}\dot{V}CO_2$  indicated a linear increase with a time delay without the effect of lactate increase in incremental-load exercise.

*(Appl Human Sci, 16(1): 15-18, 1997)*

**Keywords:** mixed venous CO<sub>2</sub> pressure, blood lactate, stored CO<sub>2</sub>, excess CO<sub>2</sub>

### Introduction

The kinetics of mixed venous CO<sub>2</sub> pressure ( $\bar{P}\dot{V}CO_2$ ) has been studied in human during exercise. Namely, when being indirectly measured in the constant-load exercise, it was reported that the  $\bar{P}\dot{V}CO_2$  indicated almost no change from the resting value for a few seconds at the onset of exercise, then increased up to the new steady state value (Edwards et al., 1972). Thereafter, when being directly measured using pulmonary artery catheterization, the  $\bar{P}\dot{V}CO_2$  was reported to indicate something of an increase even at a very early phase (Casaburi et al., 1989). Besides, Yano et al. (1995) pointed out that the later increase of  $\bar{P}\dot{V}CO_2$  overshoot when exercise intensity was high from the former reports

(Cerretelli et al., 1966; Davies et al., 1972; Yano, 1986), and examined whether this overshoot was related to early lactate produced at the onset of exercise. The result was that the effect could not be observed at least when increase of blood lactate was low.

In incremental-load exercise, the beginning phase of constant-load exercise continuously appears in response to increase of work load so that  $\bar{P}\dot{V}CO_2$  is considered to increase continuously during the exercise. It is also known that blood lactate starts to increase above a certain load in the incremental-load exercise (Beaver et al., 1986 a, b). Accordingly, in the incremental-load exercise, the effect of blood lactate may overlap with increase of  $\bar{P}\dot{V}CO_2$  due to increase of work load. Thus, the kinetics of  $\bar{P}\dot{V}CO_2$  was examined in the present study.

### Methods


The subjects were eight trained males belonging to sport clubs of the college. Their age was  $21 \pm 1.41$  yr (mean  $\pm$  SD), height  $170 \pm 4.58$  cm, weight  $59 \pm 4.48$  kg, and maximal oxygen uptake  $3.52 \pm 0.295$  l/min. All subjects were informed of the purpose of the experiment, the procedure, and the risks before their consent was obtained.

The exercise was performed using a Monark bicycle ergometer. The work load started from 180 kpm/min and thereafter was increased by 180 kpm/min every minute until the subject could not follow the pedaling rate of 60 rpm (exhaustion test). Two incremental-load exercises up to 540 and 1080 kpm/min were also performed later (submaximal tests).

At the exhaustion test, expired gas was collected for 5 min at rest, and every minute during the exercise with Douglas bags. Then fractions of O<sub>2</sub> and CO<sub>2</sub> in the Douglas bags were analyzed by a Mass spectrometer (Perkin Elma MGA-1100) and the volume was measured by a dry gas meter to give the minute ventilation. From these data, O<sub>2</sub> uptake ( $\dot{V}O_2$ ) and CO<sub>2</sub> output ( $\dot{V}CO_2$ ) were calculated. CO<sub>2</sub> rebreathing (Defares et al., 1961) was performed at rest and immediately after the exhaustion test, and immediately after 540 and 1080 kpm/min of submaximal tests. The rebreathing gas contained 5 % of CO<sub>2</sub> in O<sub>2</sub> balance. The CO<sub>2</sub> pressure obtained from the

CO<sub>2</sub> rebreathing was used to estimate mixed venous CO<sub>2</sub> pressure (detail in Yano et al., 1984). Blood was collected from the median cubital vein at rest and 5 min after the submaximal test up to 1080 kpm/min and exhaustion test, and was analyzed by Lactate Analyzer LA 640 (Roche).

The difference between  $\dot{V}O_2$  or  $\dot{V}CO_2$  in exercise and at rest ( $\Delta\dot{V}O_2$  or  $\Delta\dot{V}CO_2$ ) is shown in Fig. 1.  $\Delta\dot{V}CO_2$  indicated a linear increase at low work rate, and this increasing rate changed. Before this changing point, there was a definite difference between  $\Delta\dot{V}O_2$  and  $\Delta\dot{V}CO_2$  at each work rate. This difference was defined to be the respiratory phase of CO<sub>2</sub> store (stored $\dot{V}CO_2$ ). The regression line between  $\Delta\dot{V}CO_2$  and the work rate was obtained at the period of linear increase of  $\Delta\dot{V}CO_2$ . This regression line was extrapolated to obtain the prediction of  $\Delta\dot{V}CO_2$  after the changing point (pred $\dot{V}CO_2$ ). The difference between  $\Delta\dot{V}O_2$  and pred $\dot{V}CO_2$  was also defined as the stored $\dot{V}CO_2$  after the change of rate of increase of  $\Delta\dot{V}CO_2$ . Then stored $\dot{V}CO_2$  was added along different lapses. This was defined as the amount of CO<sub>2</sub> store (stored CO<sub>2</sub>). Furthermore, the difference between measured  $\Delta\dot{V}CO_2$  and pred $\dot{V}CO_2$  was defined as the respiratory phase of CO<sub>2</sub> excess (excess $\dot{V}CO_2$ ). The added value of excess $\dot{V}CO_2$  along different lapses was defined as the amount of excessive CO<sub>2</sub> expiration (excess CO<sub>2</sub>).



**Fig. 1** The differences between O<sub>2</sub> uptake and CO<sub>2</sub> output in exercise and at rest ( $\Delta\dot{V}O_2$  and  $\Delta\dot{V}CO_2$ ) are shown against work rate. Filled circles indicate  $\Delta\dot{V}O_2$ . Circles indicate  $\Delta\dot{V}CO_2$ .


The relationship between independent and dependent variables was expressed as the Pearson coefficient of correlation. A significant level was accepted at  $p < 0.05$ .

## Results

The excess $\dot{V}CO_2$  started from 1080 kpm/min for 3 subjects, 1260 kpm/min for 4 subjects, and 1440 kpm/min for 1 subject. Blood lactate was  $1.11 \pm 0.16$  mM at rest,  $1.79 \pm 0.66$  mM after 1080 kpm/min of submaximal test, and  $9.45 \pm 1.96$  mM after exhaustion test. The difference between blood lactates after exercise and at rest was significantly related to excessCO<sub>2</sub> ( $r = 0.828$ ,  $p < 0.001$ ).

Figure 2 shows the kinetics of stored CO<sub>2</sub> against the rise of work rate. A slope of the linear regression lines obtained between  $\Delta\dot{V}CO_2$  and work rate was not always the same as increasing rate of  $\Delta\dot{V}O_2$ . When the slope was steep, stored $\dot{V}CO_2$  became small at upper work rate and vice versa. Therefore, estimates were not always excellent, resulting in scatter estimates at high work rate. The stored CO<sub>2</sub> linearly increased against after the delay although stored CO<sub>2</sub> at 720 kpm/min indicated somewhat high value. This delay corresponded to about one minute.

Figure 3 shows the kinetics of  $P\bar{v}CO_2$  against the rise of work rate.  $P\bar{v}CO_2$  seemed to indicate a linear increase after a somewhat slow rise at the beginning phase of the incremental-load exercise. The rising rate of  $P\bar{v}CO_2$  did not change at the starting point of excess $\dot{V}CO_2$ . The  $P\bar{v}CO_2$  at exhaustion did not deviate from the straight line obtained between 540 and 1080 kpm/min although a lot of



**Fig. 2** Kinetics of stored CO<sub>2</sub> during incremental-load exercise. (mean  $\pm$  SD)

blood lactate was produced at exhaustion.

When the linear portion of  $\bar{P}\dot{V}CO_2$  rise was extrapolated to the lower work rate,  $\bar{P}\dot{V}CO_2$  became the value at rest when the work rate was 180 kpm/min corresponding to one minute delay. As this trend of  $\bar{P}\dot{V}CO_2$  kinetics was similar to that of stored  $CO_2$ , this relationship was examined. As shown in Fig. 4,  $\bar{P}\dot{V}CO_2$  significantly related to stored  $CO_2$  ( $r=0.876$ ,  $p<0.01$ ).

## Discussion

The purpose of the present study was to examine the kinetics of  $\bar{P}\dot{V}CO_2$  in incremental-load exercise.  $\bar{P}\dot{V}CO_2$  seemed to indicate a linear increase after a somewhat slow increase. Stored  $CO_2$  was calculated from the  $\Delta\dot{V}O_2$  and  $\Delta\dot{V}CO_2$ . The stored  $CO_2$  indicated linear increase after a time delay.  $\bar{P}\dot{V}CO_2$  significantly related to the stored  $CO_2$ . Blood lactate seemed to start to increase at around




Fig. 3 Kinetics of mixed venous  $CO_2$  pressure ( $\bar{P}\dot{V}CO_2$ ) during incremental-load exercise and at rest. (mean  $\pm$  SD)




Fig. 4 Relationship between mixed venous  $CO_2$  pressure ( $\bar{P}\dot{V}CO_2$ ) and stored  $CO_2$ .

1080 kpm/min. In spite of this changes, the linear increase of  $\bar{P}\dot{V}CO_2$  was unchanged. Blood lactate rise significantly related to excess  $CO_2$ .

In the present study, stored  $CO_2$  was obtained through the difference between  $\Delta\dot{V}O_2$  and  $\Delta\dot{V}CO_2$ . This is because if the absolute values of  $\dot{V}O_2$  and  $\dot{V}CO_2$  were to be used for the calculation of stored  $\dot{V}CO_2$ , the stored  $CO_2$  would be calculated at rest. In order to avoid this contradiction, the difference between  $\dot{V}O_2$  and  $\dot{V}CO_2$  at rest ( $\dot{V}O_{2rest} - \dot{V}CO_{2rest}$ ) is eliminated from  $\dot{V}O_2$  during exercise. Its value is, then, assumed to be equal to the  $CO_2$  product ( $\dot{V}CO_{2prod}$ ). That is,

$$\dot{V}CO_{2prod} = \dot{V}O_2 - (\dot{V}O_{2rest} - \dot{V}CO_{2rest})$$

$$\text{As stored } \dot{V}CO_2 = \dot{V}CO_{2prod} - \dot{V}CO_2,$$

$$\begin{aligned} \text{stored } \dot{V}CO_2 &= \dot{V}O_2 - (\dot{V}O_{2rest} - \dot{V}CO_{2rest}) - \dot{V}CO_2 \\ &= \Delta\dot{V}O_2 - \Delta\dot{V}CO_2 \end{aligned}$$

This is the present way of calculation on stored  $\dot{V}CO_2$ .

As respiratory quotient (RQ) is  $\dot{V}CO_{2prod}/\dot{V}O_2$ ,

$$\begin{aligned} RQ &= (\dot{V}O_2 - (\dot{V}O_{2rest} - \dot{V}CO_{2rest}))/\dot{V}O_2 \\ &= 1 - (\dot{V}O_{2rest} - \dot{V}CO_{2rest})/\dot{V}O_2 \end{aligned}$$

This equation becomes  $\dot{V}CO_{2rest}/\dot{V}O_{2rest}$  at rest as  $\dot{V}O_2$  is equal to  $\dot{V}O_{2rest}$ . This is corresponding to respiratory gas exchange ratio (RER) at rest. As  $\dot{V}O_2$  increases in exercise, the second term of the equation approaches zero, and RQ approaches from RER (=RQ) at rest to 1.

Thus the stored  $CO_2$  was obtained under these conditions.

The obtained stored  $CO_2$  related to  $\bar{P}\dot{V}CO_2$ . Therefore, the kinetics of  $\bar{P}\dot{V}CO_2$  at light and moderate exercise intensities can be estimated from the stored  $CO_2$ . The estimated kinetics of  $\bar{P}\dot{V}CO_2$  seems to be a linear increase with a time delay.

In the incremental exercise, it is known that blood lactate starts to increase above a certain work load (Beaver et al., 1986 a, b). This lactic acid is partly buffered by the bicarbonate system (Hultman and Sahlin, 1980; Yano, 1987). In this chemical reaction, the bicarbonate ion shifts toward  $CO_2$ . If the shifted  $CO_2$  is not perfectly removed,  $CO_2$  pressure increases. This was predicted particularly in the active muscle producing lactic acid, and the increase of  $CO_2$  pressure in the active muscle could induce the rise of mixed venous  $CO_2$  pressure. However, the effect of lactate increase on  $\bar{P}\dot{V}CO_2$  was not confirmed in the present study.

The reason for  $\bar{P}\dot{V}CO_2$  being unaffected may relate to the velocity of chemical reaction between  $CO_2$  and bicarbonate ion in bicarbonate buffering system. This velocity is slow in the muscle cell (Sahlin et al., 1978) as the enzyme to catalyze the reaction is inactive. Therefore, the buffer by bicarbonate system does not produce the quick shift toward  $CO_2$ . If during this delay the bicarbonate ion is removed from muscle cell,  $CO_2$  in the muscle cell is not increased.

The present study indicated that  $CO_2$  was excessively expired in response to the increase of blood lactate.

Unless the obtained excessCO<sub>2</sub> included an amount of the bicarbonate ion removed from active muscle cell, CO<sub>2</sub> pressure in the active muscle cell, consequently  $\bar{P}\dot{V}\text{CO}_2$  could have been affected.

Thus, within the present results and assumptions it seemed that  $\bar{P}\dot{V}\text{CO}_2$  indicated a linear increase with a time delay without the effect of lactate increase in incremental-load exercise.

## References

- Beaver WL, Wasserman K, Whipp BJ (1986a) Bicarbonate buffering of lactic acid generated during exercise. *J Appl Physiol* 60: 472-478
- Beaver WL, Wasserman K, Whipp BJ (1986b) A new method for detecting anaerobic threshold by gas exchange. *J Appl Physiol* 60: 2020-2027
- Casaburi R, Daly J, Hansen JE, Effros RM (1989) Abrupt changes in mixed venous blood gas composition after the onset of exercise. *J Appl Physiol* 67: 1106-1112
- Cerretelli P, Sikand R, Farhi LE (1966) Readjustments in cardiac output and gas exchange during onset of exercise and recovery. *J Appl Physiol* 21: 1345-1350
- Davies CTM, DiPrampo PE, Cerretelli P (1972) Kinetics of cardiac output and respiratory gas exchange during exercise and recovery. *J Appl Physiol* 32: 618-625
- Defares JM, Weise ME, Duyff JW (1961) New indirect Fick procedure for the determination of cardiac output. *Nature* 192: 760-761
- Edwards RHT, Denison DM, Jones G, Davies CTM, Campbell EJM (1972) Changes in mixed venous gas tensions at start of exercise in man. *J Appl Physiol* 32: 165-169
- Hultman E, Sahlin K (1980) Acid-base balance during exercise. *Sport Sci Rev* 8: 41-128
- Sahlin K, Alvestrand A, Brandt R, Hultman E (1978) Intracellular pH and bicarbonate concentration in humans muscle during recovery from exercise. *J Appl Physiol* 45: 474-480
- Yano T (1986) Immediate CO<sub>2</sub> storage capacity at the onset of exercise. *Jpn J Physiol* 36: 1241-1251
- Yano T (1987) A theoretical approach to excess CO<sub>2</sub> expiration due to lactate production in exercise. *Jpn J Physiol* 37: 937-940
- Yano T, Kumazaki Y, Asano K (1995) Kinetics of mixed venous CO<sub>2</sub> pressure in constant-load exercise. *Appl Human Sci* 14: 155-156
- Yano T, Norsk P, Bond-Petersen F (1984) Cardiac output determined by the CO<sub>2</sub> rebreathing method with correction of lung-bag volume shrinkage. *Jpn J Physiol* 34: 883-891

---

Received: April 1, 1996

Accepted: October 15, 1996

Correspondence to: Tokuo Yano, Laboratory of Human Movement Science, Faculty of Education, Hokkaido University, Kita 12 Nishi 7, Kita-ku, Sapporo 001, Japan