



Title	Involvement of AtNAP1 in the regulation of chlorophyll degradation in <i>Arabidopsis thaliana</i>
Author(s)	Nagane, Tomohiro; Tanaka, Ayumi; Tanaka, Ryouichi
Citation	Planta, 231(4), 939-949 <a href="https://doi.org/10.1007/s00425-010-1099-8">https://doi.org/10.1007/s00425-010-1099-8</a>
Issue Date	2010-03
Doc URL	<a href="https://hdl.handle.net/2115/44896">https://hdl.handle.net/2115/44896</a>
Rights	The original publication is available at <a href="http://www.springerlink.com">www.springerlink.com</a>
Type	journal article
File Information	SupplementalMaterial-2.pdf, Supplemental Material 2



Involvement of AtNAP1 in the regulation of chlorophyll degradation in *Arabidopsis thaliana*

Planta

Tomohiro Nagane, Ayumi Tanaka and Ryouichi Tanaka

Institute of Low Temperature Science, Hokkaido University, N19W8, Kita-ku, Sapporo 060-0819, Japan

e-mail: Correspondence email: rtanaka@lowtem.hokudai.ac.jp; Tel: +81 11 706 5496, Fax: +81 11 706 5493

## Online Resource 2 Pigment contents before and after dark treatment

	Chlorophyll a (nmol/gFW)	Chlorophyll b (nmol/gFW)	HMchl (nmol/gFW)	Pheophorbide a (nmol/gFW)	Total chlorophyll (nmol/gFW)	Chlorophyll a/b ratio
Before dark treatment						
Wild type	2021±367	652±111	1.17±0.33	17.5±9.8	2672±478	3.1±0.05
<i>hmc1</i>	1186±154	357±38	25.9±4.5	3.2±5.8	1543±191	3.31±0.12
<i>pao1</i>	1476±218	457±71	1.44±0.63	2.8±4.3	1933±289	3.23±0.07
After dark treatment						
Wild type	855±196	224±54	5.59±2.47	50±18.2	1079±239	3.9±0.67
<i>hmc1</i>	609±214	204±39	20.2±5.77	342±192	813±252	2.9±0.45
<i>pao1</i>	896±208	311±82	5.39±1.19	768±158	1208±290	2.9±0.1

The levels of chlorophyll and intermediate molecules in the wild type, *hmc1* and *pao1* plants. Plants were grown for 3 weeks under continuous illumination. The third whorl of the leaves were harvested, and then the remaining plants were transferred to darkness and kept for 5 days. The same whorl of the leaves were harvested. The harvested leaves were grinded in acetone and the pigments were extracted and subjected to HPLC analysis. SD of five independent samples from a single harvest is shown.