



HOKKAIDO UNIVERSITY

Title	Sperm dimorphism in terms of nuclear shape and microtubule accumulation in <i>Cyrtanthus mackenii</i>
Author(s)	Hirano, Tomonari; Hoshino, Yoichiro
Citation	Sexual Plant Reproduction, 23(2), 153-162 https://doi.org/10.1007/s00497-009-0123-2
Issue Date	2010-03
Doc URL	https://hdl.handle.net/2115/44961
Rights	The original publication is available at www.springerlink.com
Type	journal article
File Information	497_2009_123_MOESM2_ESM.pdf, Supplemental Table 1



Supplemental Table 1 Distances between vegetative nucleus and sperm nucleus and between sperm nuclei.

Pollen culture periods (h)	Vegetative nucleus	
	to Svn (μm)	Svn-Sua (μm)
12	$2.1 \pm 1.8^{\text{ns}}$	$14.0 \pm 4.0^{\text{a}}$
24	$1.8 \pm 0.7^{\text{ns}}$	$3.2 \pm 1.1^{\text{b}}$
36	$2.1 \pm 1.6^{\text{ns}}$	$4.0 \pm 1.6^{\text{b}}$

Data represent mean $\pm$ standard deviations of 15 MGUs. In each column, values indicated with the same superscript letter are not significantly different at the 0.05 level. NS: not significant.