



Title	Stabilizing effect of Ca <sup>2+</sup> on myosin and myofibrils of squid mantle muscle as affected by heating conditions
Author(s)	Ohno, Takaaki; Kinoshita, Yasunori; Konno, Kunihiko
Citation	Fisheries Science, 77(3), 425-430 <a href="https://doi.org/10.1007/s12562-011-0347-6">https://doi.org/10.1007/s12562-011-0347-6</a>
Issue Date	2011-05
Doc URL	<a href="https://hdl.handle.net/2115/45422">https://hdl.handle.net/2115/45422</a>
Rights	© 2011 公益社団法人日本水産学会
Type	journal article
File Information	FS77-3_425-430.pdf



**Stabilizing effect of Ca<sup>2+</sup> on myosin and myofibrils of squid mantle muscle as affected by heating conditions.**

Takaaki Ohno<sup>\*1</sup> · arunori Kinoshita<sup>\*1,\*2</sup> · unihiko Konno<sup>\*1,\*3</sup>

<sup>\*1</sup> *Faculty of Fisheries Sciences, Hokkaido University, Hakodate, Hokkaido 041-8611, Japan*

<sup>\*2</sup> *Hokkaido Industrial Technology Center, Hakodate, Hokkaido 041-0801, Japan*

<sup>\*3</sup> Corresponding author

E-mail: konno@fish.hokudai.ac.jp

**Abstract** Suppressive effect of  $\text{Ca}^{2+}$  on the thermal denaturation of myosin and on myofibrils of squid mantle muscle was compared. Its stabilization effect on myosin was smaller than that on myofibrils, and was not affected by KCl concentration. The stabilizing effect of  $\text{Ca}^{2+}$  on myosin decreased with lowering the temperature for heating showing no stabilization at 20 °C, while the effect on myofibrils was evenly detected at all temperatures tested. Stabilizing effect of  $\text{Ca}^{2+}$  on myosin disappeared even at 30 °C in the presence of sorbitol showing a small inactivation rate, while the effect of  $\text{Ca}^{2+}$  was equally detected on myofibrils irrespective of the reduction of inactivation rate with sorbitol. Stabilization of myosin by  $\text{Ca}^{2+}$  again appeared even in the presence of sorbitol by raising the heating temperature to 38 °C. It was suggested that  $\text{Ca}^{2+}$  confers stabilization on myosin only when myosin is in under unstable conditions. Stabilization effect of  $\text{Ca}^{2+}$  on myosin was enhanced upon F-actin binding, namely  $\text{Ca}^{2+}$  bound myosin was more significant stabilized by F-actin binding, and the effect was no longer affected by the conditions for heating.

**Keywords** squid · myosin · denaturation · sorbitol ·  $\text{Ca}^{2+}$

## Introduction

Thermal stability of fish myosin is deeply dependent on the environmental temperature for habitation. Consequently, the stability of myosin from cold water fish is much more labile than that from warm water fish species when compared under the identical conditions [1]. This was true for squid species [2, 3]. Squid of cold water or deep sea contained unstable myosin than one of warm water. We have reported unique properties in the thermal denaturation process of squid myofibrils and myosin subfragment-1 (S-1) that myosin in squid mantle muscle myofibrils was significantly stabilized upon addition of  $\text{Ca}^{2+}$  to the heating medium. The stabilization by  $\text{Ca}^{2+}$  was much more remarkable with myofibrils, myosin bound to F-actin, than with myosin S-1. Stabilization by  $\text{Ca}^{2+}$  with myofibrils at around 0.1 M KCl was around 100 times, while one with S-1 was as small as 5 times [4]. We studied the stabilization by  $\text{Ca}^{2+}$  by using various species of squid and cuttlefish, and concluded that the same stabilizing effect was generally observable with all species examined [3].

It is well established with fish myofibrils that high concentration of salt such as 1.5-2 M NaCl or KCl resulted in the selective denaturation of F-actin [5], and consequently myosin loses the protection by F-actin there. With squid myofibrils, salt concentration required for a complete loss of protection by F-actin occurred at about 0.75 - 0.8 M which was roughly a half concentration needed for fish myofibrils. The magnitude of the accelerated inactivation rate caused by increasing

KCl concentration or as a result of loss of protection was about 300 - 500 times, which was much greater than that obtained with fish myofibrils [2, 5]. In the present paper, we further studied the stabilizing effect of  $\text{Ca}^{2+}$  on myosin alone and on myofibrils by changing heating conditions considering the thermal inactivation rate itself to understand what factors are involved in the stabilizing action by  $\text{Ca}^{2+}$  on myosin, and how stabilizing effect of  $\text{Ca}^{2+}$  and one by F-actin binding are related to understand the significant stabilizing effect of  $\text{Ca}^{2+}$  on myofibrils.

## **Materials and Methods**

Myofibrils were prepared from the mantle muscle of Japanese common squid *Todarodes pacificus* as described earlier [2]. Myofibrils were suspended in 0.1 M KCl, 20 mM Tris-HCl (pH 7.5). Myosin was prepared from the myofibrils by applying the ammonium sulfate fractionation in the presence of 2 mM  $\text{MgCl}_2$ -ATP, the identical procedures for isolation of myosin from carp myofibrils as has been reported [6]. In the present study, myofibrils and myosin suspended in 0.1 M KCl, 20 mM Tris-HCl (pH 7.5) were used. For studying the stabilizing effect of  $\text{Ca}^{2+}$  on myosin and myofibrils, heating was conducted in the presence of 1 mM either  $\text{CaCl}_2$  (Ca-medium) or EDTA (EDTA-medium). Free  $\text{Ca}^{2+}$  concentrations in the heating medium were set by using Ca-EGTA buffer using fixed concentration of 0.1 mM  $\text{CaCl}_2$  and varied concentrations of EGTA. EGTA concentration required

was calculated by assuming the binding constant of  $1 \times 10^6$  at pH 7.5 [7]. Heating temperature was optionally changed dependent on the purposes. In some experiments, the heating medium contained sorbitol to increase the stability or to reduce the inactivation rate at the temperature [8]. Thermal denaturation of myosin was monitored by following the  $\text{Ca}^{2+}$ -ATPase inactivation.  $\text{Ca}^{2+}$ -ATPase was assayed in a medium of 0.5 M KCl, 25 mM Tris-maleate (pH 7.0), 5 mM  $\text{CaCl}_2$ , and 1 mM ATP at 25°C. The inactivation rate ( $\text{sec}^{-1}$ ) was calculated by assuming the inactivation as the first order reaction kinetics.

## Results

$\text{Ca}^{2+}$ -ATPase inactivation rates of squid myosin both in two media containing either  $\text{Ca}^{2+}$  or EDTA was compared with those for myofibrils by changing KCl concentrations (Fig. 1a). To estimate the inactivation rate of myosin accurately, the heating temperature was set to 25°C. The inactivation rate for myosin was slightly affected by KCl concentration both in Ca- and EDTA-media and the inactivation rate in EDTA-medium was always greater than that in Ca-medium. The stabilization by  $\text{Ca}^{2+}$  estimated from the rates was about 5 times. The inactivation rates of myofibrils in Ca- and EDTA-media above 0.8 M KCl were the same as those of myosin in respective media. It was confirmed that squid myosin lost the protection by F-actin at such high concentrations of KCl. The

rates for myofibrils below 0.4 M (Ca-medium) and 0.2 M KCl (EDTA-medium) were too small to be measured at 25 °C. Stabilizing extent by Ca at low concentration of KCl on myofibrils was shown in Fig. 1b, in which the heating temperature was raised to 35 °C. The result showed that strengthened F-actin binding to myosin by lowering KCl concentrations further stabilized myosin in the presence of Ca<sup>2+</sup>. An apparent stabilization by Ca<sup>2+</sup> at 0.05 M KCl was about 50 times, which was 10 times greater than that obtained with myosin. Moreover, the changes in the rates by increasing KCl concentration from 0.05 to 0.8 M were about 1000 times and about 40 times in Ca- and EDTA-media, respectively. The differences in the rates between two media with myofibrils decreased gradually with increasing KCl concentration due to a gradual decrease in the affinity between myosin and F-actin.

We studied whether the stabilization by Ca<sup>2+</sup> on myofibrils and myosin is affected by heating temperature. Thermal inactivation rates of myosin and myofibrils at 0.1 M KCl in Ca- and EDTA-media were estimated by varying the heating temperature and the rates were analyzed by using Arrhenius plot (Fig. 2). Myofibrils gave straight lines with the identical slopes both in Ca- and EDTA-media indicating practically the same stabilizing effect of Ca<sup>2+</sup> (roughly 80 times) at all temperatures studied. The estimated activation energies from the slopes were about 110 kcal/mol/degree both in Ca- and EDTA-media. However, myosin showed a complex temperature dependency giving a breaking point at around 20 °C in Ca- and EDTA-media. At high temperature

range, the slope for EDTA-medium was steeper than that for Ca-medium. Consequently, the difference in the rates between Ca- and in EDTA-media became smaller and smaller with lowering the heating temperature. Below the breaking point, the slope was more gradual than that at high temperature range. Moreover, no stabilizing effect of  $\text{Ca}^{2+}$  was observed any more at lower temperature range. The activation energies obtained with myosin at higher temperature range in Ca- and EDTA-medium were 80 and 100 kcal/mol/degree, respectively and one at lower temperature zone was 45 kcal/mol/degree, which was roughly a half that of high temperature range. Inactivation rate of myosin at 20 °C where no stabilizing effect of  $\text{Ca}^{2+}$  on myosin was detected was  $5 \times 10^{-5} \text{ sec}^{-1}$ , while the stabilizing effect by  $\text{Ca}^{2+}$  was still detectable with myofibrils with the same inactivation rates in Ca-medium. The inactivation rates of myofibrils below 30 °C were too small to measure, and the stabilizing effect of  $\text{Ca}^{2+}$  on myofibrils at such low temperatures could not be studied.

As the stabilizing effect of  $\text{Ca}^{2+}$  on myosin and on myofibrils was distinguished, we measured  $\text{Ca}^{2+}$  concentration requirement for the stabilization for both samples. Free  $\text{Ca}^{2+}$  concentration was set by using Ca-EGTA buffer system. Relative stabilization by  $\text{Ca}^{2+}$  obtained with myosin and with myofibrils was calculated by assuming the stabilization at 1 mM  $\text{CaCl}_2$  and EDTA as 100 % and 0 %, respectively (Fig. 3). Myosin was fully stabilized above 10  $\mu\text{M}$  with a half maximal stabilization at 12.6 nM. The concentration was 240 times lower than that required for the Ca-regulation of Mg-ATPase of squid myosin in which a half activation of Mg-ATPase was observed

at 3  $\mu\text{M}$   $\text{Ca}^{2+}$  [9]. Thus, the  $\text{Ca}^{2+}$  for regulation and for stabilization was clearly distinguished each other. Myofibrils were fully stabilized above 10  $\mu\text{M}$  with half stabilization at 200 nM. The concentration required for achieving the stabilization for myofibrils was higher than that for myosin by about 16 times. The result indicated that the increased extent of stabilization by  $\text{Ca}^{2+}$  upon F-actin binding was accompanied by the lowered sensitivity to  $\text{Ca}^{2+}$ .

In Fig. 2, we found that the stabilization by  $\text{Ca}^{2+}$  disappeared at low temperatures where the inactivation rates were very small. We wondered whether stabilization by  $\text{Ca}^{2+}$  is determined by the inactivation rates themselves or heating temperatures. We tried to answer the above questions by changing inactivation rates of myosin and myofibrils artificially. Our first attempt was to study the effect of sorbitol addition to the heating medium so as to reduce the inactivation rate at the heating temperature. Heating temperatures were chosen to give similar inactivation rates for myosin and for myofibrils, 30 °C for myosin and 40 °C for myofibrils, respectively. The inactivation rates in the presence of various concentrations of sorbitol were estimated in Ca- and EDTA-media and their logarithmic values were plotted against sorbitol concentrations according to Ooizumi *et al.* [8]. As Ooizumi *et al.* have reported with fish myofibrils, the plotting gave straight lines for all cases (Fig. 4a, b). The inactivation rates for myofibrils both in Ca- and EDTA- media decreased similarly with increasing sorbitol concentration, and the same stabilization by  $\text{Ca}^{2+}$  was observed at any concentrations of sorbitol or with any inactivation rates (Fig. 4a). For example, inclusion of 1.2 M

sorbitol decreased the rate to 1/60 that without sugar, while the same stabilization by  $\text{Ca}^{2+}$  was observed. Squid myosin was also stabilized upon addition of sorbitol in a concentration dependent manner both in Ca- and EDTA-media (Fig. 4b). However, slopes for Ca- and EDTA media were clearly different from each other. The slope with  $\text{Ca}^{2+}$  (1.29/mol of sorbitol) was a little smaller than that obtained with EDTA (1.67/mol sorbitol). Consequently, the difference in the rates in Ca- and EDTA-media became smaller and smaller with increasing the concentration, and finally the rates in two media was identical at 1.8 M sorbitol. We noticed that the inactivation rate in the presence of 1.8 M sorbitol with  $\text{Ca}^{2+}$  and with EDTA was about 370 and 2170 times smaller than those without sorbitol, respectively. The rates at 30 °C with 1.8 M sorbitol ( $1.42 \times 10^{-5} \text{ sec}^{-1}$ ) corresponded to the rates at 15 °C in the absence of sorbitol in Fig. 2. As the myofibrils with a similar inactivation rate upon addition of sorbitol still showed the stabilization effect of  $\text{Ca}^{2+}$ , the stabilized effect of  $\text{Ca}^{2+}$  was determined by the inactivation itself not by the heating temperature.

Inclusion of 1.8 M sorbitol made it possible to estimate the inactivation rates of myosin at higher temperatures, a similar temperature range for heating of myofibrils without sorbitol. Inactivation rates of myosin in Ca- and EDTA-media with 1.8 M sorbitol were estimated at various temperatures and the rates were analyzed by using Arrhenius plot (Fig. 5). The profiles without sorbitol were taken from Fig. 2 for a comparison. Stabilizing effect of  $\text{Ca}^{2+}$  on myosin appeared again even in the presence of sorbitol when the temperature was raised to 38 °C. Heating of myosin

with sorbitol at 38 °C gave similar rates to ones without sorbitol at 30 °C in respective two media. The stabilization by  $\text{Ca}^{2+}$  obtained was 4.4 times under the conditions. A gradual decrease of stabilizing effect by  $\text{Ca}^{2+}$  with lowering the heating temperature was reproduced again. The inactivation rates with  $\text{Ca}^{2+}$  and with EDTA became the same below 31 °C in the presence of 1.8 M sorbitol. The rates at 31 °C with sorbitol were rather similar to those without sorbitol at 20 °C. A general feature of temperature dependent inactivation rates for myosin with sorbitol was similar to that without sorbitol, although the slopes were different from each other. The slopes for myosin with sorbitol were steeper than those for ones without sorbitol. The addition of sorbitol changed the activation energies for myosin denaturation both in Ca- and EDTA-media. These were 107 and 144 kcal/mol, respectively, which were larger than those without sorbitol (81 and 104 kcal/mol for Ca- and EDTA-media). At 30 °C, the inactivation rates of myofibrils were larger than those of myosin with sorbitol (see Fig. 2). Appearance of the stabilizing effect of  $\text{Ca}^{2+}$  on myosin in the presence of sorbitol at high temperature and its loss at relatively low temperature indicated that the stabilizing effect of  $\text{Ca}^{2+}$  was not determined by the temperature for heating but by the denaturation rate itself.

## **Discussion**

As the KCl concentration for heating of myosin alone did not affect the stabilization by  $\text{Ca}^{2+}$  (about 5 times), a significantly large stabilization by  $\text{Ca}^{2+}$  observed with myofibrils (roughly

100 times) at low-salt medium was a consequence of binding with F-actin. Since F-actin binding stabilizes myosin significantly, it is certain that F-actin changes the conformation of myosin so as to be resistant to heating. We confirmed that apparent stabilization upon F-actin binding in the presence of  $\text{Ca}^{2+}$  was more than 500 times, while the one in the presence of EDTA was only about 40 times (Fig. 1b). The fact indicated that the stabilization by  $\text{Ca}^{2+}$  and by F-actin binding is not a simple sum. When we assume the stabilization extent by F-actin in the presence of EDTA is the magnitude of stabilization truly achieved by F-actin binding, the increased stabilization by F-actin was achieved with  $\text{Ca}^{2+}$  bound myosin. Protective effect of F-actin on myosin was negligible above 0.8 M KCl and small protection was found at around 0.4-0.5 M, while the apparent stabilization effect by  $\text{Ca}^{2+}$  for the sample was nearly the same as found with myosin alone (Fig. 1a). Thus the two stabilizing effects were clearly distinguished.

F-actin bound myosin, myofibrils, was different from myosin in several points in terms of stabilization by  $\text{Ca}^{2+}$ . Lowering temperature lost the stabilization by  $\text{Ca}^{2+}$  for myosin, while such change in temperature unaffected the effect by  $\text{Ca}^{2+}$  with myofibrils. Decreased inactivation rate rather than heating temperature itself was the reason for the loss of stabilization there because the inclusion of sorbitol so as to reduce the inactivation rate diminished the stabilization by  $\text{Ca}^{2+}$  at the same temperature. The conclusion was confirmed by the fact that the stabilization by  $\text{Ca}^{2+}$  appeared again even with sorbitol by raising the temperature so as to give large inactivation rate. Under these

conditions, stabilization by  $\text{Ca}^{2+}$  was equally detectable with myofibrils. These results indicated that structural change caused by  $\text{Ca}^{2+}$  binding to myosin is detectable only when myosin is placed under unstable conditions. The change might not be large because the effect was easily lost by shifting the conditions. However, myofibrils in the presence of sorbitol showing very small inactivation rates still showed the same stabilization by  $\text{Ca}^{2+}$  as observed without sorbitol. The stable structure of  $\text{Ca}^{2+}$  bound myosin seemed sensed by F-actin binding. Stabilizing effect of  $\text{Ca}^{2+}$  on myosin and on myofibrils was summarized in Table 1. With myosin, stabilization was observed only when the inactivation rates were  $1-3 \times 10^{-3} \text{ (sec}^{-1}\text{)}$ , and never be observed with the inactivation rate of  $1-5 \times 10^{-5} \text{ (sec}^{-1}\text{)}$ , 100 times smaller. However, stabilizing effect on myofibrils was equally observed irrespective of the inactivation rates. The rate with  $\text{Ca}^{2+}$  at  $40^\circ\text{C}$  in the presence of 1.3 M sorbitol was as low as  $1-5 \times 10^{-5} \text{ (sec}^{-1}\text{)}$ . Nevertheless, the stabilizing effect was the same.

Moreover, the effect of  $\text{Ca}^{2+}$  on myosin itself and on F-actin bound myosin was also distinguished by  $\text{Ca}^{2+}$  requirement for the stabilization (Fig. 3). Although the stabilization extent by  $\text{Ca}^{2+}$  was greater with F-actin bound myosin, myofibrils required higher  $\text{Ca}^{2+}$  concentration for a full stabilization than myosin itself. As the  $\text{Ca}^{2+}$  concentration for stabilization was clearly lower than that for Ca-regulation,  $\text{Ca}^{2+}$  binding to regulatory light chain is not involved in the mechanism (Fig. 3). Different concentration requirement with myofibrils (F-actin-bound myosin) from myosin also suggested different mechanism from Ca-regulation because  $\text{Ca}^{2+}$  requirements for Mg-ATPase

activation with myosin and actin-myosin complex are the same [9]. At present we have no information on the  $\text{Ca}^{2+}$  binding region on myosin for stabilization.

As the  $\text{Ca}^{2+}$  concentration required for the stabilization was remarkably lower than that for Ca-regulation, myosin in muscle is always kept stable irrespective of contraction and relaxation. After the death of squid, ATP consumption leads to a rigor contraction resulted in a strong binding of F-actin to myosin [10]. The complex formation makes squid myosin very stable during the storage of squid muscle. Indeed squid myosin itself is not so stable, but two significant stabilizing factors,  $\text{Ca}^{2+}$  and F-actin, make squid myosin uniquely stable. The properties of squid myosin contribute the long life of squid muscle without loss of quality during the frozen storage.

## References

1. Hashimoto, A., Kobayashi, A., Arai, K. (1982) Thermostability of fish myofibrillar Ca-ATPase and adaptation to environmental temperature. *Nippon Suisan Gakkaishi* 48:671-684 (in Japanese with English Abstract)
2. Konno, K. (1991) Thermal denaturation of squid myofibrils: Effects of calcium ion and EDTA. *Nippon Suisan Gakkaishi* 57:2145-2149
3. Konno, K., Yuasa, M. (1993) Comparative aspect of thermal stability of squid myofibrils. *Comp Biochem Physiol* 105A: 427-430
4. Konno, K. (1991) Complex formation between regulatory and essential light chain on squid mantle myosin subfragment-1 revealed by thermal denaturation method. *J Biochem* 109:816-821
5. Wakameda, A. Arai, K. (1986) Dissociation of carp myosin B into actin and myosin in the presence of a high concentration of NaCl. *Nippon Suisan Gakkaishi* 52:293-300 (in Japanese with English Abstract)

6. Koseki, H., Kato, S., Konno, K. (1993) Myosin extractability as a sensitive probe for the thermal denaturation of carp myofibrils. *Nippon Suisan Gakkaishi* 59:515-518 (in Japanese with English Abstract)
7. Harafuji, H., Ogawa, Y. (1980) Re-examination of the apparent binding constant of ethylene glycol bis ( $\beta$ -aminoethyl ether)-N,N,N',N'-tetraacetic acid with calcium around neutral pH. *J Biochem* 87:1305-1312
8. Ooizumi, T., Hashimoto, A., Ogura, J., Arai, K. (1981) Quantitative aspect for protective effect of sugar and sugar alcohol against denaturation of fish myofibrils. *Nippon Suisan Gakkaishi* 47:901-908 (in Japanese with English Abstract)
9. Konno, K., Arai, K., Yoshida, M., Watanabe, S. (1981) Calcium regulation in squid mantle and scallop adductor muscles. *J Biochem* 89:581-569
10. Yoshioka, T., Kinoshita, Y., Yoshino, H., Park, S., Konno, K., Seki, N. (2003) Changes in translucency of squid mantle muscle upon storage. *Fish Sci* 69:408-413

## Legends of figures

Fig. 1 Effect of KCl concentration on the stabilizing effect of  $\text{Ca}^{2+}$ . (a); Squid myosin (circles) or myofibrils (triangles) were heated in the presence of 1 mM  $\text{CaCl}_2$  (open symbols) or EDTA (closed symbols) at 25 °C, and the inactivation rates were estimated. (b); The same experiments as in (a) were conducted only with myofibrils in the presence of 1 mM  $\text{CaCl}_2$  (open symbols) or EDTA (closed symbols) at 35 °C

Fig. 2 Temperature dependent stabilizing effect of  $\text{Ca}^{2+}$  on myosin and myofibrils. Myosin (circles) and myofibrils (triangles) of squid were suspended in a medium containing 0.1 M KCl, 20 mM Tris-HCl (pH 7.5). Thermal inactivation rates for the two preparations were estimated by changing heating temperatures. Open and closed symbols denote the heating in media containing 1 mM  $\text{CaCl}_2$  and EDTA, respectively.

Fig. 3  $\text{Ca}^{2+}$  concentration dependent stabilization of squid myosin and myofibrils. Squid myosin (triangles) and myofibrils (circles) were suspended in a medium containing 0.1 M KCl, 20 mM Tris-HCl (pH 7.5) and 0.1 mM  $\text{CaCl}_2$  and desired concentrations of EGTA to give desired free  $\text{Ca}^{2+}$  concentrations. Heating temperatures for myosin and for myofibrils were 25 and 35 °C, respectively.

Fig. 4 Effect of sorbitol on the stabilizing effect of  $\text{Ca}^{2+}$  on myosin and myofibrils. Squid myofibrils (a) and myosin (b) were suspended in 0.1 M KCl, 20 mM Tris-HCl (pH 7.5) in media containing 1 mM  $\text{CaCl}_2$  (open) or EDTA (closed), and heated in the presence of various concentrations of sorbitol. Heating temperatures for myosin and myofibrils were 30 and 40 °C, respectively. The inactivation rates in logarithmic values were plotted against sorbitol concentrations added.

Fig. 5 Temperature dependent stabilizing effect of  $\text{Ca}^{2+}$  on myosin with and without sorbitol. Squid myosin with either 1 mM  $\text{CaCl}_2$  (open) or EDTA (closed) was heated as in Fig. 2. The heating media also contained 1.8 M sorbitol (squares). The same results without sorbitol (circles) as presented in Fig. 2 were shown again.

Table 1 Stabilizing effect of Ca<sup>2+</sup> on myofibrils and on myosin under various heating conditions.

Sample	Temperature (°C)	Sorbitol (M)	Inactivation rate in Ca-medium(sec <sup>-1</sup> )	Stabilization*
Myofibrils	40		2.5 x 10 <sup>-5</sup>	100
Myosin	40	1.3	1 x 10 <sup>-5</sup>	100
	30		3.5 x 10 <sup>-3</sup>	5
	20		4.5 x 10 <sup>-5</sup>	0
	30	1.8	1.2 x 10 <sup>-5</sup>	0
	40	1.8	1.4 x 10 <sup>-3</sup>	4.4

\* The extent was expressed as "Rate in EDTA-medium/Rate in Ca-medium"









