



HOKKAIDO UNIVERSITY

Title	Formation of 5-formyl-2'-deoxycytidine from 5-methyl-2'-deoxycytidine in duplex DNA by Fenton-type reactions and gamma-irradiation
Author(s)	Murata-Kamiya, Naoko; Kamiya, Hiroyuki; Karino, Naoko et al.
Citation	Nucleic Acids Research, 27(22), 4385-4390 https://doi.org/10.1093/nar/27.22.4385
Issue Date	1999-11
Doc URL	https://hdl.handle.net/2115/46962
Type	journal article
File Information	236_Kamiya NAR 99.pdf



Formation of 5-formyl-2'-deoxycytidine from 5-methyl-2'-deoxycytidine in duplex DNA by Fenton-type reactions and γ -irradiation

Naoko Murata-Kamiya, Hiroyuki Kamiya¹, Naoko Karino², Yoshihito Ueno², Hiroshi Kaji, Akira Matsuda² and Hiroshi Kasai^{1,*}

Department of Health Policy and Management and ¹Department of Environmental Oncology, Institute of Industrial Ecological Sciences, University of Occupational and Environmental Health, 1-1, Iseigaoka, Yahatanishi-ku, Kitakyushu 807-8555, Japan and ²Graduate School of Pharmaceutical Sciences, Hokkaido University, Kita-12, Nishi-6, Kita-ku, Sapporo 060-0812, Japan

Received August 10, 1999; Revised and Accepted October 1, 1999

ABSTRACT

5-Methyl-2'-deoxycytidine (5-Me-dC) is formed by the enzymatic methylation of dC, primarily in CpG sequences in DNA, and is involved in the regulation of gene expression. In the present study, 5-Me-dC and double-stranded DNA fragments containing 5-Me-dC were either γ -irradiated or aerobically treated with Fenton-type reagents, Fe(II)-EDTA, Fe(II)-nitritotriacetic acid, Fe(III)-EDTA-H₂O₂-catechol or ascorbic acid-H₂O₂ under neutral conditions. The formation of 5-formyl-2'-deoxycytidine (5-CHO-dC) was observed upon treatment of both 5-Me-dC and DNA fragments containing 5-Me-dC. The yields of 5-CHO-dC from 5-Me-dC and those of 5-formyl-2'-deoxyuridine from dT were comparable. These results suggest that 5-Me-dC in DNA is as susceptible to oxidation as dT in cells, and raise the possibility that 5-CHO-dC may contribute to the high mutagenic rate observed in CpG sequences in genomic DNA.

INTRODUCTION

5-Methyl-2'-deoxycytidine (5-Me-dC) is a minor component found in most eukaryotic DNA. In humans, 5% of all dC residues are enzymatically methylated. Methylation of dC occurs after DNA synthesis, primarily in CpG sequences (1,2). The 5-Me-dC in a CpG sequence is involved in the regulation of gene expression (3–6) and the silencing of invading viral genomes (7,8). Interestingly, CpG sequences are hot spots for mutations: the predominant mutation in the human p53 gene is a C to T transition in a CpG sequence (9,10). This event has been thought to be the result of spontaneous or methylase-catalyzed hydrolytic deamination of 5-Me-dC to thymidine (11–13). However, this C to T transition could occur by other mechanisms as well as by the deamination of 5-Me-dC.

Recently, it was reported that UV-A irradiation of 5-Me-dC in the presence of a photosensitizer leads to the formation of various compounds, including 5-formyl-2'-deoxycytidine (5-CHO-dC)

(14). In addition, the formation of 5-formylcytosine was observed upon exposure of 5-methylcytosine to UV light (15). 5-CHO-dC in DNA may potentially be as miscoding and mutagenic as the dT analog, 5-formyl-2'-deoxyuridine (5-CHO-dU) (16–20). The presence of an intramolecular hydrogen bond between the carbonyl of the 5-formyl group and the 4-amino function may affect the equilibrium between the amino- and imino-tautomers. (21,22). We supposed that the formation of 5-CHO-dC by reactive oxygen species (ROS) might be involved in the mutation process in the CpG sequences. Thus, it is important to know the formation rates of 5-CHO-dC from 5-Me-dC in various ROS-generating systems using Fenton-type reactions, which are models for oxidation reactions in cells. In particular, an investigation of the formation of 5-CHO-dC in DNA is of great importance.

In the present study, 5-Me-dC and double-stranded DNA fragments containing 5-Me-dC were either γ -irradiated or aerobically treated with Fenton-type reagents [Fe(II)-EDTA, Fe(II)-nitritotriacetic acid (NTA), Fe(III)-EDTA-H₂O₂-catechol or ascorbic acid-H₂O₂] under neutral conditions, and the formation of 5-CHO-dC from 5-Me-dC and that of 5-CHO-dU from dT were compared. We found that 5-CHO-dC was formed as efficiently as 5-CHO-dU, a major oxidation product of dT, in the monomeric form and in DNA fragments. These results suggest that 5-CHO-dC may be formed in CpG sequences in cells and may induce C to T transition mutations.


MATERIALS AND METHODS

Nucleosides

The 5-CHO-dU standard was synthesized as described previously (16). The 5-CHO-dC (5) standard was prepared as shown in Scheme 1. Pd(II)-catalyzed vinylation of the 5-iodouracil derivative **1**, followed by transformation of the 4-oxo group into an amino group using a known method, gave the 5-vinylcytosine derivative **3** in a 65% yield in two steps (23). Compound **3** was then converted to the 5-(1,2-dihydroxyethyl)cytosine derivative **4** in a 71% yield with OsO₄ in the presence of *N*-methylmorpholine *N*-oxide. After deprotection of the TBS

*To whom correspondence should be addressed. Tel: +81 93 691 7468; Fax: +81 93 601 2199; Email: h-kasai@med.uoeh-u.ac.jp

groups at the sugar moiety with NH_4F in MeOH, the resulting nucleoside was treated with NaIO_4 to give the desired 5-CHO-dC (**5**) in a 92% yield as a crystalline product. Previously, the photosensitized oxidation of 5-Me-dC by 2-methyl-1,4-naphthoquinone was reported to give **5** in a 13% yield, which was assigned by NMR and mass spectrometric analyses (14), and these data are consistent with those of **5** described in this paper.



Scheme 1. Preparation of 5-CHO-dC: (a) tributyl(vinyl)tin, $(\text{PPh}_3)_2\text{PdCl}_2$, DMF, 80°C ; (b) (1) TPSCl , Et_3N , DMAP, CH_3CN , (2) 25% NH_4OH , room temperature, 65% from **1**; (c) OsO_4 , *N*-methylmorpholine *N*-oxide, aqueous acetone + *t*-BuOH, room temperature, 71%; (d) (1) NH_4F , MeOH, reflux, (2) NaIO_4 , H_2O , room temperature, 92%.

3',5'-Bis-O-(tert-butylidimethylsilyl)-5-vinyl-2'-deoxycytidine (3). A mixture of **1** (3.32 g, 5.69 mmol), $(\text{PPh}_3)_2\text{PdCl}_2$ (399 mg, 0.569 mmol) and tributyl(vinyl)tin (2.00 ml, 6.84 mmol) in DMF (30 ml) was stirred at 80°C for 1.5 h. The reaction mixture was filtered through a Celite pad, and the filtrate was taken up in EtOAc (120 ml). The organic layer was washed with H_2O (3×40 ml) and brine (40 ml), dried (Na_2SO_4) and evaporated under reduced pressure. The residue was roughly purified by column chromatography (SiO_2 , 5–20% EtOAc in hexane) to give solids (2.40 g) as a mixture of **2** and the tributyltin derivatives. A solution of the above solids (2.40 g), Et_3N (2.08 ml, 14.9 mmol), DMAP (1.82 g, 14.9 mmol) and 2,4,6-triisopropylbenzenesulfonyl chloride (4.51 g, 14.9 mmol) in CH_3CN (40 ml) was stirred at room temperature for 24 h. The mixture was cooled in an ice-bath. Concentrated NH_4OH (25%, 60 ml) was added, and the mixture was stirred at room temperature for 2 h. The mixture was concentrated under reduced pressure and was taken up in EtOAc (100 ml), which was washed with H_2O (2×40 ml) and brine (40 ml). The separated organic layer was dried (Na_2SO_4) and evaporated under reduced pressure. The residue was purified by column chromatography (SiO_2 , 0–4% MeOH in CHCl_3) to give **3** (1.78 g, 65% from **1** as a foam): $^1\text{H NMR}$ (270 MHz, CDCl_3) δ 7.80 (s, 1H, H-6), 6.39–6.27 (m, 2H, H-1' and 5-CH=CH₂), 5.44 (d, 1H, 5-CH=CH₂, $J = 17.2$ Hz), 5.30 (d, 1H, 5-CH=CH_E, $J = 11.2$ Hz), 4.36 (m, 1H, H-3'), 3.97 (m, 1H, H-4'), 3.87 (dd, 1H, H-5'a, $J = 3.3$, 11.2 Hz), 3.76 (dd, 1H, H-5'b, $J = 2.6$, 11.2 Hz), 2.48 (ddd, 1H, H-2'a, $J = 3.3$, 5.9, 13.2 Hz), 1.98 (ddd, 1H, H-2'b, $J = 6.6$, 6.6, 13.2 Hz), 0.89 and 0.89 (each s, each 9H, Me), 0.08–0.06 (m, 12H, Me); FABMS m/z 482 (M+H)⁺; analysis calculated

for $\text{C}_{23}\text{H}_{43}\text{N}_3\text{O}_4\text{Si}_2$: C, 57.34; H, 9.00; N, 8.72. Found: C, 57.11; H, 8.84; N, 8.47.

3',5'-Bis-O-(tert-butylidimethylsilyl)-5-(1,2-dihydroxyethyl)-2'-deoxycytidine (4). A solution of OsO_4 in *t*-BuOH (5 mg/ml in *t*-BuOH, 9.35 ml, 0.184 mmol) was added to a solution of **3** (1.77 g, 3.68 mmol) and *N*-methylmorpholine *N*-oxide (647 mg, 5.52 mmol) in acetone- H_2O -*t*-BuOH (4:1:1, 80 ml), and the resulting mixture was stirred at room temperature for 7 h. After the reaction was quenched with saturated aqueous $\text{Na}_2\text{S}_2\text{O}_3$ (20 ml), the mixture was extracted with EtOAc (100 ml). The organic layer was washed with saturated aqueous $\text{Na}_2\text{S}_2\text{O}_3$ (2×40 ml) and brine (30 ml), dried (Na_2SO_4) and evaporated under reduced pressure. The residue was purified by column chromatography (SiO_2 , 0–10% MeOH in CHCl_3) to give **4** (1.36 g, 71% as a foam): $^1\text{H NMR}$ (270 MHz, CDCl_3) δ 7.67 and 7.65 (each s, 0.3H and 0.7H, H-6), 6.23 (m, 1H, H-1'), 4.58 (m, 1H, 5-CHOH), 4.34 (m, 1H, H-3'), 3.94 (m, 1H, H-4'), 3.87–3.72 (m, 4H, H-5'a,b and 5-CHOH-CH₂OH), 2.39 (m, 1H, H-2'a), 1.91 (m, 1H, H-2'b), 0.91 and 0.88 (each s, each 9H, Me), 0.10–0.06 (m, 12H, Me); HRMS (FAB) calculated for $\text{C}_{23}\text{H}_{46}\text{N}_3\text{O}_6\text{Si}_2$ 516.2925 (M+H)⁺, found 516.2935.

5-Formyl-2'-deoxycytidine (5). A mixture of **4** (820 mg, 1.59 mmol) and NH_4F (1.18 g, 31.9 mmol) in MeOH (20 ml) was heated under reflux for 16 h and was evaporated under reduced pressure. The residue was partitioned between CHCl_3 (30 ml) and H_2O (50 ml). The aqueous layer was washed with CHCl_3 (2×30 ml) and was concentrated under reduced pressure. Sodium periodate (680 mg, 3.18 mmol) was added to the above solution, which was stirred at room temperature for 30 min. After the reaction was quenched with saturated aqueous $\text{Na}_2\text{S}_2\text{O}_3$ (5 ml), H_2O (90 ml) was added to dissolve the precipitates. The mixture was absorbed onto an activated charcoal column, which was washed well with H_2O and then with 50–100% MeOH in H_2O to give **5** (374 mg, 92% as a white powder, which was crystallized from MeOH): no definite melting point (colored from 180 to 300°C); UV (MeOH) λ_{max} 283 nm (ϵ 11 100), 226 nm (ϵ 28 700), λ_{min} 266 nm (ϵ 7400); $^1\text{H NMR}$ (400 MHz, $\text{DMSO}-d_6$) δ 9.39 (s, 1H, CHO), 8.77 (s, 1H, H-6), 8.04 (br s, 1H, 4-NH, D_2O exchangeable), 7.79 (br s, 1H, 4-NH, D_2O exchangeable), 6.00 (t, 1H, H-1', $J = 6.0$ Hz), 5.18 (br s, 2H, 3',5'-OH, D_2O exchangeable), 4.16 (m, 1H, H-3'), 3.80 (dd, 1H, H-4', $J = 3.6$, 7.3 Hz), 3.61 (dd, 1H, H-5'a, $J = 3.6$, 12.0 Hz), 3.53 (dd, 1H, H-5'b, $J = 3.9$, 12.0 Hz), 2.24 (ddd, 1H, H-2'a, $J = 4.8$, 5.6, 13.1 Hz), 2.03 (ddd, 1H, H-2'b, $J = 5.9$, 6.6, 13.1 Hz); HRMS (FAB) calculated for $\text{C}_{10}\text{H}_{14}\text{N}_3\text{O}_5$ 256.0933 (M+H)⁺, found 256.0920. Analysis calculated for $\text{C}_{10}\text{H}_{13}\text{N}_3\text{O}_5$: C, 47.06; H, 5.13; N, 16.46. Found: C, 46.91; H, 5.07; N, 16.18.

Oligonucleotides

3'-O-[2-Cyanoethyl(*N,N*-isopropyl)phosphinyl]-5'-O-dimethoxytrityl-*N*⁶-benzoyl-2'-deoxy-5-methylcytidine was prepared from 5-Me-dC using a known method (24,25). Oligodeoxyribonucleotides containing 5-Me-dC were synthesized and purified as described previously (26). The following three self-complementary oligonucleotides were synthesized: 5'-dCGMGAATTMGCG-3' (Seq-1); 5'-dCGCGAATTMGCG-3' (Seq-2); and 5'-dCGMG-AATTCGCG-3' (Seq-3) (M represents 5-methylcytosine). These were annealed to form duplexes before the treatments.

Table 1. Yields of oxidized products formed by reactive oxygen species^a

Treatment	Nucleosides		DNA	
	5-CHO-dC	5-CHO-dU	5-CHO-dC	5-CHO-dU
Fe(II)-EDTA ^b	2.7 ± 0.1	4.1 ± 0.1	1.4 ± 0.03	1.6 ± 0.1
Fe(II)-NTA ^c	1.8 ± 0.1	2.7 ± 0.1	0.94 ± 0.01	1.3 ± 0.1
Fe(III)-EDTA-H ₂ O ₂ -catechol ^d	5.8 ± 0.2	0.38 ± 0.04	1.3 ± 0.1	0.87 ± 0.22
Ascorbic acid-H ₂ O ₂ ^e				
7.5 mM H ₂ O ₂	0.35 ± 0.03	0.44 ± 0.01	ND	ND
75 mM H ₂ O ₂	0.67 ± 0.18	0.03 ± 0.01	ND	ND
γ-ray ^f	0.97 ± 0.04	0.24 ± 0.02	1.4 ± 0.1	0.82 ± 0.11

^aYields shown are the ratios (%) of an oxidized nucleoside versus total (oxidized plus starting) nucleosides. The values represent the average of three separate experiments, with standard deviations.

^bNucleosides or a DNA fragment (Seq-1) were treated with Fe(II)-EDTA for 30 min.

^cNucleosides or a DNA fragment (Seq-1) were treated with Fe(II)-NTA for 30 min.

^dNucleosides or a DNA fragment (Seq-1) were treated with Fe(III)-EDTA-H₂O₂-catechol for 120 min.

^eNucleosides were treated with ascorbic acid-H₂O₂ for 4 h.

^fNucleosides or a DNA fragment (Seq-1) were treated with 400 Gy γ-ray irradiation.

ND, not determined.

Treatment of deoxyribonucleosides and double-stranded DNA fragments with Fenton-type reagents

A single deoxyribonucleoside, dT or 5-Me-dC (1 mM), or a double-stranded DNA fragment (11 μM) was treated with Fe(II)-EDTA (5 mM FeSO₄ and 5 mM EDTA), Fe(II)-NTA (5 mM FeSO₄ and 20 mM NTA), Fe(III)-EDTA-H₂O₂-catechol (0.1 mM FeCl₃, 0.5 mM EDTA, 5 mM H₂O₂ and 1 mM catechol) or ascorbic acid-H₂O₂ (14.2 mM ascorbic acid and 7.5 or 75 mM H₂O₂) in 50 mM sodium phosphate buffer (pH 7.0) at 37°C (for deoxyribonucleosides) or 25°C (for DNA fragments) for the desired period of time. The FeSO₄ solution was prepared immediately before addition to the other components. No pH change was observed either before or after the incubation.

γ-Irradiation

Irradiations of a single deoxyribonucleoside, dT or 5-Me-dC (1 mM), or a double-stranded DNA fragment (11 μM) in 50 mM sodium phosphate buffer (pH 7.0) were carried out with a Gammacell 40 Exactor with a ¹³⁷Cs γ-ray source (Nordion International Inc.) at a dose rate of 1 Gy/min (total 100–400 Gy).

HPLC analysis

Before the analysis by HPLC, the treated DNA fragments were purified with a Sep-Pak plus (Millipore) column. They were digested with snake venom phosphodiesterase (0.15 U/ml, Boehringer Mannheim) and calf intestine alkaline phosphatase (50 U/ml, Boehringer Mannheim) in 50 mM HEPES-NaOH buffer (pH 7.2) at 37°C for 2 h. After the incubation, the reaction mixture was passed through a Micropure-EZ filter (Millipore) to remove the enzymes.

The resulting nucleoside mixture from the DNA fragments was injected into a reverse-phase HPLC column [YMC-pack ODS-AM303 (4.6 × 250 mm)] connected with a photo-diode array UV detector (Hewlett Packard 1100 HPLC Detection System). In the case of 5-Me-dC or dT treated by ROS, the reaction mixture was injected directly. The following linear

gradient of CH₃CN concentrations in 10 mM NaH₂PO₄ (pH 7.0) was used: 0–25 min, linear gradient of CH₃CN (0–12.5%); 25–26 min, linear gradient of CH₃CN (12.5–100%); and 26–31 min, 100% CH₃CN. The flow rate was 1.0 ml/min and the column temperature was 25°C.

The reactions and analyses were conducted in triplicate, and good reproducibilities were obtained.

RESULTS

Treatment of deoxyribonucleosides and DNA fragments with Fe(II)-EDTA

We first treated the nucleosides, 5-Me-dC and dT, separately with a Fenton-type reagent, Fe(II)-EDTA, and analyzed the reaction mixtures directly by reverse-phase HPLC. When 5-Me-dC was treated with Fe(II)-EDTA at 37°C for 30 min, a peak corresponding to 5-CHO-dC was detected (Fig. 1A). In the case of dT, 5-CHO-dU was identified as reported previously (27) (Fig. 1B). These assignments were made by comparison of their chromatographic and spectral properties with those of the standard samples.

We then examined the time course of 5-CHO-dC and 5-CHO-dU formation from 5-Me-dC and dT, respectively. 5-Me-dC or dT was treated with Fe(II)-EDTA at 37°C for 5, 10, 20 and 30 min. As shown in Figure 2, the yield of each oxidized product reached an almost maximal level after 5 min. In this system, it is likely that auto-oxidation of Fe(II) to Fe(III) ions occurs within 5 min. During the 30 min incubation, the yield of 5-CHO-dC reached 2.7%, almost two-thirds that of 5-CHO-dU (Table 1).

Next, we treated a double-stranded DNA fragment (Seq-1) with Fe(II)-EDTA. This sequence is the same as that of Dickerson's dodecamer, which adopts the canonical B-DNA conformation (28–30) and contains four CpG sequences. We introduced 5-Me-dC at positions 3 and 9 (Materials and Methods). We thought that this duplex would be a good model for methylated DNA.




Figure 1. HPLC analysis of the reaction mixture of (A) 5-Me-dC and (B) dT with Fe(II)-EDTA at 37°C for 30 min. The following linear gradients of CH₃CN concentrations in 10 mM NaH₂PO₄ (pH 7.0) were used: 0–25 min, linear gradient of CH₃CN (0–12.5%); 25–26 min, linear gradient of CH₃CN (12.5–100%); 26–31 min, 100% CH₃CN. Detection was performed with UV absorbance at 284 nm. The UV spectra of 5-CHO-dC and 5-CHO-dU are also shown.




Figure 2. Time course of 5-CHO-dC (open circles) and 5-CHO-dU (closed circles) formation by Fe(II)-EDTA from 5-Me-dC and dT, respectively. The 5-Me-dC and dT were treated as described in Materials and Methods.

When this DNA fragment was treated with Fe(II)-EDTA, and then digested by a nuclease and a phosphatase, the formation of 5-CHO-dC was observed. Thus, 5-CHO-dC was generated by the oxidation of 5-Me-dC in the DNA. The yield of 5-CHO-dC was similar to that of 5-CHO-dU. The yields of both products were higher in the monomeric form than in the DNA fragment.

To investigate whether 5-CHO-dC formation has sequence specificity, we compared the yields of 5-CHO-dC in the two DNA fragments, Seq-2 and Seq-3. Seq-2 has 5-Me-dC at position 9 (5'-TMG-3': M represents 5-Me-dC), whereas Seq-3 has 5-Me-dC at position 3 (5'-GMG-3'). The yields of 5-CHO-dC were 1.7%

for Seq-2 and 1.3% for Seq-3. The yields of 5-CHO-dU were 2.2% for Seq-2 and 2.0% for Seq-3. The ratios of 5-CHO-dC to 5-CHO-dU were 0.77 for Seq-2 and 0.65 for Seq-3, suggesting the absence of a neighboring base effect.

Treatment of deoxyribonucleosides and a DNA fragment with Fe(II)-NTA

We then treated the nucleosides and a double-stranded DNA fragment with Fe(II)-NTA. The formation of 5-CHO-dC and 5-CHO-dU was observed upon treatment of 5-Me-dC and dT, respectively, with Fe(II)-NTA (Table 1). These products were also detected when the duplex Seq-1 was treated with Fe(II)-NTA (Table 1). As with Fe(II)-EDTA, 5-CHO-dC was produced as efficiently as 5-CHO-dU.

Treatment of deoxyribonucleosides and a DNA fragment with Fe(III)-EDTA-H₂O₂-catechol

Next, we examined the time course of 5-CHO-dC and 5-CHO-dU formation from 5-Me-dC and dT, respectively, in another Fenton-type reaction. 5-Me-dC and dT were treated with Fe(III)-EDTA-H₂O₂-catechol at 37°C for 30, 60, 90 and 120 min. The Seq-1 duplex was tested similarly at 25°C. As shown in Figure 3, the yield of each oxidized product increased in a time-dependent manner during the 120 min incubation, for both the nucleosides and the DNA fragment. With Fe(III)-EDTA-H₂O₂-catechol, 5-CHO-dC was produced more efficiently than 5-CHO-dU (15-fold in the monomeric form and 1.5-fold in the DNA fragment) (Table 1). The yield of 5-CHO-dU was higher in DNA than in the monomeric form, whereas 5-CHO-dC was generated more efficiently in the monomeric form than in the DNA.




Figure 3. Time course of 5-CHO-dC (open circles) and 5-CHO-dU (closed circles) formation from 5-Me-dC and dT, respectively, and that of 5-CHO-dC (open squares) and 5-CHO-dU (closed squares) formation in duplex DNA by Fe(III)-EDTA-H₂O₂-catechol. The 5-Me-dC, dT and duplex DNA were treated as described in Materials and Methods.




Figure 4. Time course of 5-CHO-dC (open circles) and 5-CHO-dU (closed circles) formation from 5-Me-dC and dT, respectively, by ascorbic acid-7.5 mM H₂O₂, and that of 5-CHO-dC (open triangles) and 5-CHO-dU (closed triangles) formation from 5-Me-dC and dT, respectively, by ascorbic acid-75 mM H₂O₂. 5-Me-dC and dT were treated as described in Materials and Methods.

Treatment of deoxyribonucleosides with ascorbic acid-H₂O₂

5-Me-dC or dT was treated with ascorbic acid-H₂O₂ (14.2 mM ascorbic acid and 7.5 or 75 mM H₂O₂) at 37°C for 1, 2, 3 and 4 h. With both concentrations of H₂O₂, the formation of 5-CHO-dC and 5-CHO-dU was observed. As shown in Figure 4, the yield of 5-CHO-dC increased in a time-dependent manner during a 4 h incubation with 7.5 mM H₂O₂. As in the case of 5-CHO-dC, the steady formation of 5-CHO-dU was observed with 7.5 mM H₂O₂. However, the amount of 5-CHO-dU was decreased after 1 h with 75 mM H₂O₂. In the case of 5-CHO-dC, its yield increased up to 3 h, and then decreased with 75 mM H₂O₂. Paradoxically, the yield of 5-CHO-dU was higher with the lower H₂O₂ concentration than with the higher H₂O₂ concentration, indicating that the 5-CHO-dU was degraded during the incubation with ascorbic acid-75 mM H₂O₂. 5-CHO-dC was produced at least as efficiently as 5-CHO-dU by this type of treatment (Table 1).




Figure 5. The formation of 5-CHO-dC (open circles) and 5-CHO-dU (closed circles) from 5-Me-dC and dT, respectively, and that of 5-CHO-dC (open squares) and 5-CHO-dU (closed squares) in duplex DNA by γ -irradiation. The 5-Me-dC, dT and duplex DNA were exposed to γ -rays as described in Materials and Methods.

Treatment of deoxyribonucleosides and a DNA fragment with γ -rays

Next, we tested whether 5-CHO-dC was generated by ionizing radiation. As expected, the formation of 5-CHO-dC and 5-CHO-dU was observed upon exposure of 5-Me-dC and dT, respectively, to γ -rays. As shown in Figure 5, the yield of each oxidized product increased up to 400 Gy-irradiation. A linear increase in the formation of 5-CHO-dC was observed in the range 0–400 Gy. However, the formation of 5-CHO-dU from dT showed no linearity. Similar results were obtained when the DNA fragment, Seq-1, was γ -irradiated (Fig. 5). The yield of each oxidized product was higher in the DNA fragment than in the monomeric form. 5-CHO-dC was produced more efficiently than 5-CHO-dU.

DISCUSSION

In the present study, we investigated the formation of 5-CHO-dC from 5-Me-dC by Fenton-type reactions and γ -irradiation of both the monomeric form and duplex DNA fragments. We found that 5-CHO-dC was produced as efficiently as 5-CHO-dU. In particular, the formation of 5-CHO-dC was demonstrated, for the first time, in DNA fragments. The ratios of 5-CHO-dC to 5-CHO-dU differed, depending on the ROS-generating system used (Table 1). These results suggest that the ROS generated by these five reactions are not identical. Similar phenomena were observed for other oxidative products of DNA components (27).

Fenton-type reactions are models for oxidation reactions in cells. We found that 5-CHO-dC was formed with three Fenton-type reactions, Fe(II)-EDTA, Fe(II)-NTA and Fe(III)-EDTA-H₂O₂-catechol, under neutral conditions. With both Fe(II)-EDTA and Fe(II)-NTA, 5-CHO-dC was produced as efficiently as 5-CHO-dU (Table 1). With Fe(III)-EDTA-H₂O₂-catechol, 5-CHO-dC was produced more efficiently than 5-CHO-dU, especially in the monomeric form (Table 1). Because we previously found that 5-CHO-dU is generated more efficiently than 8-hydroxy-2'-deoxyguanosine (8-OH-dG) with Fenton-type reactions (27), 5-CHO-dC appears to be formed as efficiently as 8-OH-dG. These results

suggest that 5-CHO-dC is generated in cytosine methylation sites in cellular DNA by the endogenous formation of ROS.

With ascorbic acid-H₂O₂, the ratio of 5-CHO-dC to 5-CHO-dU dramatically changed depending upon the H₂O₂ concentration. With 7.5 mM H₂O₂, the production of 5-CHO-dC was similar to that of 5-CHO-dU (Table 1). However, with 75 mM H₂O₂, the yield of 5-CHO-dC was 8- to 27-fold greater than that of 5-CHO-dU (Fig. 4). The fact that the yield of 5-CHO-dU was higher with 7.5 than 75 mM H₂O₂ suggests that the 5-CHO-dU was degraded under the more stringent conditions. 5-CHO-dU was produced as efficiently as 5-CHO-dC with 100 Gy γ -irradiation. However, the relative yield of 5-CHO-dU to 5-CHO-dC was decreased with increased doses of irradiation (Fig. 5). Taken together, these results suggest that 5-CHO-dU was susceptible to further oxidation.

We found that 5-CHO-dC was formed as efficiently as 5-CHO-dU. 5-CHO-dU has been detected in γ -irradiated DNA (16,31). Mammalian and *Escherichia coli* cells possess repair activities for 5-CHO-dU (32–34). 5-Hydroxymethyl-2'-deoxycytidine, another type of oxidized 5-Me-dC, is present in calf thymus DNA (35), and a 5-hydroxymethylcytosine DNA glycosylase activity was found in mammalian tissue (36). Thus, it would be very interesting to determine whether 5-CHO-dC is formed and if repair activities for 5-CHO-dC are present in mammalian cells.

We supposed that 5-CHO-dC formation might be one cause of the C to T transition mutations frequently found in cytosine methylation sites. It was reported that an intramolecular hydrogen bond between the carbonyl of the 5-formyl group and the 4-amino function is present in the 5-formylcytosine moiety of its ribonucleoside (21,22). The equilibrium between the amino- and imino-tautomers may be affected by this hydrogen bonding, and may allow the formation of a pair with adenine. This possibility and the recognition of 5-CHO-dC by DNA polymerases are of great interest.

CONCLUSION

5-CHO-dC was found to be formed from 5-Me-dC as efficiently as 5-CHO-dU from dT in duplex DNA under various oxidation reactions. This oxidation product of 5-Me-dC might be one cause of the mutations at cytosine methylation sites. We are currently investigating the mutagenicity of this oxidized base to understand its role in mutagenesis and carcinogenesis.

ACKNOWLEDGEMENTS

We would like to thank Dr Hiroshi Sugiyama of the Tokyo Medical and Dental University for his advice. This work was supported in part by a Grant-in-Aid from the Ministry of Education, Science, Sports and Culture, Japan.

REFERENCES

- Razin,A. and Riggs,A.D. (1980) *Science*, **210**, 604–610.
- Verdine,G.L. (1994) *Cell*, **76**, 197–200.
- Cedar,H. (1988) *Cell*, **53**, 3–4.
- Hergersberg,M. (1991) *Experientia*, **47**, 1171–1185.
- Boyes,J. and Bird,A. (1991) *Cell*, **64**, 1123–1134.
- Barlow,D.P. (1993) *Science*, **260**, 309–310.
- Groudine,M., Eisenman,R. and Weintraub,H. (1981) *Nature*, **292**, 311–317.
- Doerfler,W. (1991) *Biol. Chem. Hoppe Seyler*, **372**, 557–564.
- Rideout,W.M.,III, Coetzee,G.A., Olumi,A.F. and Jones,P.A. (1990) *Science*, **249**, 1288–1290.
- Magewu,A.N. and Jones,P.A. (1994) *Mol. Cell. Biol.*, **14**, 4225–4232.
- Shen,J.C., Rideout,W.M.,III and Jones,P.A. (1994) *Nucleic Acids Res.*, **22**, 972–976.
- Wang,R.Y., Kuo,K.C., Gehrke,C.W., Huang,L.H. and Ehrlich,M. (1982) *Biochim. Biophys. Acta*, **697**, 371–377.
- Yebra,M.J. and Bhagwat,A.S. (1995) *Biochemistry*, **34**, 14752–14757.
- Bienvu,C., Wagner,J.R. and Cadet,J. (1996) *J. Am. Chem. Soc.*, **118**, 11406–11411.
- Privat,E. and Sowers,L.C. (1996) *Chem. Res. Toxicol.*, **9**, 745–750.
- Kasai,H., Iida,A., Yamaizumi,Z., Nishimura,S. and Tanooka,H. (1990) *Mutat. Res.*, **243**, 249–253.
- Ono,A., Okamoto,T., Inada,M., Nara,H. and Matsuda,A. (1994) *Chem. Pharm. Bull. (Tokyo)*, **42**, 2231–2237.
- Yoshida,M., Makino,K., Morita,H., Terato,H., Ohyama,Y. and Ide,H. (1997) *Nucleic Acids Res.*, **25**, 1570–1577.
- Zhang,Q.M., Sugiyama,H., Miyabe,I., Matsuda,S., Saito,I. and Yonei,S. (1997) *Nucleic Acids Res.*, **25**, 3969–3973.
- Fujikawa,K., Kamiya,H. and Kasai,H. (1998) *Nucleic Acids Res.*, **26**, 4582–4587.
- Kawai,G., Yokogawa,T., Nishikawa,K., Ueda,T., Hashizume,T., McCloskey,J.A., Yokoyama,S. and Watanabe,K. (1994) *Nucl. Nucl.*, **13**, 1189–1199.
- LaFrancois,C.J., Fujimoto,J. and Sowers,L.C. (1998) *Chem. Res. Toxicol.*, **11**, 75–83.
- Sugiyama,H., Matsuda,S., Kino,K., Zhang,Q.M., Yonei,S. and Saito,I. (1996) *Tetrahedron Lett.*, **37**, 9067–9070.
- Sinha,N.D., Biernat,J. and Koster,H. (1983) *Tetrahedron Lett.*, **24**, 5843–5846.
- Atkinson,T. and Smith,M. (1984) In Gait,M.J. (ed.), *Oligonucleotides Synthesis: A Practical Approach*. IRL Press, Oxford, pp. 35–81.
- Haginoya,N., Ono,A., Nomura,Y., Ueno,Y. and Matsuda,A. (1997) *Bioconj. Chem.*, **8**, 271–280.
- Murata-Kamiya,N., Kamiya,H., Muraoka,M., Kaji,H. and Kasai,H. (1997) *J. Radiat. Res. (Tokyo)*, **38**, 121–131.
- Giessner-Prettre,C. and Pullman,B. (1982) *Biochem. Biophys. Res. Commun.*, **107**, 1539–1544.
- Chen,Y.Z., Mohan,V. and Griffey,R.H. (1998) *J. Biomol. Struct. Dyn.*, **15**, 765–777.
- Ikeda,H., Fernandez,R., Wilk,A., Barchi,J.J.,Jr, Huang,X. and Marquez,V.E. (1998) *Nucleic Acids Res.*, **26**, 2237–2244.
- Douki,T., Delatour,T., Paganon,F. and Cadet,J. (1996) *Chem. Res. Toxicol.*, **9**, 1145–1151.
- Bjelland,S., Birkeland,N.K., Benneche,T., Volden,G. and Seeberg,E. (1994) *J. Biol. Chem.*, **269**, 30489–30495.
- Bjelland,S., Eide,L., Time,R.W., Stote,R., Eftedal,I., Volden,G. and Seeberg,E. (1995) *Biochemistry*, **34**, 14758–14764.
- Zhang,Q.M., Fujimoto,J. and Yonei,S. (1995) *Int. J. Radiat. Biol.*, **68**, 603–607.
- Steinberg,J.J., Cajigas,A. and Brownlee,M. (1992) *J. Chromatogr.*, **574**, 41–55.
- Cannon,S.V., Cummings,A. and Teebor,G.W. (1988) *Biochem. Biophys. Res. Commun.*, **151**, 1173–1179.