



Title	Redescription of <i>Cyphon patiens</i> Klausnitzer based on the Japanese Specimens
Author(s)	Yoshitomi, Hiroyuki; Hayashi, Masakazu
Description	Biodiversity and Biogeography of the Kuril Islands and Sakhalin vol.3
Citation	北海道大学総合博物館研究報告, 5, 79-81
Issue Date	2009-03
Doc URL	https://hdl.handle.net/2115/47858
Type	departmental bulletin paper
File Information	v. 3-5.pdf



Redescription of *Cyphon patiens* Klausnitzer based on the Japanese Specimens

Hiroyuki Yoshitomi¹ and Masakazu Hayashi²

¹ Ehime University Museum, Bunkyo-chô 3, Matsuyama, 790-8577 Japan

E-mail: hymushi@agr.ehime-u.ac.jp;

² Hoshizaki Green Foundation, Okinoshima 1659-5, Sono, Izumo, 691-0076 Japan

Abstract *Cyphon patiens* Klausnitzer is newly recorded from Japan, with redescription and figures.

Key words: Scirtidae, Coleoptera, *Cyphon patiens*, redescription, Hokkaido

Introduction

Recently the second author found out that two variations of the male genitalia in *Cyphon ainu* Nakane distributed in Hokkaido, Japan. After the careful examination of our collections and type specimens, it become clear that one of them is true *Cyphon ainu* Nakane and another is *Cyphon patiens* Klausnitzer known from Kunashir Islands.

In the present paper, we redescribe *Cyphon patiens* Klausnitzer based on the Japanese specimens.

Methodology and abbreviations were shown in the previous study (Yoshitomi, 2005).

Taxonomy

Cyphon patiens Klausnitzer, 1982

Cyphon patiens Klausnitzer, 1982, 280 [Type: in ZIL, examined].

Cyphon ainu: Yoshitomi, 2005, 103 [a part, misidentification].

[Japanese name: Nise-ainu-chibi-maruhananomi]

Materials examined. Type materials – Holotype, 1 male, Kunashir, Tretiakovo, 29–VI–1973, Kerzhner leg. (written by Russian characters). Misidentification to *Cyphon ainu* in Yoshitomi (2005) – 11 males & 2 females, Sarufutsu-gawa, Sarufutsu-mura, 1–VIII–1996, K. Mizota leg.; 1 male, Shiretoko-rindô, 24–VII–1986, K. Ishida leg.; 1 male, near Shiretoko-tôge, 500–750m alt., Rausu-chô, 26–VII–1994, K. Akita leg.; 1 male, Wakoto, Kucharo, 5–VII–1958, M. Miyatake leg.

Additional materials examined – 1 male & 1 female, Shiranuka, near Kushiro, 11–VII–1992, B. Kuznetsov leg.; 3 males & 1 female, Shiretoko-rindô, Utoro, Shari-chô, 8–VII–1992, B. Kuznetsov leg.; 1 male, Kitayama

camp station, Abashiri, 23–VI–1998, K. Ijima leg.; 1 male, Kabuto-numa, Toyotomi-chô, 28–VI–1999, M. Hayashi leg.; 1 male, Hyôtan-numa, Temninkyô, Higashikawa-chô, 22–VII–1993, M. Hayashi leg.; 2 males, Shitaosobetsu, Shibeche, 28–VI–1995, K. Ijima leg.

Redescription. Male. The external features fully described by Klausnitzer (1982); PW/PL 2.00–2.21 (2.11); EL/EW 1.53–1.85 (1.71); EL/PL 4.83–5.52 (5.17); EW/PW 1.33–1.58 (1.44); TL/EW 1.84–2.20 (2.05). Caudal margin of sternite VII lightly convex. Tergite VIII well sclerotized; median plate forming lobe-like projection, finely punctate; lateral projections short, curved interiorly, obtuse at apices, closely covered with large punctures in apical 1/3. Tergite IX well sclerotized, consisting of a pair of rod-like hemitergite, as long as tergite VIII; apices obtuse, bearing short setae. Tegmen short, well sclerotized; parameres short, curved dorsally, pointed at apices. Penis short, well sclerotized; parameroid forming lobe-like projection, lacking any projections in lateral margin, obtuse at apex.

Female. Sexual dimorphism indistinct. PW/PL 2.13–2.25 (2.19); EL/EW 1.69–1.80 (1.75); EL/PL 5.08–5.54 (5.32); EW/PW 1.32–1.50 (1.39); TL/EW 2.00–2.13 (2.08). Caudal margin of sternite VII arcuate. Tergite VIII moderately sclerotized, trapezoidal, shallowly concave in caudal margin, which bearing short spines, sparsely covered with minute setae in caudal part; apodemes long and slender. Sternite VIII moderately sclerotized, bearing short setae in apical part, rather straight in apical margin. Ovipositor long, two pairs of apical setae in stylus; relative length of stylus, coxite and baculus as (n = 1) 1.0 : 4.1 : 18.3. Prehensor weakly sclerotized, V-shaped.

Measurement. Male (n = 8): TL 3.39–4.08 (3.71)




Figure 1.
Habitus of *Cyphon ainu*
Nakane (A) and *Cyphon*
patiens Klausnitzer (B).

mm; PW 1.15–1.40 (1.27) mm; PL 0.52–0.68 (0.60) mm; EL 2.87–3.40 (3.10) mm; EW 1.75–1.90 (1.81) mm. Female (n = 3): TL 3.60–3.80 (3.70) mm; PW 1.20–1.35 (1.28) mm; PL 0.55–0.61 (0.59) mm; EL 3.05–3.20 (3.12) mm; EW 1.75–1.80 (1.78) mm.

Distribution. Kunashir, Hokkaido (east and northern parts).

Remarks. This species belongs to subgroup A of *Cyphon collaris* species-group (sensu Yoshitomi, 2005). This is closely related to *Cyphon ainu* Nakane distributed sympatrically (see Figure 3), and differs from the latter by the following characteristics: 1) lateral projections of tergite VIII shorter than median plate, closely covered with large punctures in apical 1/3; 2) parameroid of penis forming lobe-like projection; 3) apical margin of sternite VIII of female rather straight; 4) tergite VIII of female shallowly concave in caudal margin.

In the original description of this species (p. 280–282 in Klausnitzer, 1982), the following mistakes were presented: sternite IX (also fig. 23) – lateral projections of tergite VIII; tegmen (also fig. 24) – penis and upside-down; penis (also fig. 25) – tegmen.

Cyphon ainu Nakane, 1963

Cyphon ainu Nakane, 1963, 31 [Type: in SEHU, examined]. Other synonyms see Yoshitomi (2005).

[Japanese name: Ainu-chibi-maruhananomi]

Additional materials examined. 1 male, Iwabokki, Kushiro, 6–VII–1992, B. Kuznetsov leg.; 1 male, Kottaro, Kushiro, 6–VII–1992, B. Kuznetsov leg.; 1 male, Shiraisawa-rindô, Otaru, 1–VII–1992, B. Kuznetsov leg.; 1 male, Kushiro-shitsugen, 4–VII–1992, B. Kuznetsov leg.; 3 males, Kabuto-numa, Toyotomi-chô, 28–VI–1999, M. Hayashi leg.; 33 exs.,

Shibetsu-chô, 26–VI–2001, H. Yoshitomi leg.; 4 males, Bushigawa, Okushiri Is., 6~9–VI–2003, S. Hori leg.

Other materials were recorded by Yoshitomi (2005) except for the specimens of misidentification to *Cyphon patiens* Klausnitzer, 1982 (see above).

Discussion

We can not find the differential characters between *C. ainu* and *C. patiens* in the external features, so we have to check the male and female genitalia.

The following keys are addition to “Key to species of *Cyphon collaris* species-group of Japan” (Yoshitomi, 2005, p. 100–103).

1. Coloration of body almost yellowish-orange throughout; distributed in Hokkaido. 1'
- . Coloration of body almost blackish-brown throughout, but some species have yellowish-orange in pronotum only; distributed in Honshu, Shikoku and Kyushu. 2
- 1'. Lateral projections of tergite VIII of male shorter than median plate, closely covered with large punctures in apical 1/3; parameroid of penis forming lobe-like projection; apical margin of sternite VIII of female rather straight; tergite VIII of female shallowly concave in caudal margin.
..... *C. patiens* Klausnitzer
- . Lateral projections of tergite VIII of male longer than median plate, closely covered with short setae; parameroid of penis expanded laterally in apical part; apical margin of sternite VIII of female arcuate; tergite VIII of female arcuate in caudal margin. *C. ainu* Nakane




Figure 2.

Cyphon patiens Klausnitzer, male (A–D) and female (E–I). A and E: sternites V–VII. B: tergite VIII. C: tergite IX. D: tegmen and penis. F: sternite VIII. G: tergite VIII. H: ovipositor. I: prehensor.




Figure 3.

Distribution map of *Cyphon patiens* Klausnitzer and *Cyphon ainu* Nakane.

Acknowledgements

We wish to express our sincere gratitude to Dr. Nikolai Nikitsky (ZIL), and Dr. Masahiro Ôhara (SEHU) for their permission to loan the museum collections, and to Dr. Andreas Pütz, Mr. Shigehisa Hori, Mr. Nobuki Yasuda, Mr. Kazuo Ijima for their kindness in offering us invaluable specimens used in this study.

References

- KLAUSNITZER, B., 1982. Zur Kenntnis der Helodidae von Sachalin und den Kurilen-Inseln, UdSSR (Coleoptera). 62. Beitrag zur Kenntnis der Helodidae. *Polskie Pismo entomologiczne* 52, 275–285.
- NAKANE, T., 1963. A list of Coleoptera from the Siretoko Peninsula, Hokkaido, Japan (Insecta). *The Scientific Reports of Kyoto Prefectural University (Natural Science & Living Science)* 3 (5), 23–31.
- YOSHITOMI, H., 2005. Systematic revision of the family Scirtidae of Japan, with phylogeny, morphology and bionomics (Insecta: Coleoptera, Scirtoidea). *Japanese Journal of Systematic Entomology*. Monographic Series 3, 1–212.