



HOKKAIDO UNIVERSITY

Title	Versatile Photorearrangement of Photocycloadducts from 5-Fluoro-1,3-dimethyluracil and Naphthalene
Author(s)	Ohkura, Kazue; Akizawa, Hiromichi; Kudo, Mikiko et al.
Citation	Heterocycles, 84(2), 1057-1065 https://doi.org/10.3987/COM-11-S(P)83
Issue Date	2012-01-01
Doc URL	https://hdl.handle.net/2115/49042
Type	journal article
File Information	Het84-2_1057-1065.pdf



VERSATILE PHOTOREARRANGEMENT OF PHOTOCYCLOADDUCTS FROM 5-FLUORO-1,3-DIMETHYLURACIL AND NAPHTHALENE

Kazue Ohkura,^{*,a} Hiromichi Akizawa,^a Mikiko Kudo,^a Tetsuya Ishihara,^a Nobuhiro Oshima,^a and Koh-ichi Seki^{*,b}

^a Faculty of Pharmaceutical Sciences, Health Sciences University of Hokkaido, Ishikari-Tobetsu, Hokkaido 061-0293, Japan

E-mail: ohkura@hoku-iryo-u.ac.jp

^b Central Institute of Radio Isotope Science, Hokkaido University, Kita-15, Nishi-7, Kita-ku, Sapporo 060-0815, Japan

Abstract— Direct UV-irradiation of 5-fluoro-1,3-dimethyluracil (5-FDMU) and naphthalene (**1**) with a 500 W high-pressure mercury lamp in a degassed Pyrex tube ($\lambda > 300$ nm) predominantly afforded benzopyrimidobarrelene derivative (**2**) through 1,4-addition, while irradiation in the presence of piperylene in singlet excited states preferentially afforded naphthocyclobutapyrimidine derivative (**3**) via 1,2-addition. Upon 254 nm light-irradiation of **2** gave rise to the formation of benzopyrimidosemibullvalene (**4**) in fair yields. The reaction pathway for the formation of **4** is reasonably explained in the terms of di- π -methane rearrangement. Adduct **3** was newly converted to the corresponding barrelene derivative (**2**) by long-wave-length irradiation in the presence of a triplet sensitizer.

INTRODUCTION

In recent years, significant attention has been paid to the photocycloaddition of naphthalenes with alkenes as a useful procedure for constructing certain unique ring systems by way of 1,2-,^{1,2} 1,4-,^{1,2} 1,3-,³ 1,8-,⁴ and [4 + 4] additions.^{1,5} We previously reported that UV-irradiation of 5-fluoro-1,3-dimethyluracil (5-FDMU) and naphthalene (**1**) with a 500 W high-pressure mercury lamp in a degassed Pyrex tube ($\lambda > 300$ nm) predominantly afforded the 1,4-adduct, namely benzopyrimidobarrelene derivative (**2**) in high yield.⁶ A time course study of the photoreaction of an acetonitrile solution of 5-FDMU and **1** under the same conditions showed that the 1,2-cycloadduct, naphthocyclobutapyrimidine derivative (**3**) is formed as the major product at the initial stage, which however, is quite labile to the UV-light used, to


revert rapidly back to 5-FDMU and **1**, while the 1,4-adduct (**2**) was produced competitively with **3**, though less effectively but insensitive to the light, to accumulate in the reaction mixture as the irradiation time is prolonged.⁷ The addition of the triplet quencher piperylene to the reaction mixture of **1** and 5-FDMU dramatically changed the mode of cycloaddition, resulting in the preferential formation of **3** through 1,2-addition.⁸ Thus, 1,4- and 1,2- adducts of the pyrimidine ring and naphthalene have been synthesized mode-selectively in fair yields upon long-wave-length irradiation by a high-pressure mercury lamp with a Pyrex filter ($\lambda > 300$ nm). Furthermore, we have obtained 1,3-cycloadducts, semibullvalene derivatives (**4**), by the UV-irradiation with a low-pressure mercury lamp ($\lambda = 254$ nm). The semibullvalenes were found to be secondary products derived from initially formed 1,4-adducts under UV-irradiation, although the precise reaction mechanism remained unclear.⁹ Thus, we have hitherto demonstrated a mode selective synthesis of three types of cycloadducts, 1,2-, 1,3-, *albeit* formally, and 1,4-adducts, by controlling the reaction conditions appropriately.

Although versatile photorearrangements of 1,4-adducts and 1,2-adducts of aromatic hydrocarbons with alkenes have extensively been explored,¹⁰⁻¹² little is known about the photochemical behavior of heteroaromatic-fused barrelenes except pyrazinobarrelenes and benzoquinoxalinobarrelenes.¹³ Hence, our attention was focused to elucidate an aspect of the reciprocal valence isomerization of the adducts of 5-FDMU with naphthalene. In the present paper we describe the feature of the mutual conversion of the cycloadducts between 5-FDMU and **1**.

RESULTS AND DISCUSSION

Photoisomerization of benzopyrimidobarrelene (**2**)

First, we examined photochemical behavior of 1,4-adduct (**2**). Upon UV-irradiation with a high-pressure



Scheme 1

mercury lamp ($\lambda > 300$ nm), starting barrelene **2** was recovered unchanged, while irradiation of **2** in cyclohexane for 5 min with 254 nm light from a 60 W low-pressure mercury lamp gave rise to the formation of semibullvalene (**4**) in fair yields (40%), together with a trace amount of novel dehydrofluorinated semibullvalene derivative (**5**), and cyclo-reversionary products, 5-FDMU and **1**, in 20% yield (Scheme 1).

Time course experiments of the present photolysis with ^1H -NMR spectroscopy showed that prompt formation of **4** occurred first, while the formation of **5** was barely detectable at the initial stage (Figure





Figure 1

1). The formation of **4** reached a maximum at 5 min. With the decrease of **4**, formation of **5** rose. These results indicate that semibullvalene (**4**) may be converted to the dehydrofluorinated derivative (**5**) during the irradiation.


Indeed **4** gave semibullvalene **5** as a major product upon irradiation in cyclohexane with 254 nm light for 30 min (Scheme 2).



Scheme 2

Photoisomerization of naphthocyclobutapyrimidine (3)

We then examined the photochemical behavior of 1,2-adduct (3). Direct irradiation of 3 with a high pressure mercury lamp (10 min) only restored the starting materials, 5-FDMU and 1, in high efficiency (63 %). By contrast, the UV-irradiation of 3 in the presence of a triplet sensitizer, benzophenone, gave rise to the formation of benzopyrimidobarrelene (2) (10%), in competition with cyclo-reversion to the original 1 and 5-FDMU (55%) (Scheme 3).




Scheme 3

The transformation from 1,2-adduct (3) to 1,4-adduct (2) can be explained by a mechanism involving biradical intermediate in the excited triplet state, as proposed by J. Aretz et al.¹⁴


Mechanistic studies for the formation of semibullvarene (4) from benzopyrimidobarrelene (2)

The formation of semibullvarene 4 from 2 can be accounted either in terms of di- π -methane rearrangement¹⁰ (pathway 1, participating atoms/bonds are denoted as a-b-c-d-e in Scheme 4), or by a mechanism involving a biradical intermediate (pathway 2 in Scheme 4). It would be readily clarified by carrying out the photoreaction with 8,12-dimethylbenzopyrimidobarrelene (6), whereby either 11a-fluoro-4a,10,11-trihydro-2,4,7,10-tetramethyl-5,10,11-metheno-5H-benzo[5,6]cyclohepta[1,2-d]pyrimidine-1,3-dione (7) via di- π -methane rearrangement, or 11a-fluoro-4a,10,11-trihydro-2,4,7,10-tetramethyl-5,10,11-metheno-5H-benzo[5,6]cyclohepta [1,2-d]pyrimidine-1,3-dione (8) through a biradical intermediate should be produced.



Scheme 4

UV-Irradiation of **6** with a low-pressure mercury lamp resulted in the formation of **7** (18 %) yield, clearly demonstrating that di- π -methane rearrangement is involved in the present photo-transformation. In this reaction a novel type semibullvalene, 4a-Fluoro-10,11,11a-trihydro-1,3,8,12-tetramethyl-5,10,11-metheno-5*H*- benzo[4,5]cyclohepta[1,2-*d*]pyrimidine-1,3-dione (**9**) was also produced through di- π -methane rearrangement shown in Scheme 5 (participating atoms/bonds are denoted as b-a-f-e-d) (14%). Although methyl groups on barrelene derivative may be influential ~~in~~ on radical reactions, these



Scheme 5

results exhibit significant participation of di- π -methane rearrangement in the reaction pathway for the formation of **4**.

CONCLUSION

The present study clearly demonstrates the reciprocal photochemical valence isomerization of the cycloadducts of 5-FDMU and naphthalene. 1,4-adduct **2** which is intact to a long-wave-length light ($\lambda > 300$ nm), is readily converted into benzopyrimidosemibullvalene (**4**) with a short-wave-length light ($\lambda = 254$ nm). By contrast, 1,2-cycloadduct (**3**) is sensible to the light from a high-pressure mercury lamp ($\lambda > 300$ nm), to restore 5-FDMU and naphthalene through cyclo-reversion, whereas UV-irradiation with a high-pressure mercury lamp in the presence of a triplet sensitizer effected valence isomerization of **3** into **2**. Thus, 1,2-, 1,3-, and 1,4-cycloadducts of 5-FDMU with naphthalene are shown to be produced mode-selectively.

From the viewpoint of photochemistry of heteroaromatic-fused cycloadducts, it is worthy of note ~~attention~~ that adducts **2** and **3** photo-isomerized with the fluoride atom remaining intact at the original position, while photolysis of semibullvalene (**4**) gave pyrimidine-fused-semibullvalene (**5**) via ~~by~~ dehydrofluorination.

EXPERIMENTAL

NMR spectra were measured with a JEOL JNM-EA500 (500 MHz) spectrometer, and ^1H -NMR chemical shifts were given on the δ (ppm) scale based on those of the solvent signals. MS spectra and high-resolution MS (HRMS) spectra were recorded with a LEOL JMS-HX110 (FAB). HPLC was conducted on a Shim-pac PREP-Sil (H) (25 cm x 20 mm *i. d.*) (silica gel), using a LC-10A apparatus (Shimadzu, Kyoto) with monitoring at 254 nm. UV-Irradiation was carried out externally with a 60 W low-pressure lamp (Eiko-sha, Osaka) in a quartz tube under argon ($\lambda = 254$ nm) or a 500 W high-pressure mercury lamp (Eiko-sha, Osaka) in a degassed Pyrex tube ($\lambda > 300$ nm) on a merry-go-round apparatus. Yields were determined using ^1H -NMR spectroscopy with p-dinitrobenzene as an internal standard. Benzopyrimidobarrelene (**2**) was synthesized by the photoreaction of 5-FDMU and **1** in cyclohexane solution with a high-pressure mercury lamp ($\lambda > 300$ nm).⁶ Naphthocyclobutapyrimidine (**3**) was prepared by the similar photoreaction of 5-FDMU and **1** in the presence of piperylene.⁷

Photolysis of 2 in cyclohexane----- A solution (0.051 mmol) of **2** in cyclohexane (70 mL) was put

portion-wise (10 mL each) into 7 quartz tubes, and irradiated externally with a 60 W low-pressure lamp under an argon atmosphere at room temperature. The reaction mixture was concentrated *in vacuo*, and the residual oil was subjected to HPLC with 30% ethyl acetate in hexane.

11a-Fluoro-4a,10,11-trihydro-2,4-dimethyl-5,10,11-metheno-5H-benzo[5,6]cyclohepta[1,2-d]pyrimidine-1,3-dione(4): ¹H-NMR(methanol-*d*₄) δ:2.47(3H, s, N²-CH₃), 2.48(1H, ddd, *J*= 6.3, 8.0, 18.3Hz, H-11), 2.74 (1H, dd, *J*= 6.3, 8.0 Hz, H-10), 3.07 (1H, ddd, *J*= 6.3, 6.3, 6.3 Hz, H-12), 3.14 (3H, s, N⁴-CH₃), 4.07 (1H,ddd, *J*= 1.7, 6.3, 6.9 Hz, H-5), 4.44 (1H,dd, *J*= 6.9, 30.4 Hz, H-4a), 6.85 (1H, d, *J*= 7.5 Hz, H-6), 7.02 (1H, ddd, *J*= 1.2, 7.5, 7.5 Hz, H-7), 7.12 (1H, ddd, *J*= 1.2, 7.5, 7.5 Hz, H-8), 7.22 (1H, d, *J*=7.5 Hz, H-9). NOE: H-11with H-10 (19.7%), H-12 (3.6%); H-10 with H-12 (3.9%), H-9 (1.9%); H-12 with H-5 (3.6%), H-10 (2.4%); N⁴-CH₃ with H-4a (2.3%), H-5(1.3%), H-6(1.1%); H-5 with H-4a (4.6%), H-6 (2.0%), H-12 (4.3%), N⁴- CH₃ (2.0%). FAB-MS (*m/z*): [M + H]⁺ 287. HRFAB-MS (*m/z*): [M + H]⁺ Calcd for C₁₆H₁₆FN₂O₂, 287.1196; Found, 287.1204.

10,11-Dihydro-2,4-dimethyl-5,10,11-metheno-5H-benzo[5,6]cyclohepta[1,2-d]pyrimidine-1,3-dione (5):¹H-NMR(acetone-*d*₆) δ:3.03 (1H, t, *J*= 6.9Hz, H-11), 3.08 (3H, s, N²-CH₃), 3.15 (1H, dd, *J*= 6.3, 6.9Hz, H-10), 3.44 (1H, ddd, *J*= 6.3, 6.9, 6.9Hz, H-12) ,3.58 (3H, s, N⁴-CH₃), 4.71 (1H, d, *J*= 6.9 Hz, H-5), 7.11 (1H, dt, *J*= 1.2,7.5 Hz, H-7), 7.18 (1H, dt, *J*= 1.2, 7.5 Hz, H-8), 7.38 (1H, d, *J*= 7.5Hz, H-6), 7.40 (1H, d, *J*= 7.5Hz, H-9). NOE: H-11 with H-10 (1.0%), H-12 (1.8%). FAB-MS (*m/z*): [M + H]⁺ 267.HRFAB-MS (*m/z*): [M + H]⁺ Calcd for C₁₆H₁₅N₂O₂, 267.1133; Found, 267.1129.

Photolysis of 4 in cyclohexane----- A solution (0.034 mmol) of **4** in cyclohexane (70 mL) was UV irradiated under the same conditions described above to afford **5** (28%) together with unreacted **4** (37%).

Photolysis of 6 in cyclohexane----- A solution (0.027 mmol) of **2** in cyclohexane (50 mL) was put portion-wise (10 mL each) into 5 quartz tubes, and irradiated externally with a 60 W low-pressure lamp under an argon atmosphere for 5 min at room temperature. The reaction mixture was concentrated *in vacuo*, and the residual oil was subjected to HPLC with 1% ethyl acetate in CH₂Cl₂ to give **7**(18%) and **9** (14%), together with unreacted **6** (26%).

11a-Fluoro-4a,10,11-trihydro-2,4,7,10-tetramethyl-5,10,11-metheno-5H-benzo[5,6]cyclohepta[1,2-d]pyrimidine-1,3-dione (7): ¹H-NMR(CDCl₃) δ:1.54 (3H, s, C¹⁰-CH₃), 2.25 (3H, s, C⁷-CH₃), 2.38 (1H, dd, *J*= 6.3, 18.9Hz, H-11), 2.53 (3H, s, N²-CH₃), 2.72 (1H, dd, *J*= 5.2, 6.3 Hz, H-12), 3.14 (3H, s, N⁴-CH₃), 3.87(1H, dd, *J*= 5.2, 6.3 Hz, H-5), 4.16 (1H, dd, *J*= 6.3, 29.8 Hz, H-4a), 6.60 (1H, s, H-6), 6.98 (1H, d, *J*= 7.5 Hz, H-8), 7.08 (1H, d, *J*= 8.0Hz, H-9). NOE: C¹⁰-CH₃ with H-11(1.6%), H-12 (1.0%), H-9 (1.4%); H-4a with N⁴-CH₃ (5.3%), H-5 (6.3%), H-12 (1.3%); H-5 with H-12 (5.6%), N⁴-CH₃(2.8%),

H-4a (7.1%), H-6 (3.1%); N⁴-CH₃ with H-5 (1.4%), H-4a (2.4%), H-6 (1.6%). FAB-MS (*m/z*): [M + H]⁺ 315. HRFAB-MS (*m/z*): [M + H]⁺ Calcd for C₁₈H₂₀FN₂O₂, 315.1508; Found, 315.1510.

4a-Fluoro-10,11,11a-trihydro-1,3,8,12--tetramethyl-5,10,11-metheno-5*H*-benzo[4,5]cyclohepta[1,2-*d*]pyrimidine-1,3-dione (9): ¹H-NMR(CDCl₃) δ : 1.67 (3H, s, C¹²-CH₃), 2.13 (1H, dd, *J*= 6.3, 7.5 Hz, H-11) 2.26 (3H, s, C⁸-CH₃), 2.41 (1H, d, *J*=7.5 Hz, H-10), 2.74 (3H, s, N¹-CH₃), 2.95 (3H, s, N³-CH₃), 3.81(1H, dd, *J*= 2.9, 12.6 Hz, H-5), 4.28 (1H, ddd, *J*= 2.9, 6.3, 26.9 Hz, H-11a), 6.68 (1H, d, *J*= 7.5Hz, H-6), 6.81(1H, d, *J*=7.5Hz, H-7), 7.04 (1H, s, H-9). NOE: C¹²-CH₃ with H-11 (1.6%), H-10 (1.6%), H-5 (2.3%); H-11 with C¹²-CH₃ (2.4%), H-10 (7.3%), N¹-CH₃ (2.8%), H-11a (8.8%); C⁸-CH₃ with H-7 (2.1%), H-9 (2.2%); H-10 with C¹²-CH₃(1.6%), H-11(7.3%), N¹-CH₃ (1.2%), H-9 (3.7%); H-5 with C¹²-CH₃ (2.4%), H-6 (3.2%); H-11a with H-11(7.4%), N¹-CH₃(3.5%); H-6 with N³-CH₃ (1.6%), H-5 (3.9%), H-7 (7.7%); H-9 with C⁸-CH₃ (4.6%), H-10 (3.8%), N¹-CH₃ (1.4%). FAB-MS (*m/z*): [M + H]⁺ 315. HRFAB-MS (*m/z*): [M + H]⁺ Calcd for C₁₈H₂₀FN₂O₂, 315.1508; Found, 315.1512.

Photosensitization of 3 ----- An acetonitrile solution of **3** (0.025 M) was irradiated in the presence of benzophenone (0. 83 M) with a 500 W high-pressure mercury lamp in a degassed Pyrex tube to give **2** (10%) together with 5-FDMU (55%).

REFERENCES

1. J. J. Maccullough, *Chem. Rev.*, **1987**, 811.
2. A. Kugerberg, D. Döpp, and H. Görner, *J. Inf. Recording*, 2000, **25**, 187.
3. N. Zuparcic and B. Sket, *J. Chem. Soc., Perkin Trans. 1*, 1992, 179.
4. Y. Kubo, T. Inoue, and H. Sakai, *J. Am. Chem. Soc.*, 1992, **114**, 7660.
5. S. M. Sieburth, K F. Jr. McGee, F. Zhang, and Y. Chen, *J. Org. Chem.*, 2000, **65**, 1972; K. Chiyonobu, G. Konishi, Y. Inoue, and K. Mizuno, *J. Chem. Res.*, 2001, 135.
6. K. Ohkura, T. Sugaoi, K. Nishijima, Y. Kuge, and K. Seki, *Tetrahedron Lett.*, 2002, **43**, 3113.
7. K. Ohkura, T. Sugaoi, T. Ishihara, and K. Seki, *Heterocycles*, 2004, **64**, 57.
8. K. Ohkura, T. Ishihara, K. Nishijima, J. Diakur, and K. Seki, *Chem. Pharm. Bull.*, 2005, **53**, 258.
9. K. Ohkura, M. Kudo, T. Ishihara, K. Nishijima and K. Seki, *Heterocycles*, 2005, **65**, 2583.
10. H. E. Zimmerman, The di- π -methane (Zimmerman) rearrangement, in *Rearrangements in Ground and Excited States*, vol. 3, ed. by P. de Mayo, Academic Press, New York, 1980, pp131-166; H. E. Zimmerman, The di- π -methane rearrangement, in *Organic Photochemistry*, vol. 11, ed. by A. Padwa, Mercel Dekker, New York, 1991, pp1-36; H. E. Zimmerman and D. Armesto, *Chem. Rev.*,

- 1996, **96**, 3065; S. S.Hixson, P. S. Mariano, and H. E. Zimmerman, *Chem. Rev.*, 1973, **73**, 531; H. E. Zimmerman, R. S.Givens, and R. M.Pagni, *J. Am. Chem. Soc.*, 1968, **90**, 6096; J. Luo, H. Ihmels, H. Deiseroth, and M. Schlosser, *Can. J. Chem.*, 2009, **87**, 619.
11. J. R. Scheffer, and J. Yang, The photochemistry of dibenzobarrelenes (9,10-ethenoanthracene) and related and its derivatives, in *Handbook of Photochemistry and Photobiology*, eds. by W. Horspool and P. –S. Song, CRC press, Boca Raton, FL, 1995, chap. 16; C. O. Bender and D. W. Brooks, *Can J. Chem.*, 1975, **53**, 1684; D. Ramaiah, M. C. Sajimon, J. Joseph, and M. V. George, *Chem. Soc. Rev.*, 2005, **34**, 48.
 12. H.-D. Scharf and J. Mattay, *Tetrahedron Lett.*, 1977, **5**, 401; J. Grota and J. Mattay, *Photochem. Photobiol. Sci.*, 2005, **4**, 625.
 13. C. –C. Liao and R. K. Peddinti, M. Oelgemoller, and A. G. Griesbeck, ‘Photochemistry of Heteroarene-Fused Barrelenes, in *CRC Handbook of Organic Photochemistry and Photobiology*, 2nd Edition, eds. by W. Horspool and F.Lenci, CRC press, Boca Raton, 2004, pp.32/1-32/17.
 14. H.-D. Scharf, H. Leismann, W. Erb, H. W. Gaidetzka, and J. Aretz, *Pure Appl. Chem.*, 1975, **41**, 581.