



# HOKKAIDO UNIVERSITY

|                  |                                                                                                                                        |
|------------------|----------------------------------------------------------------------------------------------------------------------------------------|
| Title            | Adult Leigh Disease Without Failure to Thrive                                                                                          |
| Author(s)        | Sakushima, Ken; Tsuji-Akimoto, Sachiko; Niino, Masaaki et al.                                                                          |
| Citation         | Neurologist, 17(4), 222-227<br><a href="https://doi.org/10.1097/NRL.0b013e318217357a">https://doi.org/10.1097/NRL.0b013e318217357a</a> |
| Issue Date       | 2011-07                                                                                                                                |
| Doc URL          | <a href="https://hdl.handle.net/2115/49478">https://hdl.handle.net/2115/49478</a>                                                      |
| Rights           | This is a non-final version of an article published in final form in Neurologist, Jul. 2011, 17(4), 222-227.                           |
| Type             | journal article                                                                                                                        |
| File Information | Neu17-4_222-227.pdf                                                                                                                    |



## Case report

### Title:

Adult Leigh disease without failure to thrive : a case report and review of the literatures

### Authors:

Ken Sakushima, M.D., M.P.H., <sup>1</sup> Sachiko Tsuji-Akimoto, M.D., Ph. D., <sup>1</sup> Masaaki Niino, M.D.,  
Ph.D., <sup>1</sup> Shinji Saitoh, M.D., Ph.D., <sup>2</sup> Ichiro Yabe, M.D., Ph.D., <sup>1</sup> and Hidenao Sasaki, M.D.,  
Ph. D. <sup>1</sup>

1. *Department of Neurology, Hokkaido University Graduate School of Medicine, Hokkaido, Japan.*
2. *Department of Pediatrics, Hokkaido University Graduate School of Medicine, Hokkaido, Japan.*

### Correspondence:

Ichiro Yabe, M.D., Ph.D.

Department of Neurology, Hokkaido University Graduate School of Medicine, Kita-15,

Nishi-7, Kita-ku, Sapporo 060-8638 Japan

e-mail; yabe@med.hokudai.ac.jp

### short title:

Adult Leigh disease

### Disclosure of funding:

None

### Conflict of interest:

None

## **Abstract**

### **Introduction:**

Most Leigh disease (LD) patients die before reaching adulthood, but there are reports of 'adult LD'. The clinical features of adult LD were quite different from those in infant or childhood cases. Here, we describe a normally developed patient with adult LD, who presented with spastic paraplegia that was followed several years later by acute encephalopathy. We also conducted a systemic literature search on adult LD and integrated its various manifestations to arrive at a diagnostic procedure for adult LD.

### **Case report:**

A 26-year-old woman presented with acute encephalopathy after spastic paraplegia. On her first admission, she exhibited bilateral basal ganglia lesion on magnetic resonance images and normal serum lactate levels. On second admission, she had acute encephalopathy with lactic acidosis and bilateral basal ganglia and brainstem lesions. A muscle biopsy revealed cytochrome c oxidase(COX) deficiency, and a diagnosis of adult LD was made. Despite treatment in the intensive care unit, she died nine days after admission.

### **Conclusions:**

A review of the literature describing adult LD revealed that developmental delay, COX deficiency, serum lactate elevation, and basal ganglia lesions occurred less frequently than they did in children with LD. Cranial nerve disturbance, pyramidal signs, and cerebellar dysfunction were the primary symptoms in adult LD. Thus, the many differences between childhood and adult LD may be helpful for diagnosing adult LD.

### **Key Words:**

Leigh disease, adult, mitochondrial deficits, Rahman's criteria

## 1)Introduction

Leigh disease (LD) is a metabolic disorder of the central nervous system usually affecting infants or children. The characteristic features of LD are developmental delay, lactic acidosis, and typical lesions in the brainstem and basal ganglia<sup>1</sup>. LD is commonly caused by deficits in mitochondrial function, such as those involving respiratory chain complexes, ATPase 6, and components of the pyruvate dehydrogenase complex<sup>2</sup>. Typical pathologic features consist of spongiform lesions, microglial and histiocytic proliferation, and vascular proliferation in the basal ganglia and brainstem. Stringent diagnostic criteria for LD in children were proposed by Rahman et al.<sup>3</sup>, which included (1) progressive neurological disease with motor and intellectual developmental delay; (2) signs and symptoms of brainstem and/or basal ganglia disease; (3) raised lactate levels in blood and/or cerebrospinal fluid(CSF); and (4) one or more of the following: (a) characteristic features of neuroradioimaging (symmetrical hypodensities in the basal ganglia on computed tomography [CT] or hyperintense lesions on T2-weighted magnetic resonance imaging [MRI]), (b) typical neuropathological changes at postmortem, or (c) typical neuropathology in a similarly affected sibling. In addition, they defined patients with a high probability of LD who did not fulfill the criteria as “Leigh-like” syndrome.

Most LD patients die before reaching adulthood, but there are reports of ‘adult LD’<sup>4-28</sup>. The clinical features of adult LD were quite different from those in infant or childhood cases<sup>29, 30</sup>. Here, we describe a normally developed patient with adult LD, who presented with spastic paraplegia that was followed several years later by acute encephalopathy. Furthermore, we conducted a systematic literature search on adult LD and integrated its various manifestations

to arrive at a diagnostic procedure for adult LD.

## **2) Case Report**

A 26-year-old Japanese female was admitted to our hospital because of diarrhea and fever followed by coma. She was 162 cm in height on admission. In the past, she had asthma, a meniscal tear, chronic alveolar osteitis, and a periapical cyst. She had no family history of LD or mitochondrial diseases. She was born without any complications and had developed without mental retardation. At age 13, she was affected with spastic paraplegia five days after an asthmatic attack. At 18, she acquired a sensory disturbance of the lower extremities. At 24, she was hospitalized for a diagnostic workup of hearing loss after an asthmatic attack(1<sup>st</sup> admission). Neurological examination at that time revealed right facial palsy, sensorineural hearing loss, spastic paraplegia, hyperreflexia of the lower extremities, bilateral extensor planter response, and sensory disturbance in the lower extremities. In a laboratory test, her serum lactate level was within normal limits. Brain CT and MRI showed small bilateral lesions of the basal ganglia (Fig. 1). At age 26, she developed dysarthria and gait disturbance, and subsequently diarrhea and high fever. Three days after the onset of fever, a disturbance of consciousness with lactic acidosis had progressed and she was transferred to the intensive care unit (ICU) (2<sup>nd</sup> admission). She had deep coma, generalized hyporeflexia, and loss of the oculocephalic reflexes. Neither anisocoria nor neck stiffness were observed. She was treated with steroids, multivitamins, and coenzyme Q10 (COQ10) in the ICU. Brain CT showed diffuse cerebral swelling and edema with bilateral hypodensity in the basal ganglia and brainstem (Fig. 2). Laboratory findings ( Table 1) revealed elevations of creatinine kinase

(CK), and amylase as well as lactic acidosis. CSF examination showed high CSF pressure and mild pleocytosis that did not indicate any infectious disease. A muscle biopsy revealed cytochrome c oxidase (COX) deficiency without inflammation that indicated a mitochondrial abnormality, but we did not observe strongly succinate dehydrogenase-reactive blood vessels (SSV) and ragged red fiber (RRF). She died nine days after admission. No autopsy was performed. Later, a genetic analysis of the whole mitochondrial genome and known mutations of nuclear genes (*SURF1*, *COX10*) were performed, but no mutations were detected.

### **3) Discussion**

#### **3-1. Systematic literature review**

##### **Search methods and results of the systematic literature review**

Adult LD was defined as patients who survived longer than 18 years<sup>2</sup>. We searched reports in English on Medline from 1966 to October 2009 using the MESH terms "Leigh disease" and "adult", on EMBASE from 1968 to October 2009 using "Leigh disease AND adult", and the bibliographies of the retrieved articles. We applied two concepts for the accurate diagnosis of LD: (1) the patient fulfilled all four items of Rahman's criteria for LD, or three of the four items with neuroradiological or pathological abnormalities for Leigh-like syndrome (Rahman's criteria group: RCG) and (2) the patient had an identified mitochondrial abnormality in a muscle biopsy or other procedures or gene mutations (laboratory-diagnosed group: LDG). In all, 40 reports describing 58 patients were identified (Table 2); of these 19 belong to the RCG, 24 to the LDG, and 12 patients fulfilled both sets of criteria. In the RCG, 4

patients fulfilled all 4 items of Rahman's LD criteria and 15 of them fulfilled 3 of the 4 Leigh-like items.

### **Characteristics and clinical manifestation**

The RCG included 13 males and 6 females. The mean age of onset was  $10.4 \pm 10.9$  years (mean  $\pm$  SD). At the time of the reports, 14 of the patients were alive. Nine patients had a familial history of LD. The LDG included 12 males and 12 females. The mean age of onset was  $12.7 \pm 14.8$  years. At the time of the reports, 18 of these patients were alive. Sixteen patients in the LDG had a familial history of LD. The proportions of each symptom in the two groups are summarized in Fig. 3-a. In both groups, the major symptoms reported were cerebellar dysfunction and lower (VII-XII) cranial nerve disturbances. In the LDG, pyramidal signs were more frequent and mental retardation was less frequent than in the RCG..

On the other hand, in children with LD, the major symptoms are mental retardation, poor feeding, swallowing difficulty, and respiratory disturbance. Ophthalmoplegia and ataxia were reported in only 35% and 31% of children with LD, respectively<sup>29, 30</sup>.

### **Biochemical and neuroradiological features**

The percentages of adult patients with elevated lactate and neuroradiological abnormalities are shown in Fig. 3-b. In laboratory examinations, elevations in lactate levels were more frequently detected in serum than in CSF; however, serum or CSF lactate levels were not described in some patients, and others had normal serum lactate levels during the uneventful phase. As the resting levels of serum lactate were often within the normal range , the lactate

stress test (LST) that measures increases in serum lactate during slight exercise may be useful for detecting elevated lactate levels in the uneventful phase<sup>31, 32</sup>. In the reports that we reviewed, the LST was not performed or described.

Neuroradiological findings of basal ganglia lesions on MRI or CT were reported in only 67% of the LDG patients. On the other hand, brainstem lesions were more frequent in the LDG than the RCG. In contrast to the relatively low proportion of neuroradiological abnormalities reported in these cases, such abnormalities were present in most children with LD<sup>29, 33, 34</sup>.

### **Muscle biopsy histology, enzymatic deficiencies, and mutations**

Muscle biopsies were performed in 18 of the reported adult cases. Pathological studies comprised light microscopic assessment of frozen sections using hematoxylin and eosin (H&E), modified Gomori trichrome, succinate dehydrogenase (SDH), and COX staining. A COX deficiency was observed in only 4 patients and subsarcolemmal aggregations of abnormal mitochondria (Ragged-red fiber: RRF) were observed in 5 patients. In enzymatic analyses of muscle specimens which mainly evaluated the oxidation capacity of NADH, COX, succinate, and pyruvate dehydrogenase complex (PDHc) using standard methods<sup>35, 36</sup>, decreased respiratory chain activity (NADH, COX, and succinate) was observed in 3 patients and decreased PDHc activity was observed in another.

Mutations were detected using direct nucleotide sequence of causative genes including the mitochondrial genome or those identified by positional cloning<sup>2</sup>. The genetic analyses indicated that 18 patients had one or more gene mutations: the T8993C mutation was detected

in 4 patients from different families, 1 family had 4 cases of adult LD with the T9185C mutation, 3 patients in different families had a T8993G mutation, and 2 patients in the same family a G1644T mutation. There were also single occurrences of heterozygous 14459, heterozygous 14484, homozygous 3460, G13513A, and C7028T+G9547A.

COX deficiency is a typical abnormality in child LD patients with the *SURF1* gene mutation<sup>37</sup>, but RRF is absent in most childhood cases<sup>30, 38</sup>. Other reported means of evaluating mitochondrial dysfunction are to analyze enzyme activities in skin fibroblasts, muscle supernatants, lymphocytes, and the liver<sup>3</sup>. The major mutations known to occur in LD patients are T8993C, T8993G, T10191C, G13513A, A8344G, and A3243G in mitochondrial genes, and *SURF1* in the nuclear genome. Currently, there are 24 known mutations in mitochondrial genes and 21 in nuclear genes<sup>2</sup>.

### **Differential diagnosis and treatment**

Most of the reported cases excluded infections, autoimmune diseases, and toxins. Some cases were first misdiagnosed as multiple sclerosis and other cases were indistinguishable from Wernicke's encephalopathy. From the perspective of the basal ganglia lesions, hypoxic encephalopathy, Wilson's disease, mitochondrial myopathy, encephalopathy, lactic acidosis, stroke-like episodes (MELAS), and neurofibromatosis type1 should be considered as differential diagnoses.

In some cases, thiamin was administered as a nonspecific treatment. Disease specific treatments consisted of Coenzyme Q10 (150-300mg/day) administered to four patients with mitochondrial abnormalities, three of which improved and the other was unchanged, and a

ketogenic diet was administered to 2 patients; both of which improved.

For children and adults, thiamin could be available for the initial treatment for LD, but its efficacy is not as well established as either coenzyme Q10 or a ketogenic diet. Currently, there is no established treatment for LD<sup>2</sup>.

### **3-2. Conclusion**

In our case, in spite of the presence of bilateral basal ganglia lesion, a diagnosis of LD was quite difficult, due to the normal lactate levels and spastic paraplegia, both are considered as atypical symptoms for LD, and the lack of gene mutations. Adult LD could not be diagnosed until a histochemical examination of the biopsy specimen revealed the COX deficiency without inflammation that indicated a mitochondrial abnormality. Using Rahman's criteria, our patient could be consistent with a "Leigh-like" syndrome. A systematic literature review revealed that adult LD was rare and its clinical manifestations were different from those of children. These differences indicate that additional investigations should be considered when LD is suspected in adults. Together with our systematic literature review, we assembled the following characteristic features of adult LD for early diagnosis:

1. History of cryptogenic thrive failure or signs of mental retardation, pyramidal signs, cerebellar disturbances, ophthalmoplegia, deafness, dysarthria, or other neurological symptoms are present.
2. Bilateral basal ganglia lesions or brainstem lesions with serum or CSF lactate elevation are present. (LST should be considered when resting lactate levels are normal .)
3. Mitochondrial abnormalities are present in muscle pathology or in biochemical analyses, or

known LD gene mutations are present.

4. Metabolic disorders, toxins, infection, multiple sclerosis, and Wernicke's encephalopathy can be excluded.

Early diagnosis and further investigations of treatments for adult LD are expected to improve its prognosis.

## Figure legends

### Figure 1

- a) Brain CT (left) and T2WI MRI (right) on the first admission. Brain CT showed bilateral low density areas (arrow) in the basal ganglia. MRI showed bilateral and slightly asymmetric T2 high intensity areas (arrow) in the putamen.
- b) Brain CT on the second admission. Brain CT showed diffuse parenchymal edema accompanied by typical bilateral low density areas (arrow) in the putamen and cerebellum. The putaminal lesions were larger than those seen at the first admission.
- c) Histochemical study of the muscle biopsy specimens stained for succinate dehydrogenase (left) and cytochrome c oxidase (center), and a positive control for cytochrome c oxidase (right). Succinate dehydrogenase-reactive blood vessels were not observed. Cytochrome c oxidase was completely deficient

### Figure 2

- a) Clinical symptoms and signs observed in the reported cases of adult LD
- b) Laboratory data and neuroradiological abnormalities observed in reported cases of adult LD. Bars represent the percentages of total patients expressing the various symptoms and signs. RCG: Rahman's criteria group, LDG: laboratory diagnosed group.

## References

1. Leigh D. Subacute necrotizing encephalomyelopathy in an infant. *J Neurol Neurosurg Psychiatry* 1951;14:216-21.
2. Finsterer J. Leigh and Leigh-like syndrome in children and adults. *Pediatr Neurol* 2008;39:223-35.
3. Rahman S, Blok RB, Dahl HH, et al. Leigh syndrome: clinical features and biochemical and DNA abnormalities. *Ann Neurol* 1996;39:343-51.
4. Feigin I, Goebel HH. "Infantile" subacute necrotizing encephalopathy in the adult. *Neurology* 1969;19:749-59.
5. Sipe JC. Leigh's syndrome: the adult form of subacute necrotizing encephalomyelopathy with predilection for the brainstem. *Neurology* 1973;23:1030-8.
6. Feigin I, Budzilovich GN. Further observations on subacute necrotizing encephalomyelopathy in adults. *J Neuropathol Exp Neurol* 1977;36:128-39.
7. Ulrich J, Fankhauser-Mauri C. Subacute necrotizing encephalopathy (Leigh) in an adult. *Eur Neurol* 1978;17:241-6.
8. Whetsell WO, Jr., Plaitakis A. Leigh's disease in an adult with evidence of "inhibitor factor" in family members. *Ann Neurol* 1978;3:519-24.
9. Ho KL, Piligian JT, Chason JL. Adult form of subacute necrotizing encephalomyelopathy. *Arch Pathol Lab Med* 1979;103:344-7.
10. Kalimo H, Lundberg PO, Olsson Y. Familial subacute necrotizing encephalomyelopathy of the adult form (adult Leigh syndrome). *Ann Neurol* 1979;6:200-6.
11. Gray F, Louarn F, Gherardi R, Eizenbaum JF, Marsault C. Adult form of Leigh's disease: a clinico pathological case with CT scan examination. *J Neurol Neurosurg Psychiatry* 1984;47:1211-5.
12. Maso E, Ferrer I, Herraiz J, Roquer J, Serrano S. Leigh's syndrome in an adult. *J Neurol* 1984;231:253-7.
13. Bourgeois M, Duret A, Vital C, Assens F, Mazerat P. [Leigh disease in an adult. A case with psychiatric symptoms]. *Ann Med Psychol (Paris)* 1986;144:485-94.
14. Roquer J, Herraiz J. [Leigh syndrome in the adult]. *Arch Neurobiol (Madr)* 1986;49:129-35.
15. Bianco F, Floris R, Pozzessere G, Rizzo PA. Subacute necrotizing encephalomyelopathy (Leigh's disease): clinical correlations with computerized tomography in the diagnosis of the juvenile and adult forms. *Acta Neurol Scand* 1987;75:214-7.
16. Cavanagh JB. Necrotising hemorrhagic encephalopathy in an adult: Leigh's disease? *J Neurol Neurosurg Psychiatry* 1987;50:1245-6.
17. Delgado G, Gallego J, Tunon T, Zarranz JJ, Villanueva JA. Necrotising haemorrhagic encephalomyelopathy in an adult: Leigh's disease. *J Neurol Neurosurg Psychiatry* 1987;50:224-7.
18. Kissel JT, Kolkin S, Chakeres D, Boesel C, Weiss K. Magnetic resonance imaging in a case of autopsy-proved adult subacute necrotizing encephalomyelopathy (Leigh's disease). *Arch Neurol* 1987;44:563-6.
19. Fulham M, Lawrence C, Harper C. Diagnostic clues in an adult case of Leigh's disease. *Med J Aust* 1988;149:320-2.
20. Reynaud P, Loiseau H, Coquet M, Vital C, Loiseau P. [An adult case of Leigh's subacute necrotizing encephalomyelopathy]. *Rev Neurol (Paris)* 1988;144:259-65.
21. Halmagyi GM, Pampflett R, Curthoys IS. Seesaw nystagmus and ocular tilt reaction due to adult Leigh's disease. *Neuro-ophthalmology* 1992;12:1-9.
22. Spranger M, Schwab S, Wiebel M, Becker CM. [Adult Leigh syndrome. A rare differential diagnosis of central respiratory insufficiency]. *Nervenarzt* 1995;66:144-9.
23. Chalmers RM, Lamont PJ, Nelson I, et al. A mitochondrial DNA tRNA(Val) point mutation associated with adult-onset Leigh syndrome. *Neurology* 1997;49:589-92.
24. Kumagai R, Ichikawa K, Yasui T, Kageyama Y, Miyabayashi S. Adult leigh syndrome: treatment with intravenous soybean oil for acute central respiratory failure. *Eur J Neurol* 1999;6:613-5.
25. Nagashima T, Mori M, Katayama K, et al. Adult Leigh syndrome with mitochondrial DNA mutation at 8993. *Acta Neuropathol* 1999;97:416-22.
26. Malojcic B, Brinar V, Poser C, Djakovic V. An adult case of Leigh disease. *Clin Neurol Neurosurg* 2004;106:237-40.
27. Piao YS, Tang GC, Yang H, Lu DH. Clinico-neuropathological study of a Chinese case of familial adult Leigh syndrome. *Neuropathology* 2006;26:218-21.
28. Lekha P, Gayatri N, Lathika S, Sree K. Adult onset Leigh syndrome. *Annals of Indian Academy of Neurology* 2007;10:55-7.

- 29.Lee HF, Tsai CR, Chi CS, Lee HJ, Chen CC. Leigh syndrome: clinical and neuroimaging follow-up. *Pediatr Neurol* 2009;40:88-93.
- 30.Yang YL, Sun F, Zhang Y, et al. Clinical and laboratory survey of 65 Chinese patients with Leigh syndrome. *Chin Med J (Engl)* 2006;119:373-7.
- 31.Finsterer J, Milvay E. Lactate stress testing in 155 patients with mitochondriopathy. *Can J Neurol Sci* 2002;29:49-53.
- 32.Finsterer J. Reproducibility of the lactate stress test. *Metab Brain Dis* 2003;18:155-60.
- 33.Goldenberg PC, Steiner RD, Merkens LS, et al. Remarkable improvement in adult Leigh syndrome with partial cytochrome c oxidase deficiency. *Neurology* 2003;60:865-8.
- 34.Arii J, Tanabe Y. Leigh syndrome: serial MR imaging and clinical follow-up. *AJNR Am J Neuroradiol* 2000;21:1502-9.
- 35.Fischer JC, Ruitenbeek W, Berden JA, et al. Differential investigation of the capacity of succinate oxidation in human skeletal muscle. *Clin Chim Acta* 1985;153:23-36.
- 36.Fischer JC, Ruitenbeek W, Trijbels JM, et al. Estimation of NADH oxidation in human skeletal muscle mitochondria. *Clin Chim Acta* 1986;155:263-73.
- 37.Pronicki M, Matyja E, Piekutowska-Abramczuk D, et al. Light and electron microscopy characteristics of the muscle of patients with SURF1 gene mutations associated with Leigh disease. *J Clin Pathol* 2008;61:460-6.
- 38.Shtilbans A, Shanske S, Goodman S, et al. G8363A mutation in the mitochondrial DNA transfer ribonucleic acidLys gene: another cause of Leigh syndrome. *J Child Neurol* 2000;15:759-61.

**Table 1. Laboratory data of our case**

| <b>Hematologic study</b> |                              | <b>Coagulation study</b>              |                         |
|--------------------------|------------------------------|---------------------------------------|-------------------------|
| WBC                      | 19,400/ $\mu$ l              | PT                                    | 12.7 seconds            |
| RBC                      | $520 \times 10^4$ / $\mu$ l  | APTT                                  | 34.0 seconds            |
| Hb                       | 15.4 g/dl                    | Fibrinogen                            | 502 mg/dl               |
| Plt                      | $35.6 \times 10^4$ / $\mu$ l | FDP                                   | 4.0 $\mu$ g/dl          |
| <b>Chemical study</b>    |                              | D-dimer                               | 2.53 $\mu$ g/dl         |
| TP                       | 6.3 g/dl                     | <b>Cerebrospinal fluid analysis</b>   |                         |
| T-Bil                    | 0.3 mg/dl                    | CSF pressure                          | >400 mmH <sub>2</sub> O |
| GOT                      | 39 IU/l                      | Cell                                  | 10/ $\mu$ l             |
| GPT                      | 18 IU/l                      | Protein                               | 60 mg/dl                |
| LDH                      | 370 IU/l                     | Glucose                               | 89 mg/dl                |
| CK                       | 2053 IU/l                    | <b>Muscle pathology</b>               |                         |
| Amylase                  | 998 IU/l                     | SSV                                   | negative                |
| BUN                      | 32 mg/dl                     | COX                                   | complete deficiency     |
| Cr                       | 1.7 mg/dl                    | <b>Gene analysis</b>                  |                         |
| Na                       | 150 mEq/l                    | whole mitochondrial genom             | normal                  |
| K                        | 3.9 mEq/l                    | <i>SURF1</i> and <i>COX10</i> mutatic | negative                |
| Cl                       | 102 mEq/l                    |                                       |                         |
| Ca                       | 7.9 mEq/l                    |                                       |                         |
| Lactate                  | 134.4 mg/dl                  |                                       |                         |
| Pyruvate                 | 9.46 mg/dl                   |                                       |                         |
| NH <sub>3</sub>          | 55 $\mu$ g/dl                |                                       |                         |

Table 2. List of searched reports


| Source                   | Year | Age | Alive | FH  | Sex | Onset | MA  | Mutation | Autopsy | Group(items*) |
|--------------------------|------|-----|-------|-----|-----|-------|-----|----------|---------|---------------|
| Feigin_1 <sup>4</sup>    | 1969 | 36  | No    | No  | M   | 0     | No  | ND       | Yes     | RCG           |
| Feigin_2 <sup>4</sup>    | 1969 | 30  | No    | No  | F   | 28    | No  | ND       | Yes     | No(2)         |
| Sipe <sup>5</sup>        | 1973 | 32  | No    | No  | F   | 5     | No  | ND       | Yes     | No(2)         |
| Feigin_1 <sup>6</sup>    | 1977 | 50  | No    | No  | M   | 49    | No  | ND       | Yes     | No(1)         |
| Feigin_2 <sup>6</sup>    | 1977 | 84  | No    | No  | F   | 84    | No  | ND       | Yes     | No(1)         |
| Feigin_3 <sup>6</sup>    | 1977 | 66  | No    | No  | M   | 58    | No  | ND       | Yes     | No(1)         |
| Feigin_4 <sup>6</sup>    | 1977 | 58  | No    | No  | M   | 58    | No  | ND       | Yes     | No(1)         |
| Ulrich <sup>7</sup>      | 1978 | 31  | No    | No  | M   | 31    | No  | ND       | Yes     | No(2)         |
| Whetsell <sup>8</sup>    | 1978 | 21  | No    | Yes | M   | 6     | No  | ND       | Yes     | RCG           |
| Kalimo_1 <sup>10</sup>   | 1979 | 62  | No    | Yes | F   | 22    | No  | ND       | Yes     | No(1)         |
| Kalimo_2 <sup>10</sup>   | 1979 | 46  | No    | Yes | M   | 13    | No  | ND       | Yes     | No(2)         |
| Kalimo_3 <sup>10</sup>   | 1979 | 43  | No    | Yes | M   | 15    | No  | ND       | Yes     | No(1)         |
| Ho <sup>9</sup>          | 1979 | 55  | No    | No  | F   | 52    | No  | ND       | Yes     | No(2)         |
| Gray <sup>11</sup>       | 1984 | 31  | No    | No  | F   | 21    | No  | ND       | Yes     | No(2)         |
| Maso <sup>12</sup>       | 1984 | 55  | No    | No  | M   | 55    | No  | ND       | Yes     | No(2)         |
| Erven_1 <sup>37</sup>    | 1987 | 25  | Yes   | Yes | M   | 3     | No  | ND       | No      | No(3)         |
| Erven_2 <sup>37</sup>    | 1987 | 23  | Yes   | Yes | F   | 9     | No  | ND       | No      | No(2)         |
| Erven_3 <sup>37</sup>    | 1987 | 20  | Yes   | Yes | F   | 4     | No  | ND       | No      | No(1)         |
| Cummiskey <sup>38</sup>  | 1987 | 34  | No    | No  | F   | ND    | No  | ND       | Yes     | No(2)         |
| Kissel <sup>18</sup>     | 1987 | 21  | No    | Yes | F   | 21    | No  | ND       | Yes     | No(2)         |
| Delgado <sup>17</sup>    | 1987 | 21  | No    | No  | M   | 21    | No  | ND       | Yes     | No(2)         |
| Bianco_1 <sup>15</sup>   | 1987 | 19  | Yes   | No  | M   | 6     | No  | ND       | No      | RCG           |
| Bianco_2 <sup>15</sup>   | 1987 | 27  | Yes   | No  | M   | 12    | No  | ND       | No      | RCG           |
| Fulham <sup>19</sup>     | 1988 | 32  | No    | No  | F   | 32    | No  | ND       | Yes     | No(2)         |
| Peiffer <sup>39</sup>    | 1988 | 34  | No    | Yes | M   | 9     | Yes | ND       | Yes     | RCG           |
| Halmagyi <sup>21</sup>   | 1992 | 31  | No    | No  | M   | 29    | No  | ND       | Yes     | No(2)         |
| deVries <sup>40</sup>    | 1993 | 56  | No    | Yes | F   | 56    | No  | T8993C   | No      | LDG           |
| Santorelli <sup>41</sup> | 1996 | 23  | Yes   | No  | F   | 12    | Yes | ND       | No      | LDG           |
| Morris <sup>42</sup>     | 1996 | 24  | No    | No  | M   | 6     | Yes | ND       | Yes     | RCG, LDG      |
| Chalmers_1 <sup>23</sup> | 1997 | 43  | Yes   | Yes | F   | 25    | Yes | G1644T   | No      | RCG, LDG      |
| Chalmers_2 <sup>23</sup> | 1997 | 38  | Yes   | Yes | M   | 20    | Yes | G1644T   | No      | RCG, LDG      |
| Suzuki <sup>43</sup>     | 1998 | 20  | No    | Yes | F   | 1     | No  | T8993C   | No      | LDG           |
| Malandrini <sup>44</sup> | 1998 | 32  | No    | No  | M   | 7     | No  | ND       | Yes     | No(2)         |

|                               |      |    |     |     |   |    |     |                   |     |          |
|-------------------------------|------|----|-----|-----|---|----|-----|-------------------|-----|----------|
| Kumagai <sup>24</sup>         | 1999 | 38 | Yes | No  | F | 18 | Yes | ND                | No  | RCG, LDG |
| Nagashima <sup>25</sup>       | 1999 | 43 | No  | No  | F | 37 | No  | T8993G            | Yes | RCG, LDG |
| Vilarinho <sup>45</sup>       | 2001 | 21 | Yes | Yes | F | 0  | No  | T8993C            | No  | RCG, LDG |
| Yamakawa <sup>46</sup>        | 2001 | 22 | Yes | No  | M | 3  | Yes | ND                | No  | RCG      |
| Funalog_1 <sup>47</sup>       | 2002 | 33 | No  | Yes | M | 4  | No  | hetro14484        | No  | LDG      |
| Funalog_2 <sup>47</sup>       | 2002 | 32 | No  | No  | M | 18 | Yes | hetro14459        | No  | LDG      |
| Funalog_3 <sup>47</sup>       | 2002 | 39 | Yes | Yes | M | 34 | No  | homo3460          | No  | LDG      |
| Maldergem_1 <sup>48</sup>     | 2002 | 29 | Yes | Yes | F | 0  | Yes | ND                | No  | RCG, LDG |
| Maldergem_2 <sup>48</sup>     | 2002 | 31 | Yes | Yes | F | 0  | Yes | ND                | No  | LDG      |
| Kirby <sup>49</sup>           | 2003 | 36 | Yes | Yes | M | ND | No  | G13513A           | No  | LDG      |
| Yoshinaga <sup>50</sup>       | 2003 | 20 | Yes | No  | F | 0  | No  | T8993G            | No  | LDG      |
| Goldenberg <sup>33</sup>      | 2003 | 22 | Yes | Yes | F | 14 | No  | C7028T<br>+G9547A | No  | RCG, LDG |
| Malojcic <sup>26</sup>        | 2004 | 21 | Yes | No  | M | ND | Yes | ND                | No  | RCG      |
| Piao <sup>27</sup>            | 2006 | 32 | No  | Yes | M | 17 | No  | ND                | Yes | No(2)    |
| Wick <sup>51</sup>            | 2007 | 21 | No  | Yes | F | 21 | No  | ND                | Yes | No(1)    |
| Child_1 <sup>52</sup>         | 2007 | 24 | Yes | Yes | M | 6  | No  | T9185C            | No  | LDG      |
| Child_2 <sup>52</sup>         | 2007 | 22 | Yes | Yes | M | 3  | No  | T9185C            | No  | LDG      |
| Child_3 <sup>52</sup>         | 2007 | 18 | Yes | Yes | M | 1  | No  | T9185C            | No  | RCG, LDG |
| Child_4 <sup>52</sup>         | 2007 | 24 | Yes | Yes | F | 8  | No  | T9185C            | No  | LDG      |
| Debray <sup>53</sup>          | 2007 | 18 | Yes | No  | M | 4  | No  | T8993C            | No  | RCG, LDG |
| Lekha <sup>28</sup>           | 2007 | 27 | Yes | No  | M | 26 | Yes | ND                | No  | RCG, LDG |
| Mkaouar-Rebai_1 <sup>54</sup> | 2009 | ND | ND  | No  | M | 8  | ND  | ND                | No  | No(2)    |
| Mkaouar-Rebai_2 <sup>54</sup> | 2009 | ND | ND  | Yes | M | 7  | ND  | ND                | No  | No(2)    |
| Mkaouar-Rebai_3 <sup>54</sup> | 2009 | ND | ND  | Yes | M | ND | ND  | ND                | No  | No(2)    |
| Sobreira <sup>55</sup>        | 2009 | 42 | Yes | Yes | M | 0  | No  | T8993G            | No  | RCG, LDG |
| Present case                  |      | 26 | No  | No  | F | 13 | Yes | ND                | No  | RCG, LDG |


\* Number of items fulfilling the Rahman's criteria, RCG: Rahman's criteria group, LDG: laboratory diagnosed group, ND: not determined, MA: mitochondrial abnormality demonstrated by muscle biopsy or other procedures, "Yes": mitochondrial abnormality exists, "No": mitochondrial abnormality was not determined or examined, FH: family history

Fig.1

a)




b)



c)



# Figure 2-a



# Figure 2-b

