



HOKKAIDO UNIVERSITY

Title	Chromatographic Rules in the Elution of Individual Molecular Species of Triglyceride
Author(s)	Takahashi, Koretaro; Hirano, Tsugihiko; Zama, Kozo
Description	Short Paper
Citation	日本水産学会誌, 49(8), 1301-1301 https://doi.org/10.2331/suisan.49.1301
Issue Date	1983
Doc URL	https://hdl.handle.net/2115/49852
Rights	© 1983 公益社団法人日本水産学会
Type	journal article
File Information	BJSSF49_1301.pdf



Short Paper

Chromatographic Rules in Elution of Individual Molecular Species of Triglyceride^{*1}


We have discovered previously that there are two rules in the elution of diglyceride acetate derived from lecithin on high performance liquid chromatography¹⁾, that is:

$$C = p_1 \cdot \log RRT + q_1 \quad C = \begin{vmatrix} x & d_1 \\ c_2 & d_2 \end{vmatrix} \quad (1)$$

$$D = p_2 \cdot \log RRT + q_2 \quad D = \begin{vmatrix} c_1 & y \\ c_2 & d_2 \end{vmatrix} \quad (2)$$

where p is the slope and q is the intersection on the ordinate. c and d are acyl carbon number and number of double bonds in each acyl group, respectively. x and y are variables of acyl carbon number and number of double bonds, respectively.

The structure of diglyceride acetate is:



And the structure of triglyceride is:



The difference between (3) and (4) is at position 3 in the molecule. But $\text{CO} \cdot \text{CH}_3$ can also be considered as the shortest chain in acyl group.

So: $\begin{array}{l} \text{—Acyl}_1 \\ \text{—Acyl}_2 \end{array} \ni \begin{array}{l} \text{—Acyl}_1 \\ \text{—Acyl}_2 \\ \text{—CO} \cdot \text{CH}_3 \end{array}$ (where $\ni$ means "include")

therefore, (1) and (2) can be rewritten as:

$$C = p_1 \cdot \log RRT + q_1 \quad C = \begin{vmatrix} x & d_1 \\ c_2 & d_2 \\ 2 & 0 \end{vmatrix} \quad (1')$$

$$D = p_2 \cdot \log RRT + q_2 \quad D = \begin{vmatrix} c_1 & y \\ c_2 & d_2 \\ 2 & 0 \end{vmatrix} \quad (2')$$

since (2, 0) in position 3 exhibits the acyl (acetyl) group which has two carbons and no double bond. So we can conclude that (1) and (2) are members of the general

rule for triglycerides under the condition of $\begin{vmatrix} x & d_1 \\ c_2 & d_2 \\ 2 & 0 \end{vmatrix}$ or

$\begin{vmatrix} c_1 & y \\ c_2 & d_2 \\ 2 & 0 \end{vmatrix}$. By adapting the general expressions to triglyceride, the following equations will finally be obtained.

$$C = p_1 \cdot \log RRT + q_1 \quad C = \begin{vmatrix} x & d_1 \\ c_2 & d_2 \\ c_3 & d_3 \end{vmatrix} \quad (5)$$

$$D = p_2 \cdot \log RRT + q_2 \quad D = \begin{vmatrix} c_1 & y \\ c_2 & d_2 \\ c_3 & d_3 \end{vmatrix} \quad (6)$$

We have adapted (5) and (6) to the data of PERKINS *et al.*^{2,3)} and discovered that the theory, that has proven mathematically, is available practically as illustrated in the two figures.

Accepted November 4, 1982

^{*1} クロマトグラフィーにおける各トリグリセリド分子種の溶出順位性について

^{*2} 高橋是太郎・座間宏一: 北海道大学水産学部・函館市港町.

^{*3} 平野次彦: 日立北海セミコンダクター株式会社・北海道七飯町.




Fig. 1. Relation between relative retention time of triglyceride types and their total acyl carbon number.

* Triolein was used as a reference peak for the relative retention time calculation.




Fig. 2. Relation between relative retention time of triglyceride types and their total double bond.

* Triolein was used as a reference peak for the relative retention time calculation.

The authors wish to thank Associate Professor Kōzō TAKAMA of Training Factory for Food Processing of our Faculty for his helpful advice.

Koretarō TAKAHASHI,^{*2} Tsugihiko HIRANO,^{*3} and Kōichi ZAMA^{*2}

Faculty of Fisheries, Hokkaido University, Minato, Hakodate 041, Japan.

References

- 1) K. TAKAHASHI, T. HIRANO, K. TAKAMA, and K. ZAMA: *Bull. Japan. Soc. Sci. Fish.*, **48**, 1803-1814 (1982).
- 2) A. H. EL-HAMDY and E. G. PERKINS: *JAACS*, **58**, 867-872 (1982).
- 3) E. G. PERKINS, D. J. HENDREN, N. PELICK, and J. E. BAUER: *Lipids*, **17**, 460-463 (1982).