



Title	A New Loop-Mediated Isothermal Amplification Method for Rapid, Simple, and Sensitive Detection of <i>Leptospira</i> spp. in Urine
Author(s)	Koizumi, Nobuo; Nakajima, Chie; Harunari, Tsunehito et al.
Citation	Journal of Clinical Microbiology, 50(6), 2072-2074 <a href="https://doi.org/10.1128/JCM.00481-12">https://doi.org/10.1128/JCM.00481-12</a>
Issue Date	2012-06
Doc URL	<a href="https://hdl.handle.net/2115/50782">https://hdl.handle.net/2115/50782</a>
Rights	© 2012 American Society for Microbiology
Type	journal article
File Information	Supplemental_material.pdf, Supplemental material



## 1 **Supplemental Methods**

### 2 **Bacterial DNA preparation**

3 To evaluate the specificity of *Lepto-rrs* LAMP, genomic DNA was extracted from bacterial  
4 cultures using the DNeasy Blood & Tissue Kit (Qiagen, Hilden, Germany) according to the  
5 manufacturer's instructions. The concentration of the extracted DNA was determined using an  
6 ND-1000 spectrophotometer (NanoDrop Technologies, Wilmington, DE, USA). Genomic DNA  
7 was serially diluted for the sensitivity test, and the number of genome copies in the *Lepto-rrs*  
8 LAMP reaction mixture was calculated based on a molecular size of 5.1 Mbp (*L. interrogans*  
9 Fiocruz L1-130) (1).

### 10 **Primer design**

11 The *rrs* sequences from pathogenic, intermediate, and non-pathogenic *Leptospira* spp.  
12 (accession nos. AB279549, AE016823, AY631876, AY631877, AY631878, AY631879,  
13 AY631880, AY631881, AY631883, AY631884, AY631885, AY631886, AY631895, AY796065,  
14 EF025496, EF612284, and Z21634) and other bacteria belonging to the family Leptospiraceae,  
15 e.g., *Leptonema illini* (AY714984) and *Turneriella parva* (AY293856), were aligned and  
16 compared using GENETYX software ver. 9 (Genetyx Cooperation, Tokyo, Japan). *Lepto-rrs*  
17 LAMP primers were designed using PrimerExplorer V4 software (available online at  
18 <https://primerexplorer.jp/lamp4.0.0/index.html>) and manually modified. The location of the

19 primers in *rrs* is shown in Fig. S1. The primer set was designed to amplify a region of *rrs* from  
20 both pathogenic and intermediate *Leptospira* spp but not from non-pathogenic ones. Nucleotides  
21 at the 5' terminus of FIP and 3' terminus of BIP primers were mismatched for *rrs* sequences of  
22 non-pathogenic spp., thereby allowing amplification of only pathogenic and intermediate  
23 *Leptospira* spp. (Fig. S1).

24

25 Reference

26

27 **1. Nascimento, A. L. T. O., S. Verjovski-Almeida, M. A. Van Sluys, C. B. Monteiro-Vitorello,**  
28 **L. E. Camargo, L. A. Digiampietri, R. A. Harstkeerl, P. L. Ho, M. V. Marques, M. C.**  
29 **Oliveira, J. C. Setubal, D. A. Haake, and E. A. L. Martins.** 2004. Genome features of  
30 *Leptospira interrogans* serovar Copenhageni. Braz. J. Med. Biol. Res. **37**:459–477.

31






Fig. S2. Sensitivity test using purified genomic DNA from *L. interrogans*. Lepto-*rrs* LAMP reactions were detected by UV fluorescence (upper panel) or by confirming a ladder-like pattern on 2% agarose gel (lower panel). M, DNA marker (OneSTEP Ladder 50; Nippon Gene, Tokyo, Japan).




Fig. S3. Phylogenetic tree based on the *Leptospira flaB* gene sequence. The sequences obtained in this study were indicated in bold type. An accession number of each sequence is indicated in parentheses. The sequences were aligned in MEGA4 (1) using CLUSTALW, and phylogenetic distances were calculated in MEGA4 using the neighbor-joining method. Numbers on nodes are bootstrap support after 1000 replicates.

Reference

1. Tamura, K., J. Dudley, M. Nei, and S. Kumar. 2007. MEGA4: Molecular evolutionary genetics analysis (MEGA) software version 4.0. *Mol. Biol. Evol.* **24**:1596–1599.