



HOKKAIDO UNIVERSITY

Title	Abel-Tauber theorems for Hankel and Fourier transforms
Author(s)	Kikuchi, Hideyuki; 菊地, 秀行
Degree Grantor	北海道大学
Degree Name	博士(理学)
Dissertation Number	甲第4600号
Issue Date	1999-03-25
DOI	https://doi.org/10.11501/3151282
Doc URL	https://hdl.handle.net/2115/51577
Type	doctoral thesis
File Information	000000336472.pdf



Abel-Tauber theorems for Hankel and
Fourier transforms

Hideyuki Kikuchi

1999

①

学 位 論 文

Abel-Tauber theorems for Hankel and
Fourier transforms

Hideyuki Kikuchi

1999

Abel-Tauber theorems for Hankel and Fourier transforms

HIDEYUKI KIKUCHI

ABSTRACT. We prove Abel-Tauber theorems for Hankel and Fourier transforms. For example, let f be a locally integrable function on $[0, \infty)$ which is eventually decreasing to zero at infinity. Let $\rho = 3, 5, 7, \dots$ and ℓ be slowly varying at infinity. We characterize the asymptotic behavior $f(t) \sim \ell(t)t^{-\rho}$ as $t \rightarrow \infty$ in terms of the Fourier cosine transform of f . Similar results for sine and Hankel transforms are also obtained. As an application, we can give an answer to a problem of R. P. Boas on Fourier series.

1. Introduction and results

As a prototype, we use Fourier cosine transforms to explain our problem. Let f be a locally integrable, eventually decreasing function on $[0, \infty)$ which tends to zero at infinity, and let F_c be its Fourier cosine transform. Let $\rho > 0$ and ℓ be slowly varying at infinity (see below). We are concerned with Abel-Tauber theorems which characterize the asymptotic behavior $f(t) \sim \ell(t)t^{-\rho}$ as $t \rightarrow \infty$ in terms of F_c . It turns out that the values $1, 3, 5, \dots$ of ρ are exceptional. For $\rho \neq 1, 3, 5, \dots$, one can obtain the desired Abel-Tauber theorems using regular variation — or Karamata theory. See Bingham-Goldie-Teugels [BGT, Ch. 4], where references to earlier work by Hardy and Rogosinski, Aljančić, Bojanić and Tomić, Vuilleumier, Zygmund and others are given. However the same theorems do not hold for $\rho = 1, 3, 5, \dots$. These exceptional values are related to the power series expansion of the kernel $\cos x$ (see Soni-Soni [SS]).

1991 *Mathematics Subject Classification*. Primary 40E05; Secondary 42A16, 44A15.

Key words and phrases. Abel-Tauber theorems, Hankel transforms, Fourier transforms, Fourier series, Π -variation.

In [I1], one of the authors showed that one could use Π -variation — or de Haan theory in the terminology of [BGT] — to obtain the desired Abel-Tauber theorem for cosine transforms when $\rho = 1$. For theorems of the same type, we refer to [I1] (cosine series and integrals), [I2] (sine series and integrals), [I3] (Fourier-Stieltjes coefficients), and Bingham-Inoue [BI] (Hankel transforms).

In this paper, we consider the remaining exceptional values, e.g., $\rho = 3, 5, \dots$ for cosine transforms. In fact, as in [BI], we consider those for Hankel transforms from the beginning; the results for cosine and sine transforms follow as special cases. As an application, we can give an answer to a problem of R. P. Boas on Fourier series.

We write R_0 for the class of slowly varying functions at infinity, that is, the class of positive measurable ℓ , defined on some neighbourhood of infinity, satisfying

$$\ell(\lambda x)/\ell(x) \rightarrow 1 \quad (x \rightarrow \infty) \quad \forall \lambda > 0.$$

For $\ell \in R_0$, the class Π_ℓ is the class of measurable f satisfying

$$\{f(\lambda x) - f(x)\}/\ell(x) \rightarrow c \log \lambda \quad (x \rightarrow \infty) \quad \forall \lambda > 0$$

for some constant c , called the ℓ -index of f . See [BGT] for background.

Let $\nu \geq -1/2$, $t^{\nu+1/2}h(t) \in L^1_{\text{loc}}[0, \infty)$, and h be ultimately decreasing to zero at infinity. We consider the *Hankel Transform*

$$H_\nu(x) := \int_0^{\infty-} h(t)(xt)^{1/2} J_\nu(xt) dt \quad (0 < x < \infty), \quad (1.1)$$

where $\int_0^{\infty-}$ denotes an improper integral $\lim_{M \rightarrow \infty} \int_0^M$ and J_ν is the Bessel function

$$J_\nu(x) = \sum_{j=0}^{\infty} c_{\nu,j} x^{\nu+2j} \quad (0 \leq x < \infty)$$

with

$$c_{\nu,j} := \frac{(-1)^j}{2^{\nu+2j} \cdot j! \cdot \Gamma(\nu + j + 1)} \quad (\nu \geq -1/2, j = 0, 1, \dots). \quad (1.2)$$

Since the improper integral on the right of (1.1) converges uniformly on each (a, ∞) with $a > 0$, H_ν is finite and continuous on $(0, \infty)$.

For $n \in \mathbb{N}$ and $x \in (0, \infty)$, we define $\bar{H}_{\nu,n}$ by

$$\bar{H}_{\nu,n}(x) := x^{\nu+\frac{1}{2}+2n} \left\{ H_\nu(1/x) - \sum_{j=0}^{n-1} c_{\nu,j} \int_0^\infty t^{\nu+\frac{1}{2}+2j} h(t) dt \cdot x^{-\nu-\frac{1}{2}-2j} \right\} \quad (1.3)$$

if $\int_0^\infty t^{\nu-\frac{3}{2}+2n} h(t) dt < \infty$.

Theorem 1. Let $\ell \in R_0$ and $n \in \mathbb{N}$. Let $\nu \geq -1/2$, $t^{\nu+\frac{1}{2}} h(t) \in L^1_{\text{loc}}[0, \infty)$, and h be ultimately decreasing to zero at infinity, with Hankel transform H_ν . Then

$$h(t) \sim t^{-\nu-\frac{3}{2}-2n} \ell(t) \quad (t \rightarrow \infty) \quad (1.4)$$

if and only if

$$\int_0^\infty t^{\nu-\frac{3}{2}+2n} h(t) dt < \infty \quad \text{and} \quad \bar{H}_{\nu,n} \in \Pi_\ell \quad \text{with } \ell\text{-index } c_{\nu,n}. \quad (1.5)$$

Note that Theorem 1 includes results for Fourier cosine and sine transforms, as

$$x^{1/2} J_{-1/2}(x) = \sqrt{\frac{2}{\pi}} \cos x, \quad x^{1/2} J_{1/2}(x) = \sqrt{\frac{2}{\pi}} \sin x.$$

For $x \in (0, \infty)$, we define \bar{H}_ν by

$$\bar{H}_\nu(x) := x^{\nu+\frac{1}{2}} H_\nu(1/x). \quad (1.6)$$

We will prove Theorem 1 by reducing the problem to the following known result (which corresponds to the case $n = 0$ of (1.4)):

Theorem A ([BI], extending [I1], [I2]). Let ν , h , H_ν and ℓ be as in Theorem 1.

Then

$$h(t) \sim t^{-\nu-\frac{3}{2}}\ell(t) \quad (t \rightarrow \infty) \quad (1.7)$$

if and only if

$$\bar{H}_\nu \in \Pi_\ell \text{ with } \ell\text{-index } c_{\nu,0}. \quad (1.8)$$

The cosine case $\nu = -\frac{1}{2}$ of Theorem A is due to [I1], the sine case $\nu = \frac{1}{2}$ to [I2], and the general case $\nu \geq -\frac{1}{2}$ to Bingham-Inoue [BI].

The theorems above treat the boundary cases to the following known Abel-Tauber theorem for Hankel transforms:

Theorem B ([RS], [SS], extending [P], [B]). Let ν , h , H_ν and ℓ be as in Theorem 1.

(1) For $0 < \rho < \nu + \frac{3}{2}$,

$$h(t) \sim t^{-\rho}\ell(t) \quad (t \rightarrow \infty) \quad (1.9)$$

if and only if

$$H_\nu(x) \sim x^{\rho-1}\ell(1/x) \cdot 2^{\frac{1}{2}-\rho} \frac{\Gamma(\frac{3}{4} + \frac{\nu}{2} - \frac{\rho}{2})}{\Gamma(\frac{1}{4} + \frac{\nu}{2} + \frac{\rho}{2})} \quad (x \rightarrow 0+). \quad (1.10)$$

(2) Let $n \in \mathbb{N}$ and $\nu - \frac{1}{2} + 2n < \rho < \nu + \frac{3}{2} + 2n$. Then (1.9) holds if and only

if $\int_0^\infty t^{\nu-\frac{3}{2}+2n}h(t)dt < \infty$ and

$$\begin{aligned} H_\nu(x) &= \sum_{j=0}^{n-1} c_{\nu,j} \int_0^\infty t^{\nu+\frac{1}{2}+2j}h(t)dt \cdot x^{\nu+\frac{1}{2}+2j} \\ &\sim x^{\rho-1}\ell(1/x) \cdot 2^{\frac{1}{2}-\rho} \frac{\Gamma(\frac{3}{4} + \frac{\nu}{2} - \frac{\rho}{2})}{\Gamma(\frac{1}{4} + \frac{\nu}{2} + \frac{\rho}{2})} \quad (x \rightarrow 0+). \end{aligned} \quad (1.11)$$

The part (1) of Theorem B is due to Pitman [P], Bingham [B], and Ridenhour-Soni [RS], while the part (2) to Soni-Soni [SS].

We focus on Fourier (cosine and sine) transforms. Let $f \in L^1_{\text{loc}}[0, \infty)$ and f be ultimately decreasing to zero at infinity. We write F_c for the *Fourier cosine transform* of f :

$$F_c(x) = \int_0^{\infty-} f(t) \cos(xt) dt \quad (0 < x < \infty). \quad (1.12)$$

Similarly, let $g(t) \in L^1_{\text{loc}}[0, \infty)$, and g be ultimately decreasing to zero at infinity.

We write G_s for the *Fourier sine transform* of g :

$$G_s(x) = \int_0^{\infty-} g(t) \sin(xt) dt \quad (0 < x < \infty). \quad (1.13)$$

Now, at least formally,

$$F_c^{(2j)}(0) = (-1)^j \int_0^{\infty} t^{2j} f(t) dt, \quad G_s^{(2j+1)}(0) = (-1)^j \int_0^{\infty} t^{2j+1} g(t) dt.$$

So for $n \in \mathbb{N}$ we define $\bar{F}_{c,n}$ by

$$\bar{F}_{c,n}(X) := x^{2n} \left\{ F_c(1/x) - \sum_{j=0}^{n-1} \frac{F_c^{(2j)}(0)}{(2j)!} x^{-2j} \right\} \quad (0 < x < \infty) \quad (1.14)$$

if $F_c \in C^{2n-2}([0, \infty))$. Similarly, for $n \in \mathbb{N}$, we define $\bar{G}_{s,n}$ by

$$\bar{G}_{s,n}(x) := x^{2n+1} \left\{ G_s(1/x) - \sum_{j=0}^{n-1} \frac{G_s^{(2j+1)}(0)}{(2j+1)!} x^{-2j-1} \right\} \quad (0 < x < \infty) \quad (1.15)$$

if $G_s \in C^{2n-1}([0, \infty))$. Here as usual, $C^m([0, \infty))$ is the class of functions which are of $C^m(I)$ -class for some open neighbourhood I of $[0, \infty)$.

Theorem 2. Let $\ell \in R_0$ and $n \in \mathbb{N}$. Let $f \in L^1_{\text{loc}}[0, \infty)$ and f be ultimately decreasing to zero at infinity, with Fourier cosine transform F_c . Then

$$f(t) \sim t^{-2n-1} \ell(t) \quad (t \rightarrow \infty) \quad (1.16)$$

if and only if

$$F_c \in C^{2n-2}([0, \infty)) \quad \text{and} \quad \bar{F}_{c,n} \in \Pi_\ell \quad \text{with } \ell\text{-index } \frac{(-1)^n}{(2n)!}. \quad (1.17)$$

Theorem 3. Let $\ell \in R_0$ and $n \in \mathbb{N}$. Let $g(t)t \in L^1_{\text{loc}}[0, \infty)$ and g be ultimately decreasing to zero at infinity, with Fourier sine transform G_s . Then

$$g(t) \sim t^{-2n-2}\ell(t) \quad (t \rightarrow \infty) \quad (1.18)$$

if and only if

$$G_s \in C^{2n-1}([0, \infty)) \quad \text{and} \quad \bar{G}_{s,n} \in \Pi_\ell \quad \text{with } \ell\text{-index } \frac{(-1)^n}{(2n+1)!}. \quad (1.19)$$

Remark. In Theorem 2, $F_c \in C^{2n-2}([0, \infty))$ implies that the limit $F_c(0+)$ exists and that F_c , with $F_c(0) := F_c(0+)$, is in $C^{2n-2}([0, \infty))$; similarly for the meaning of $G_s \in C^{2n-1}([0, \infty))$ in Theorem 3.

The proofs of Theorems 2 and 3 are based on Theorem 1.

We give an application of Theorem 3 to probability theory. Let X be a real random variable defined on a probability space (Ω, \mathcal{F}, P) . The *tail-sum* of X is the function T defined by

$$T(x) := P(X \leq -x) + P(X > x) \quad (0 \leq x < \infty).$$

Note that T is finite and decreases to zero at infinity. Now

$$\{1 - U(\xi)\}/\xi = \int_0^{\infty-} T(x) \sin(x\xi) dx \quad (0 < \xi < \infty),$$

where U is the real part of the characteristic function of X :

$$U(\xi) := E[\cos(\xi X)] \quad (\xi \in \mathbb{R})$$

(see [BGT, p. 336]). By Theorem 3, the asymptotic behavior

$$T(x) \sim x^{-2n-2} \ell(x) \quad (x \rightarrow \infty)$$

with $n \in \mathbb{N}$ and $\ell \in R_0$ is characterized in terms of U .

We can apply Theorems 1 and A to Question 7.19 of Boas [Bo]. For $f \in L^1[0, \pi]$, we define its *Fourier cosine coefficients* a_n by

$$a_n := \frac{2}{\pi} \int_0^\pi f(t) \cos(nt) dt \quad (n = 1, 2, \dots), \quad := \frac{1}{\pi} \int_0^\pi f(t) dt \quad (n = 0). \quad (1.20)$$

Similarly, for $g \in L^1[0, \pi]$, we define its *Fourier sine coefficients* b_n by

$$b_n := \frac{2}{\pi} \int_0^\pi g(t) \sin(nt) dt \quad (n = 1, 2, \dots). \quad (1.21)$$

Theorem 4. Let $f \in L^1[0, \pi]$ with Fourier cosine coefficients (a_k) . We assume that $a_k \geq 0$ for all $k \geq 0$. Let $n \in \mathbb{N}$ and $\ell \in R_0$. Then

$$\sum_{k=m}^{\infty} a_k \sim \frac{\ell(m)}{m^{2n}} \cdot \frac{1}{2n} \quad (m \rightarrow \infty) \quad (1.22)$$

if and only if

$$f \in C^{2n-2}([0, \pi]) \quad \text{and} \quad \bar{f}_n \in \Pi_\ell \quad \text{with } \ell\text{-index } \frac{(-1)^n}{(2n)!}, \quad (1.23)$$

where

$$\bar{f}_n(x) := x^{2n} \left\{ f(1/x) - \sum_{j=0}^{n-1} \frac{f^{(2j)}(0)}{(2j)!} x^{-2j} \right\} \quad (1/\pi \leq x < \infty). \quad (1.24)$$

Corollary. In Theorem 4, we further assume that (a_k) is decreasing. Then (1.23)

is equivalent to

$$a_m \sim \frac{\ell(m)}{m^{2n+1}} \quad (m \rightarrow \infty). \quad (1.25)$$

Theorem 5. Let $g \in L^1[0, \pi]$ with Fourier sine coefficients (b_k) . We assume that $b_k \geq 0$ for all $k \geq 1$. Let $n \in \mathbb{N}$ and $\ell \in R_0$. Then

$$\sum_{k=m}^{\infty} b_k \sim \frac{\ell(m)}{m^{2n+1}} \cdot \frac{1}{2n+1} \quad (m \rightarrow \infty) \quad (1.26)$$

if and only if

$$g \in C^{2n-1}([0, \pi]) \quad \text{and} \quad \bar{g}_n \in \Pi_\ell \quad \text{with } \ell\text{-index } \frac{(-1)^n}{(2n+1)!}, \quad (1.27)$$

where

$$\bar{g}_n(x) := x^{2n+1} \left\{ g(1/x) - \sum_{j=0}^{n-1} \frac{g^{(2j+1)}(0)}{(2j+1)!} x^{-2j-1} \right\} \quad (1/\pi \leq x < \infty). \quad (1.28)$$

Corollary. In Theorem 5, we further assume that (b_k) is decreasing. Then (1.27) is equivalent to

$$b_m \sim \frac{\ell(m)}{m^{2n+2}} \quad (m \rightarrow \infty). \quad (1.29)$$

Remark. We understand that $L^1[0, \pi]$ consists of equivalence classes with respect to the equivalence relation $f_1 \sim f_2 \Leftrightarrow f_1 = f_2$ a.e. So, e.g., in (1.23), $f \in C^{2n-2}([0, \pi])$ implies that there exists a function in $C^{2n-2}([0, \pi])$ which lies in the equivalence class of f and that we identify the function with f . In particular, if $\sum_{k=0}^{\infty} |a_k| < \infty$, then by [Z, Ch. III, Theorem 3.9] (Theorem of Lebesgue on Cesàro summability) $f \in C([0, \pi])$ and we may assume that $f(x) = \sum_{k=0}^{\infty} a_k \cos(kx)$ for $0 \leq x \leq \pi$. Similarly, if $\sum_{k=1}^{\infty} |b_k| < \infty$, then $g \in C([0, \pi])$ and we may assume that $g(x) = \sum_{k=1}^{\infty} b_k \sin(kx)$ for $0 \leq x \leq \pi$.

For (1.26) with $n = 0$, we have the following:

Theorem 6. Let g , (b_k) and ℓ be as in Theorem 5. We write $\bar{g}(x) := xg(1/x)$ for $x \geq 1/\pi$. Then

$$\sum_{k=m}^{\infty} b_k \sim \frac{\ell(m)}{m} \quad (m \rightarrow \infty) \quad (1.30)$$

if and only if

$$g \in C([0, \pi]) \quad \text{and} \quad \bar{g} \in \Pi_{\ell} \quad \text{with } \ell\text{-index } 1. \quad (1.31)$$

See also [I2, Theorem 1.2].

Theorems 4, 5 and 6 treat the boundary cases to the following known results due to Yong [Y]:

Theorem C ([Y]). Let f , (a_k) and ℓ be as in Theorem 4. Let $n \in \mathbb{N}$ and $2n - 1 < \rho < 2n + 1$. Then

$$\sum_{k=m}^{\infty} a_k \sim \frac{\ell(m)}{m^{\rho-1}} \cdot \frac{1}{\rho-1} \quad (m \rightarrow \infty) \quad (1.32)$$

if and only if $f \in C^{2n-2}([0, \pi])$ and

$$f(x) - \sum_{j=0}^{n-1} \frac{f^{(2j)}(0)}{(2j)!} x^{2j} \sim \frac{\pi}{2\Gamma(\rho) \cos(\rho\pi/2)} x^{\rho-1} \ell(1/x) \quad (x \rightarrow 0+). \quad (1.33)$$

Theorem D ([Y]). Let g , (b_k) and ℓ be as in Theorem 5. Let $n \in \mathbb{N}$ and $2n < \rho < 2n + 2$. Then

$$\sum_{k=m}^{\infty} b_k \sim \frac{\ell(m)}{m^{\rho-1}} \cdot \frac{1}{\rho-1} \quad (m \rightarrow \infty) \quad (1.34)$$

if and only if $g \in C^{2n-1}([0, \pi])$ and

$$g(x) - \sum_{j=0}^{n-1} \frac{g^{(2j+1)}(0)}{(2j+1)!} x^{2j+1} \sim \frac{\pi}{2\Gamma(\rho) \sin(\rho\pi/2)} x^{\rho-1} \ell(1/x) \quad (x \rightarrow 0+). \quad (1.35)$$

Theorems 4, 5 and 6, together with Theorems C and D, give an answer to [Bo, Question 7.19].

In this paper, we give a proof only to Theorem 1 which is our main theorem. For the rest, we refer to [IK].

2. Proof of Theorem A

As we stated in §1, Theorem A is due to [BI]. Since our main theorem, that is Theorem 1, is proved using Theorem A, we give its proof following [BI] for completeness.

Step 1. Choose X so large that h is positive and non-increasing on $[X, \infty)$. We first show that we lose no generality by supposing that h vanishes on $[0, X)$.

Set $\tilde{h}(t) := I_{[X, \infty)}(t)h(t)$, and let \tilde{H}_ν be its Hankel transform:

$$\tilde{H}_\nu(x) := \int_0^{\infty-} \tilde{h}(t)(xt)^{1/2} J_\nu(xt) dt \quad (x > 0).$$

By the mean-value theorem, there exists $c_1 \in (0, \infty)$ such that

$$|x^{-\nu} J_\nu(x) - y^{-\nu} J_\nu(y)| \leq c_1 |x - y| \quad (0 \leq x, y \leq 1).$$

So for $\lambda > 1$,

$$\begin{aligned} & \frac{1}{\ell(x)} |\{\tilde{H}_\nu(\lambda x) - \tilde{H}_\nu(x)\} - \{(\lambda x)^{\nu+\frac{1}{2}} \tilde{H}_\nu(1/\lambda x) - x^{\nu+\frac{1}{2}} \tilde{H}_\nu(1/x)\}| \\ &= \frac{1}{\ell(x)} \left| \int_0^X t^{\nu+\frac{1}{2}} h(t) \{(t/\lambda x)^{-\nu} J_\nu(t/\lambda x) - (t/x)^{-\nu} J_\nu(t/x)\} dt \right| \\ &\leq c_1 \frac{(1-\lambda^{-1})}{x\ell(x)} X \int_0^X t^{\nu+\frac{1}{2}} |h(t)| dt \rightarrow 0 \quad (x \rightarrow \infty). \end{aligned}$$

So (1.8) holds if and only if $x^{\nu+\frac{1}{2}} \tilde{H}_\nu(1/x) \in \Pi_\ell$ with ℓ -index $c_{\nu,0}$. Hence we may replace h by \tilde{h} — that is, we may assume that h vanishes on $[0, X)$.

Step 2: Abelian part. First we assume (1.7) and show (1.8). For $x > 0$,

$$\frac{1}{x^{\nu+\frac{3}{2}}h(x)} \left\{ \bar{H}_\nu(x) - c_{\nu,0} \int_0^x h(t)t^{\nu+\frac{1}{2}} dt \right\} = I(x) + II(x), \quad (2.1)$$

where

$$I(x) := \int_0^1 \frac{(xu)^{\nu+2}h(xu)}{x^{\nu+2}h(x)} \left\{ u^{1/2}J_\nu(u) - c_{\nu,0}u^{\nu+\frac{1}{2}} \right\} \frac{du}{u^{\nu+2}},$$

$$II(x) := \int_1^{\infty-} \frac{h(xu)}{h(x)} u^{1/2}J_\nu(u) du.$$

By the uniform convergence theorem for regularly varying functions ([BGT, Th. 1.5.2]), $(xu)^{\nu+2}h(xu)/x^{\nu+2}h(x)$ converges to $u^{1/2}$ as $x \rightarrow \infty$ uniformly in $u \in (0, 1]$.

Hence we find that

$$I(x) \rightarrow \int_0^1 \frac{1}{u^{\nu+\frac{3}{2}}} \left\{ u^{1/2}J_\nu(u) - c_{\nu,0}u^{\nu+\frac{1}{2}} \right\} du \quad (x \rightarrow \infty).$$

We note that the integral above converges absolutely since there exists $c_2 \in (0, \infty)$ such that

$$|x^{-\nu}J_\nu(x) - c_{\nu,0}| \leq c_2x^2 \quad (0 \leq x \leq 1). \quad (2.2)$$

In the same way, for any $Y > X$,

$$\int_1^Y \frac{h(xu)}{h(x)} u^{1/2}J_\nu(u) du \rightarrow \int_1^Y \frac{1}{u^{\nu+\frac{3}{2}}} u^{1/2}J_\nu(u) du \quad (x \rightarrow \infty).$$

By the second integral mean-value theorem ([WW, §4.14]), for $x > 1$,

$$\int_Y^{\infty-} \frac{h(xu)}{h(x)} u^{1/2}J_\nu(u) du = \frac{h(xY+)}{h(x)} \int_Y^{Y'} u^{1/2}J_\nu(u) du$$

for some $Y' \in (Y, \infty)$. If we set

$$c_3 := \sup \left\{ \left| \int_x^y u^{1/2}J_\nu(u) du \right| : 0 < x < y < \infty \right\},$$

then

$$\limsup_{x \rightarrow \infty} \left| \int_Y^{\infty-} \frac{h(xu)}{h(x)} u^{1/2}J_\nu(u) du \right| \leq \limsup_{x \rightarrow \infty} \frac{h(xY)}{h(x)} c_3 = \frac{c_3}{Y^{\nu+\frac{3}{2}}},$$

which can be made arbitrarily small by choosing Y large enough. So

$$II(x) \rightarrow \int_1^\infty \frac{1}{u^{\nu+\frac{3}{2}}} u^{1/2} J_\nu(u) du \quad (x \rightarrow \infty).$$

Therefore

$$\frac{1}{\ell(x)} \left\{ (\bar{H}_\nu(\lambda x) - c_{\nu,0} \int_0^{\lambda x} h(t)t^{\nu+\frac{1}{2}} dt) - (\bar{H}_\nu(x) - c_{\nu,0} \int_0^x h(t)t^{\nu+\frac{1}{2}} dt) \right\} \rightarrow 0$$

as $x \rightarrow \infty$. Hence, by the Uniform Convergence Theorem,

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{\bar{H}_\nu(\lambda x) - \bar{H}_\nu(x)}{\ell(x)} &= \lim_{x \rightarrow \infty} \frac{c_{\nu,0}}{\ell(x)} \int_x^{\lambda x} h(t)t^{\nu+\frac{1}{2}} dt \\ &= \lim_{x \rightarrow \infty} c_{\nu,0} \frac{x^{\nu+\frac{3}{2}} h(x)}{\ell(x)} \int_1^\lambda \frac{(xu)^{\nu+\frac{3}{2}} h(xu)}{x^{\nu+\frac{3}{2}} h(x)} \frac{du}{u} \\ &= c_{\nu,0} \int_1^\lambda \frac{du}{u} = c_{\nu,0} \log \lambda. \end{aligned}$$

Thus (1.7) implies (1.8).

Step 3: Tauberian part. We now prove the implication from (1.8) to (1.7). By a formula of Gegenbauer,

$$\int_0^\infty t^{\nu+\frac{1}{2}} e^{-xt} (yt)^{1/2} J_\nu(yt) dt = d_\nu \frac{xy^{\nu+\frac{1}{2}}}{(x^2 + y^2)^{\nu+\frac{3}{2}}} \quad (x > 0),$$

where $d_\nu := \pi^{-1/2} 2^{\nu+1} \Gamma(\nu + \frac{3}{2})$ ([WW, §13.2 (5)]). So by Parseval's formula for Hankel transforms ([RS]), for $x > 0$

$$\int_0^\infty h(t)t^{\nu+\frac{1}{2}} e^{-xt} dt = d_\nu \int_{0+}^{\infty-} H_\nu(y) \frac{xy^{\nu+\frac{1}{2}}}{(x^2 + y^2)^{\nu+\frac{3}{2}}} dy. \quad (2.5)$$

By the second integral mean-value theorem,

$$|H_\nu(y)| \leq c_3 h(X)/y \quad (0 < y < \infty), \quad (2.6)$$

whence

$$\int_1^\infty |H_\nu(y)| \frac{y^{\nu+\frac{1}{2}}}{(x^2+y^2)^{\nu+\frac{3}{2}}} dy < \infty.$$

On the other hand, by [BGT, Th. 3.7.4], (1.8) implies $|\bar{H}_\nu| \in R_0$, whence

$$\int_0^1 |H_\nu(y)| \frac{y^{\nu+\frac{1}{2}}}{(x^2+y^2)^{\nu+\frac{3}{2}}} dy = \int_1^\infty \frac{|\bar{H}_\nu(u)|}{(x^2u^2+1)^{\nu+\frac{3}{2}}} du < \infty.$$

Thus the integral on the right of (2.5) converges absolutely — and so the results of [BGT Ch. 4] apply.

We use Laplace transforms. Write

$$U(t) := \int_0^t h(u)u^{\nu+\frac{1}{2}} du \quad (t \geq 0),$$

$$\hat{U}(x) := \int_{[0,\infty)} e^{-xt} dU(t) = \int_0^\infty h(t)t^{\nu+\frac{1}{2}} e^{-xt} dt \quad (x > 0).$$

Then by (2.5),

$$\hat{U}(1/x) = (k * \bar{H}_\nu)(x) \quad (x > 0),$$

where

$$k(x) := d_\nu \frac{x^{2\nu+2}}{(1+x^2)^{\nu+\frac{3}{2}}} \quad (x > 0)$$

and $k * \bar{H}_\nu$ denotes the Mellin convolution

$$(k * \bar{H}_\nu)(x) = \int_0^\infty k(x/t)\bar{H}_\nu(t)dt/t.$$

The absolute convergence strip of the Mellin transform

$$\check{k}(z) := \int_0^\infty t^{-z}k(t)dt/t$$

is $-1 < \Re z < 2\nu + 2$, and for z in the strip

$$\check{k}(z) = \frac{2^\nu}{\pi^{1/2}} \Gamma(\nu + 1 - \frac{1}{2}z) \Gamma(\frac{1}{2} + \frac{1}{2}z),$$

in particular,

$$\check{k}(0) = 2^\nu \Gamma(\nu + 1).$$

By (2.6), \bar{H}_ν is locally bounded on $[0, \infty)$. So by the argument of [BGT, p. 242] (Abelian theorem for differences), we find that (1.8) implies $k * \bar{H}_\nu \in \Pi_\ell$ with ℓ -index 1 — i.e., so is $\hat{U}(1/\cdot)$. So by a Tauberian theorem of de Haan (cf. [BGT, Th. 3.9.1]), we see that $U \in \Pi_\ell$ with ℓ -index 1. Finally, by [BGT, Th. 3.6.10] (with slow decrease replaced by slow increase), we obtain (1.7), completing the proof. \square

3. Proof of Theorem 1

We note that (1.4) implies

$$\int_0^\infty t^{\nu - \frac{3}{2} + 2n} h(t) dt < \infty. \quad (3.1)$$

So, when proving the equivalence of (1.4) and (1.5), we may assume (3.1).

We define h_0, \dots, h_{n-1} by

$$\begin{aligned} h_0(t) &:= \int_t^\infty h(s) s^{\nu + \frac{1}{2}} ds \quad (0 \leq t < \infty), \\ h_j(t) &:= \int_t^\infty h_{j-1}(s) s ds \quad (0 \leq t < \infty, \quad j = 1, \dots, n-1). \end{aligned}$$

Since h is eventually non-negative, h_j are all eventually decreasing. By Fubini's theorem,

$$\begin{aligned} h_j(0) &= \int_0^\infty dt_j t_j \int_{t_j}^\infty dt_{j-1} t_{j-1} \cdots \int_{t_1}^\infty h(t_0) t_0^{\nu + \frac{1}{2}} dt_0 \\ &= \int_0^\infty dt_0 h(t_0) t_0^{\nu + \frac{1}{2}} \int_0^{t_0} dt_1 t_1 \cdots \int_0^{t_{j-1}} t_j dt_j \\ &= \frac{1}{2^j j!} \int_0^\infty h(t) t^{\nu + \frac{1}{2} + 2j} dt. \end{aligned}$$

Since

$$x^{-\mu} J_{\mu}(x) = O(1) \quad (x \rightarrow \infty),$$

$$\frac{d}{dx} \{x^{-\mu} J_{\mu}(x)\} = -x^{-\mu} J_{\mu+1}(x),$$

$$x^{-\mu} J_{\mu}(x) \rightarrow c_{\mu,0} \quad (x \rightarrow 0+)$$

for any $\mu \geq -1/2$ (see Watson [W], pages 199 and 45), we obtain, by integration by parts,

$$\begin{aligned} H_{\nu}(x) &= x^{\nu+\frac{1}{2}} \int_0^{\infty-} h(t) t^{\nu+\frac{1}{2}} \{(tx)^{-\nu} J_{\nu}(tx)\} dt \\ &= x^{\nu+\frac{1}{2}} h_0(0) c_{\nu,0} - x^{\nu+\frac{1}{2}+2} \int_0^{\infty-} h_0(t) t \{(tx)^{-\nu-1} J_{\nu+1}(tx)\} dt = \dots \\ &= \sum_{j=0}^{n-1} (-1)^j x^{\nu+\frac{1}{2}+2j} h_j(0) c_{\nu+j,0} \\ &\quad + (-1)^n x^{\nu+\frac{1}{2}+2n} \int_0^{\infty-} h_{n-1}(t) t \{(tx)^{-\nu-n} J_{\nu+n}(tx)\} dt \\ &= \sum_{j=0}^{n-1} (-1)^j x^{\nu+\frac{1}{2}+2j} h_j(0) c_{\nu+j,0} + (-1)^n x^n \int_0^{\infty-} g(t) (tx)^{1/2} J_{\nu+n}(tx) dt, \end{aligned}$$

where

$$g(t) := t^{-\nu+\frac{1}{2}-n} h_{n-1}(t) \quad (0 < t < \infty).$$

Since

$$(-1)^j h_j(0) c_{\nu+j,0} = c_{\nu,j} \int_0^{\infty} t^{\nu+\frac{1}{2}+2j} h(t) dt,$$

we have

$$\bar{H}_{\nu,n}(x) = (-1)^n x^{(\nu+n)+\frac{1}{2}} \int_0^{\infty-} g(t) (t/x)^{1/2} J_{\nu+n}(t/x) dt. \quad (3.2)$$

Now $t^{(\nu+n)+\frac{1}{2}} g(t) \in L^1_{\text{loc}}[0, \infty)$ and g is eventually decreasing to zero, whence by

Theorem A (with ν replaced by $\nu+n$) (1.5) is equivalent to

$$g(t) \sim t^{-\nu-n-\frac{3}{2}} \ell(t) \cdot \frac{(-1)^n c_{\nu,n}}{c_{\nu+n,0}} = t^{-\nu-n-\frac{3}{2}} \ell(t) \cdot \frac{1}{2^n n!} \quad (t \rightarrow \infty)$$

or

$$h_{n-1}(t) \sim t^{-2} \ell(t) \cdot \frac{1}{2^n n!} \quad (t \rightarrow \infty). \quad (3.3)$$

Since h_j is eventually decreasing, $\log \{h_j(t)t\}$ is slowly increasing, whence by the Monotone Density Theorem (see [BGT, §1.7]) (3.3) is equivalent to


$$h_0(t) = \int_t^\infty s^{\nu+\frac{1}{2}} h(s) ds \sim t^{-2n} \ell(t) \cdot \frac{1}{2n} \quad (t \rightarrow \infty). \quad (3.4)$$

By assumption, h is eventually decreasing, whence $\log \{h(t)t^{\nu+\frac{1}{2}}\}$ is slowly increasing. Again by the Monotone Density Theorem, (3.4) is equivalent to (1.4). This completes the proof. \square

REFERENCES

- [B] N. H. Bingham, *A Tauberian theorem for integral transforms of Hankel type*, J. London Math. Soc. (2) **5** (1972), 493–503.
- [BGT] N. H. Bingham, C. M. Goldie and J. L. Teugels, *Regular variation*, 2nd ed., Encycl. Math. Appl. **27**, Cambridge University Press, 1989 (1st ed. 1987).
- [BI] N. H. Bingham and A. Inoue, *An Abel-Tauber theorem for Hankel transforms*, Trends in probability and related analysis, Proceedings of SAP '96 (eds. N. Kono and N. R. Shieh), World Scientific, Singapore, 1997.
- [Bo] R. P. Boas, Jr., *Integrability theorems for trigonometric transforms*, Springer-Verlag, New York, 1967.
- [C] L. Y. Chan, *On Fourier series with non-negative coefficients and two problems of R. P. Boas*, J. Math. Anal. Appl. **110** (1985), 116–129.
- [I1] A. Inoue, *On Abel-Tauber theorems for Fourier cosine transforms*, J. Math. Anal. Appl. **196** (1995), 764–776.
- [I2] A. Inoue, *An Abel-Tauber theorem for Fourier sine transforms*, J. Math. Sci. Univ. Tokyo **2** (1995), 303–309.
- [I3] A. Inoue, *Abel-Tauber theorems for Fourier-Stieltjes coefficients*, J. Math. Anal. Appl. **211** (1997), 460–480.
- [IK] A. Inoue and H. Kikuchi, *Abel-Tauber theorems for Hankel and Fourier transforms and a problem of Boas*, submitted Hokkaido Math. J.
- [P] E. J. G. Pitman, *On the behaviour of the characteristic function of a probability distribution in the neighbourhood of the origin*, J. Austral. Math. Soc. **8** (1968), 422–443.
- [RS] J. R. Ridenhour and R. P. Soni, *Parseval relation and Tauberian theorems for the Hankel transform*, SIAM J. Math. Anal. **5** (1974), 809–821.
- [SS] K. Soni and R. P. Soni, *Slowly varying functions and asymptotic behavior of a class of integral transforms III*, J. Math. Anal. Appl. **49** (1975), 612–628.
- [Y] C. H. Yong, *On the asymptotic behavior of trigonometric series I*, J. Math. Anal. Appl. **33** (1971), 23–34.
- [W] G. N. Watson, *A treatise on the theory of Bessel functions*, 2nd ed., Cambridge University Press, 1944.
- [WW] E. T. Whittaker and G. N. Watson, *A course of modern analysis*, 4th ed., Cambridge University Press, 1927.
- [Z] A. Zygmund, *Trigonometric series*, 2nd ed., Cambridge University Press, 1959.


Hideyuki Kikuchi
Department of Mathematics
Hokkaido University
Sapporo 060, Japan
E-mail: h-kikuti@math.sci.hokudai.ac.jp



Inches 1 2 3 4 5 6 7 8
cm 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19

Kodak Color Control Patches

© Kodak, 2007 TM: Kodak



Kodak Gray Scale



© Kodak, 2007 TM: Kodak

A 1 2 3 4 5 6 **M** 8 9 10 11 12 13 14 15 **B** 17 18 19

