



HOKKAIDO UNIVERSITY

Title	Induction of viable gynogenetic progeny using eggs and UV-irradiated sperm from the Chinese tetraploid loach, <i>Misgurnus anguillicaudatus</i>
Author(s)	Li, Ya-Juan; Yu, Zhuo; Zhang, Ming-Zhao et al.
Citation	Aquaculture International, 21(4), 759-768 https://doi.org/10.1007/s10499-012-9551-3
Issue Date	2013-08-01
Doc URL	https://hdl.handle.net/2115/52956
Rights	The original publication is available at www.springerlink.com
Type	journal article
File Information	Revised ver AQU1 1545.pdf



Revised version manuscript AQU1 1545

Induction of viable gynogenetic progeny using eggs and UV irradiated sperm from the Chinese tetraploid loach, *Misgurnus anguillicaudatus*

Ya-Juan Li · Zhuo Yu · Ming-Zhao Zhang, · Cong Qian · Syuiti Abe · Katsutoshi Arai

Y.-J. Li · Z. Yu · M.-Z. Zhang · C. Qian

Life Science and Technology Institute, Dalian Ocean University, Dalian 116023, People's Republic of China

Syuiti Abe · Katsutoshi Arai (Corresponding author)

Faculty and Graduate School of Fisheries Sciences, Hokkaido University

Hakodate, Hokkaido 041-8611, Japan

e-mail: arai@fish.hokudai.ac.jp

Abstract When eggs from the Chinese tetraploid loach which had 100 chromosomes were fertilized with UV irradiated sperm, we obtained viable gynogenetic progeny without any additional treatment for the duplication of maternal chromosomes, which survived beyond first-feeding towards adult stage of development. Gynogenetic progeny were determined to be diploid since they possessed 50 chromosomes, along with two chromosomes bearing nucleolar organizing regions (NORs), detected by silver nitrate staining (Ag-NORs), chromomycin-A₃ (CMA₃) positive sites and fluorescence *in situ* hybridization (FISH) signals for rDNA loci. In contrast, when gynogens were induced using eggs from diploid loach fertilized by UV irradiated sperm, but without chromosome doubling, we found that all resultant progeny were non-viable haploid gynogens with 25 chromosomes, along with one NOR-bearing chromosome detected by Ag-NOR, CMA₃ and FISH. These observations demonstrate the true genetic tetraploid nature of the Chinese loach possessing 100 chromosomes and the potential use of this tetraploid as a source of functional diploid gametes for further ploidy manipulation experiments.

Keywords Chromosome · Gametes · FISH · Polyploid

Introduction

Diploid gametes represent key resources in order to expand the chromosome manipulation for improved breeding techniques. If we are able to use diploid gametes for breeding and subsequent manipulation, then it would allow the mass production of sterile auto- or allo-triploid lines, synthesis of new amphidiploids or allotetraploid lines, recovery of androgenotes from cryopreserved diploid sperm, and other applications for aquaculture (Arai 2000, 2001; Arai et al. 2010). However, successful examples of induced tetraploid fish as a source of diploid gametes have been limited due to the technical difficulties in producing tetraploids by inhibition of early cleavage coupled with very low survival rates of the resultant tetraploid progeny (Chourrout et al. 1986; Nam et al. 2001; Zou et al. 2004; Sakao et al. 2006). An alternative way would be to use the diploid gametes of natural tetraploid lines in certain species or hybrids.

In dojo loach or Oriental weatherfish *Misgurnus anguillicaudatus* (Teleostei: Cobitidae), most individuals of Japanese wild populations are diploid with 50 chromosomes. However, tetraploids possessing 100 chromosomes have often been identified amongst specimens recovered from Japanese markets (Ojima and Takai 1979; Arai et al. 1991a). Using chromosome manipulation techniques, Arai et al. (1991b, 1993) concluded that loaches possessing 100 chromosomes were true tetraploids possessing four sets of homologous chromosomes, because gynogenetic progeny which were artificially induced by fertilizing eggs from these market-sourced specimens with genetically-inert UV irradiated sperm were viable without requiring any treatment to duplicate the maternal chromosome set. In a similar manner, viable androgenetic progeny were successfully induced by fertilizing UV irradiated eggs with the sperm from loach possessing 100 chromosomes (Arai et al. 1995). Gynogenetic and androgenetic progeny produced from the gametes of normal diploid fish are non-viable due to the expression of abnormalities collectively referred as 'haploid syndrome', observed in many species of fish including loach (Suzuki et al. 1985; Arai et al. 1992; Fujimoto et al. 2007), common carp (Nagy et al. 1978) and salmonids (Chourrout et al. 1980; Chourrout 1982; Onozato 1982; Onozato and Yamaha 1983).

In central parts of China, natural tetraploid loach possessing 100 chromosomes co-exist, in the same habitats, with sympatric diploid loach possessing 50 chromosomes (Li et al. 2008). Cytogenetic studies including differential staining with silver nitrate for nucleolar organizing regions (Ag-NOR), fluorochromes such as chromomycin A₃ (CMA₃)/Distamycin (DA)/ 4',6-diamidino-2-phenylindole (DAPI) and fluorescence *in situ* hybridization (FISH) using human 5.8S+28S rDNA sequences suggested that a loach

possessing 100 chromosomes should be a tetraploid, based upon a karyotype showing quartets of chromosomes and the presence of four FISH signals (Li et al. 2010). However, specific genetic characteristic of these Chinese loaches have not yet been determined using the chromosome manipulation approach reported in Arai et al. (1991b, 1993, 1995). If Chinese loaches possessing 100 chromosomes are genetic tetraploids with four sets of homologous chromosomes, then it should be possible to use these as a source of diploid gametes in order to expand and improve existing breeding techniques. In the present study, we artificially induced gynogenesis using eggs from putative tetraploid females collected from the Chang Jiang River, Hubei Province, China by fertilizing with UV irradiated sperm and then carefully determined survival rates and chromosome numbers in the resultant gynogenetic progeny.

Materials and methods

Fish and gametes

Diploid (4 females, 4 males) and tetraploid (4 females) loach were collected in Chang Jiang River in Hubei Province, China and transported to Dalian Ocean University. Identification of tetraploid and diploid specimens was determined in accordance to Li et al. (2008). Artificial ovulation was induced by the injection of hCG (human Chorionic Gonadotropin, 20 IU per body gram weight, Aska Pharmaceutical Co. Ltd., Tokyo) according to Suzuki and Yamaguchi (1975). Sperm was collected into a capillary hematocrit tube by gently squeezing the male's abdomen (Morishima et al. 2002). Sperm was then diluted 1:100 with Kurokura solution (Kurokura et al. 1984).

Fertilization and gynogenesis

Eggs taken from diploid (2n) and tetraploid (4n) females were fertilized with sperm from 2n males and then, diploid female x diploid male (control cross, 2n x 2n) and tetraploid female x diploid male (triploid cross, 4n x 2n) crosses were performed. Eggs of 2n and 4n females were then inseminated with UV irradiated sperm from a 2n male (2n x UV and 4n x UV, respectively) to induce gynogenetic development. UV-irradiation of sperm from 2n males was performed according to Suzuki et al. (1985). Fertilized and treated eggs from each cross were reared in a Petri dish (90mm diameter x 15mm depth) containing freshwater (aged tap water) at room temperature.

Fertilization rate was calculated as the proportion of cleaved eggs relative to the initial number of eggs. Hatching rate was calculated as the proportion of hatched larvae relative to the number of

fertilized eggs. The normal rate was calculated as the proportion of normal larvae relative to the number of hatched larvae. Survival rate at 7 days after hatching (dah) was calculated as the proportion of surviving larvae relative to the number of hatched larvae. Water was changed every day and larvae were first fed *Artemia* from 3 days after hatching. Larvae were incubated at a temperature of 20°C.

Chromosome preparation and observation

Embryos at the optic vesicle stage (15 to 20 h after fertilization at 20°C) were used for chromosome preparation. In physiological saline, the chorion was mechanically removed and embryos treated for 45 min by soaking them in 0.0025% demecolcine (Sigma). Hypotonic treatment was performed with 0.8% citric acid for 20 min and samples were fixed with Carnoy's fixative (3 parts of methanol: 1 part of acetic acid). Chromosome preparations were prepared from developing embryos as described by Inokuchi et al. (1994). Chromosome slides were stained with Giemsa (Merck) for microscopic observation and karyotyping carried out in accordance with Levan et al. (1964).

Differential staining with CMA₃/DA/ DAPI (Schweizer 1976; Schweizer et al. 1978) and the Ag-NOR method (Howell and Black 1980) was applied in accordance with Li et al. (2010). FISH using human 5.8S+28S rDNA sequences as a probe was applied in accordance with Li et al. (2010).

Results

Viability of gynogenetic progeny

All control crosses (2n x 2n) resulted in high rates of fertilization (83-94%), hatching (75-96%), normal fry (84-97%) and survival at 7 dah (72-95%; Table 1). Triploid crosses (4n x 2n) also exhibited high rates of fertilization (77-98%), hatching (51-99%), normal fry (81-99%) and survival at 7 dah (77-90%; Table 1). Most control and triploid larvae exhibited normal appearance (Fig. 1A).

In gynogenetic crosses, differences were observed between 2n x UV and 4n x UV crosses. Gynogenetic progeny from 2n x UV crosses produced lower rates of fertilization (43-84%) and hatching (3.2-56%) than control and triploid crosses and no normal fry were apparent (Table 1). All resultant fry exhibited abnormalities characteristic of haploidy (Fig. 1B). In contrast, 4n x UV crosses produced high rates of fertilization (69-95%) and hatching (81-99%). Whilst 37 to 67% of resultant larvae were normal (Fig. 1C) or malformed (Fig. 1D), 46-60% fry survived to 7 dah (Table 1).

Chromosomal nature of gynogenetic progeny

As shown in Table 2 and Fig. 2AC, the modal chromosome number of embryos from $2n \times 2n$ crosses was diploidy ($2n=50$), while the mode of embryos from $4n \times 2n$ crosses was triploidy ($3n=75$). The $2n \times UV$ embryos exhibited a haploid range of chromosomes with a modal number of $1n=25$ (Table 2, Fig. 2B). In contrast, $4n \times UV$ embryos exhibited a diploid range of chromosomes with a modal number of $2n=50$ (Table 2, Fig. 2D). Consequently, most embryos from crosses $2n \times 2n$, $4n \times 2n$, $2n \times UV$ and $4n \times UV$ crosses were considered diploid ($2n=50$), triploid ($3n=75$), gynogenetic haploid ($1n=25$) and gynogenetic diploid ($2n=50$), respectively. However, hypo- and hyper-chromosome numbers were observed in each ploidy group (Table 2). The $2n \times UV$ embryos possessed one Ag-NOR/ CMA₃ positive region and FISH signal indicating haploidy (Fig 3A-C), while $4n \times UV$ embryos demonstrated two Ag-NOR/ CMA₃ positive regions and FISH signals indicating diploidy (Fig. 3 D-F).

Discussion

In the present study, viable gynogenetic progeny were produced without chromosome duplication from the eggs of Chinese tetraploid loach females fertilized with UV-irradiated loach sperm. Progeny possessed 50 chromosomes, equivalent to one half of the 100 chromosomes found in parental tetraploid loach specimens (Li et al. 2010). Gynogenetic fry possessing 25 chromosomes that were induced from the eggs of diploid females could not survive beyond hatching or feeding due to abnormalities. Diploidy of viable gynogenetic progeny was also demonstrated by the presence of two Ag-NORs, CMA₃ positive sites and FISH signals, whereas the haploid nature of non-viable gynogens was indicated by one site and one signal. Consequently, the production of viable gynogenetic progeny suggests that Chinese tetraploid loach are true tetraploids, possessing four sets of homologous chromosomes with the ability of generating diploid gametes. Thus, the tetraploid nature of the Chinese loach possessing 100 chromosomes was conclusively demonstrated by observing the viability of UV-induced gynogenetic progeny. Similar results have already been obtained in gynogenetic and androgenetic progeny induced from the gametes of loach possessing 100 chromosomes obtained from a Japanese market (Arai et al. 1991b, 1993, 1995), and in those induced from the gametes of natural tetraploid population in a different cobitid species *Cobitis biwae* (Kusunoki et al. 1994). Therefore, the overall genetic characteristics of Chinese tetraploid loach are essentially the same as those of tetraploid loach found in Japanese markets (Arai et al. 1991ab, 1993, 1995; Arai 2001, 2003). These results suggest the possibility that the tetraploid loach found in Japanese markets should be transported from China as food or fishing bait.

An important application of genetically tetraploid fish is as a source of diploid gametes for the expansion and improvement of existing ploidy manipulation techniques.

In Chinese loach, it is also possible to expand ploidy manipulation using diploid gametes as reported in our previous studies using natural tetraploid loach found in Japanese markets (Arai 2000, 2001, 2003). The mass production of triploids was first realized by crossing tetraploid and diploid fish (Matsubara et al. 1995; Zhang and Arai 1996) whilst a neo-tetraploid strain was produced by inhibiting second polar body release in the $2n \times 4n$ cross (Zhang and Arai 1996; Fujimoto et al. 2010). Fertile hexaploid lines were also produced by inhibiting second polar body release just after fertilization in $4n \times 4n$ crosses. Second generation hexaploid, pentaploid and tetraploid fish were then produced using fertile triploid gametes from hexaploid loaches (Arai et al. 1999). Diploid sperm from natural tetraploids is particularly useful for producing viable androgenotes by fertilizing UV-irradiated eggs (Arai et al. 1995; Yasui et al. 2010). Consequently, the Chinese tetraploid loach represents a valuable resource for inducing various levels of polyploids, gynogens and androgenotes for aquaculture. Breeding programs using diploid gametes are already implemented in allotetraploid crucian carp $\times$ common carp hybrids (Liu et al. 2001).

Acknowledgments This study was supported in part by Grants-in-Aid for Scientific Research (B) (No.21380114) from the Japan Society for the Promotion of Science (JSPS) to KA and by the Ronpaku (Dissertation Ph.D) Program (CSC-10610) from JSPS to YJL.

References

- Arai K (2000) Chromosome manipulation in aquaculture: recent progress and perspective. *Suisanzoshoku* 48:295-303
- Arai K (2001) Genetic improvement of aquaculture finfish species by chromosome manipulation techniques in Japan. *Aquaculture* 197: 205-228
- Arai K (2003) Genetics of the loach, *Misgurnus anguillicaudatus*: recent progress and perspective. *Folia biologica (Krakow)* 51 supplement:107-117
- Arai K, Matubara K, Suzuki R (1991a) Karyotype and erythrocyte size of spontaneous tetraploidy in the loach *Misgurnus anguillicaudatus*. *Nippon Suisan Gakkaishi* 57:2167-2172
- Arai K, Matubara K, Suzuki R (1991b) Chromosomes and developmental potential of progeny of spontaneous tetraploid loach *Misgurnus anguillicaudatus*. *Nippon Suisan Gakkaishi* 57:2173-2178
- Arai K, Masaoka T, Suzuki R (1992) Optimum condition of UV ray irradiation for genetic inactivation of loach eggs. *Nippon Suisan Gakkaishi* 58:1197-1201

217 Arai K, Matsubara K, Suzuki R (1993) Production of polyploids and viable gynogens
 218 using spontaneously occurring tetraploid loach, *Misgurnus anguillicaudatus*.
 219 Aquaculture 117: 227-235.

220 Arai K, Ikeno M, Suzuki R (1995) Production of androgenetic diploid loach *Misgurnus*
 221 *anguillicaudatus* using spermatozoa of natural tetraploids. Aquaculture
 222 137:131-138

223 Arai K, Taniura K, Zhang Q (1999) Production of second generation progeny of
 224 hexaploid loach. Fisheries Sci 65: 186-192

225 Arai K, Sakao S, Fujimoto T, Yamaha E (2010) Cutting edge of chromosome
 226 manipulation for aquaculture and conservation of salmonids. National Taiwan
 227 Museum Special Publication 14:49-52.

228 Chourrout D (1982) Gynogenesis caused by ultraviolet irradiation of salmonid sperm. J
 229 Exp Zool 223:175-181

230 Chourrout D, Chevassus B, Herioux F (1980) Analysis of an Hertwig effect in the
 231 rainbow trout (*Salmo gairdneri* Richardson) after fertilization with
 232 gamma-irradiated sperm. Reprod Nutr Develop 20(3A): 719-726

233 Chourrout D, Chevassus B, Krief F, Happe A, Burger G, Renard P (1986) Production of
 234 second generation triploid and tetraploid rainbow trout by mating tetraploid
 235 males and diploid females—Potential of tetraploid fish. Theor Appl Genet
 236 72:193-206

237 Fujimoto T, Sakao S, Yamaha E, Arai K (2007) Evaluation of different doses of UV
 238 irradiation to loach eggs for genetic inactivation of the maternal genome. J Exp
 239 Zool Part A 307:449-462

240 Fujimoto T, Yasui GS, Hayakawa M, Sakao S, Yamaha E, Arai K (2010) Reproductive
 241 capacity of neo-tetraploid loaches produced using diploid spermatozoa from a
 242 natural tetraploid male. Aquaculture 308: S133-S139

243 Howell WM, Black DA (1980) Controlled silver-staining of nucleolus organizer regions
 244 with a protective colloidal developer: A 1-step method. Experientia 36:1014-1015

245 Inokuchi T, Abe S, Yamaha E, Yamazaki F, Yoshida MC (1994) BrdU replication banding
 246 studies on the chromosomes in early embryos of salmonid fishes. Hereditas
 247 121:255-265

248 Kurokura H, Hirano R, Tomita M, Iwahashi M (1984) Cryopreservation of carp sperm.
 249 Aquaculture 37:267-273

250 Kusunoki T, Arai K, Suzuki R (1994) Production of viable gynogens without
 251 chromosome duplication in the spinous loach *Cobitis biwae*. Aquaculture 119:
 252 11-23

253 Levan A, Fredga K, Sandberg AA (1964) Nomenclature for centromeric position on
 254 chromosomes. *Hereditas* 52: 201-220.
 255 Li Y-J, Yin J, Wang J-B, Yuan X, Wei J, Sun X-W, Arai K (2008) A study on the
 256 distribution of polyploidy loaches in China. *Nippon Suisan Gakkaishi*
 257 74:177-182
 258 Li Y-J, Tian Y, Zhang M-Z, Tian P-P, Yu Z, Abe S, Arai K (2010) Chromosome banding
 259 and FISH with rDNA in the diploid and tetraploid loach *Misgurnus*
 260 *anguillicaudatus*. *Ichthyol Res* 57:358-366
 261 Liu S, Liu Y, Zhou G, Zhang X, Luo C, Feng H, He X, Zhu G, Yang H (2001) The
 262 formation of tetraploid stocks of red crucian carp x common carp hybrids as
 263 effect of interspecific hybridization. *Aquaculture* 192:171-186
 264 Matsubara K, Arai K, Suzuki R (1995) Survival potential and chromosomes of progeny
 265 of triploid and pentaploid females in the loach, *Misgurnus anguillicaudatus*.
 266 *Aquaculture* 131:37-48
 267 Morishima K, Horie S, Yamaha E, Arai K (2002) A cryptic clonal line of the loach
 268 *Misgurnus anguillicaudatus* (Teleostei: Cobitidae) evidenced by induced
 269 gynogenesis, interspecific hybridization, microsatellite genotyping and
 270 multilocus DNA fingerprinting. *Zool Sci* 19:565-575.
 271 Nagy A, Rajki K, Horvath L, Csanyi V (1978) Investigation on carp *Cyprinus carpio* L.
 272 gynogenesis. *J Fish Biol* 13:215-224
 273 Nam YK, Choi GC, Park DJ, Kim DS (2001) Survival and growth of induced tetraploid
 274 mud loach. *Aquacult Int* 9:6-11.
 275 Ojima Y, Takai A (1979) The occurrence of spontaneous polyploidy in the Japanese
 276 common loach, *Misgurnus anguillicaudatus*. *Proc Japan Acad* 55B:487-491
 277 Onozato H (1982) The "Hertwig effect" and gynogenesis in chum salmon *Oncorhynchus*
 278 *keta* eggs fertilized with ⁶⁰Co gamma-ray irradiated milt. *Bull Jpn Soc Sci Fish*
 279 48:1237-1244
 280 Onozato H, Yamaha E (1983) Induction of gynogenesis with ultraviolet rays in four
 281 species of salmoniformes. *Bull Jpn Soc Sci Fish* 49:693-699
 282 Sakao S, Fujimoto T, Kimura S, Yamaha E, Arai K (2006) Drastic mortality in tetraploid
 283 induction results from the elevation of ploidy in masu salmon *Oncorhynchus*
 284 *masou*. *Aquaculture* 252:147-160
 285 Schweizer D (1976) Reverse fluorescent chromosome banding with chromomycin and
 286 DAPI. *Chromosoma* 58:307-324
 287 Schweizer D, Ambros P, Andrlé M (1978) Modification of DAPI banding on human
 288 chromosomes by prestaining with a DNA-banding oligopeptide antibiotic,

distamyci A. Exp Cell Res 111:327-332

Suzuki R, Yamaguchi M (1975) Influence of water temperature on inducing spawning by hormone injection in the loach, cyprinid fish. Suisanzoshoku 22:135-139

Suzuki R, Oshirao T, Nakanishi T (1985) Survival, growth and fertility of gynogenetic diploids induced in the cyprinid loach, *Misgurnus anguillicaudatus*. Aquaculture 48:45-55

Yasui GS, Fujimoto T, Arai K (2010) Restoration of the loach, *Misgurnus anguillicaudatus*, from cryopreserved diploid sperm and induced androgenesis. Aquaculture 308: S140-S144

Zhang Q, Arai K (1996) Flow cytometry for DNA contents of somatic cells and spermatozoa in the progeny of natural tetraploid loach. Fisheries Sci 62: 870-877



Zou S, Li S, Cai W, Zhao J, Yang H (2004) Establishment of fertile tetraploid population of blunt snout bream (*Megalobrama amblycephala*). Aquaculture 238:155-164



Legends for figures

Figure 1 Normal larvae from 2n x 2n (A), abnormal larvae from 2n x genetically inactivated sperm with UV irradiation (UV) (B), normal larvae from 4n x UV (C) and malformed larvae from 4n x UV (D). Bars indicate 1 mm.

Figure 2 Metaphase spread with (A) 50 chromosomes from a 2n x 2n embryo, (B) 25 chromosomes from a 2n x UV embryo, (C) 75 chromosomes from a 4n x 2n embryo, and (D) 50 chromosomes from a 4n x UV embryo (D). Bars indicate 10 μ m.

Figure 3 Metaphase spreads in 2n x UV (A, B, C) and 4n x UV (D, E, F) dojo loach embryos. Ag-NORs (A, D), CMA₃/DA/DAPI staining positive sites (B, E) and FISH signals detected with 5.8S+28S rDNA probe (C, F) are shown. Arrows indicate NORs shown by Ag staining (A, D), CMA₃ positive sites (B, E) or FISH signals (C, F). Bars indicate 10 μ m.












Table 1. Number of eggs, fertilization rate, hatching rate, normal rate and survival rate at 7 days after hatch in different crosses

Cross	No. of eggs	Fertilization rate (%)	Hatching rate (%)	Normal rate (%)	Survival rate at 7 days after hatch (%)
2nx2n-1	412	92.95	79.36	91.14	75.79
2nx2n-2	716	83.58	74.51	96.7	94.86
2nx2n-3	448	66.07	95.61	91.17	71.73
2nx2n-4	414	93.96	78.41	84.26	73.77
mean±SD		84.14±0.13	81.97±0.09	90.82±0.05	79.04±0.11
2nxUV-1	701	75.89	3.2	0	0
2nxUV-2	766	56.27	9	0	0
2nxUV-3	433	42.96	27.96	0	0
2nx2nUV-	565	83.89	56.33	0	0
mean±SD		64.75±0.19	24.12±0.24	0	0
4nx2n-1	1210	97.69	51.02	88.72	76.78
4nx2n-2	850	77.41	91.79	98.68	89.74
4nx2n-3	778	89.2	99.14	80.52	77.91
4nx2n-4	524	90.65	94.32	84.38	84.38
mean±SD		88.74±0.08	84.06±0.22	88.07±0.08	82.20±0.06
4nxUV-1	480	95.21	81.33	66.67	54.53
4nxUV-2	489	73.62	98.89	44.1	59.55
4nxUV-3	695	68.78	96.23	37.39	49.13
4nxUV-4	744	70.97	92.42	48.57	45.7
mean±SD		77.14±0.12	92.22±0.08	49.18±0.13	52.23±0.06

Table 2. Metaphase chromosome count in Dojo loach embryos originating from different crosses

Cross	Embryo number	Cell number	Chromosome number frequency																		
			23	24	25	26	27	...	40	41	42	43	44	45	46	47	48	49	50	51	52
2n x 2n	17	72	0	0	0	0	0		1	3	2	2	2	1	4	4	5	2	41	2	3
2n x UV	11	39	5	6	26	1	1		0	0	0	0	0	0	0	0	0	0	0	0	0
4n x 2n	16	77	0	0	0	0	0		0	0	0	0	0	0	0	0	0	0	0	0	0
4n x UV	17	100	0	0	0	0	0		0	0	0	2	1	0	3	10	9	3	44	3	6

Cross	Chromosome number frequency																		
	53	54	56	57	...	66	67	68	69	70	71	72	73	74	75	76	77	78	79
2n x 2n	0	0	0	0		0	0	0	0	0	0	0	0	0	0	0	0	0	0
2n x UV	0	0	0	0		0	0	0	0	0	0	0	0	0	0	0	0	0	0
4n x 2n	0	0	0	0		1	0	2	4	5	3	8	5	2	31	5	1	4	6
4n x UV	5	11	2	1		0	0	0	0	0	0	0	0	0	0	0	0	0	0

2n x 2n: control cross from fertilization of eggs of diploid female with sperm of diploid male, 2n x UV: gynogenetic cross from fertilization of eggs of diploid female with UV-irradiated sperm, 4n x 2n: cross from fertilization of eggs of tetraploid female with sperm of diploid male, 4n x UV: gynogenetic cross from fertilization of eggs of tetraploid female with UV-irradiated sperm