



HOKKAIDO UNIVERSITY

Title	The relation between habitual sleep duration and blood pressure values in Japanese male subjects
Author(s)	Satoh, Hiroki; Nishihira, Jun; Wada, Tatsuhiko et al.
Citation	Environmental Health and Preventive Medicine, 2013(18), 215-220 https://doi.org/10.1007/s12199-012-0309-3
Issue Date	2013-09-27
Doc URL	https://hdl.handle.net/2115/53243
Rights	The original publication is available at http://www.springerlink.com .
Type	journal article
File Information	18_3_215.pdf



**The Relation between the Habitual Sleep Duration and Blood Pressure Values in
Japanese Male Subjects**

Hiroki SATOH ^{1, 2}, Jun NISHIHIRA ¹, Tatsuhiko WADA ¹, Satoshi FUJII ³

and Hiroyuki TSUTUI ²

¹ Department of Clinical Management and informatics, Hokkaido Information University,

Ebetsu, Japan

² Department of Cardiovascular Medicine, Hokkaido University Graduate School of Medicine,

Sapporo, Japan

³ Department of Molecular and Cellular Pathobiology and Therapeutics,

Nagoya City University Graduate School of Pharmaceutical Sciences, Nagoya, Japan

Correspondence to: Hiroki SATOH

Department of Clinical Management and informatics, Hokkaido Information University

59-2, Nishi-Nopporo, Ebetsu, Hokkaido, 069-8585, Japan

Telephone: +81 11 385 4411 Fax: +81 11 384 0134

E-mail: h-satoh@do-johodai.ac.jp

Keywords: sleep duration; blood pressure; hypertension; Japanese male subjects;

epidemiology.

Abstracts

Background: Previous studies have demonstrated that the sleep duration is closely associated with metabolic risk factors, however, the relationship between the habitual sleep duration and blood pressure values in Japanese population has not been fully established.

Methods: We performed a cross-sectional study of 1,670 Japanese male subjects to clarify the relationship between the habitual sleep duration and blood pressure values. The study subjects were divided into four groups such as <6, 6-, 7-, and ≥ 8 hours according to their nightly habitual sleep duration.

Results: The rate of subjects with <6, 6-, 7-, and ≥ 8 hours sleep duration was 12.0, 37.6, 38.2, and 12.2 (%), respectively. Compared to the group with 7-hours sleep duration (referent), those with <6 and ≥ 8 hours groups have significantly greater systolic and diastolic blood pressure values. The rate of hypertensive subjects, defined as systolic blood pressure more than 140 mmHg and/or diastolic blood pressure more than 90 mmHg, with sleep duration with <6, 6-, 7-, and ≥ 8 hours was 13.4, 5.7, 7.5, and 13.8 (%), respectively. Compared to the group with 7-hours sleep duration (referent), the multivariate odds ratios (95% confidence interval) of that with <6 and ≥ 8 hours for hypertension were 2.43 (1.40-4.20, $P < 0.01$) and 2.28 (1.31-3.95, $P < 0.01$), respectively, adjusted with conventional cardiovascular risk factors.

Conclusions: The present study demonstrated both long and short habitual sleep duration was

significant high blood pressure values and the occurrence of hypertension in Japanese male subjects.

1 Introduction

2 Sleep loss, long-term sleep deprivation, and alternations in sleep duration are common
3 in modern society [1], with evidence showing that we are sleeping on average only 6.8 hours
4 per night, which is 1.5 hours less than we did one hundred years ago [2]. The change of sleep
5 condition exerts deleterious effects on detectable changes in metabolic [3, 4], endocrine [5],
6 and sympathetic tone [6]. These findings suggest that alternations of the habitual sleep
7 duration may predispose to overt the change of blood pressure values. Several previous
8 studies demonstrated that shorter sleep durations were found to be related to hypertension,
9 compared with subjects with 7 hours of sleep per night in Caucasian populations [7-10]. The
10 first United States National Health and Nutrition Examination Survey (NHANES-I)
11 elucidated that sleeping 5 hours or less per night was associated with a 60% increased risk of
12 incident hypertension in Caucasian middle-aged subjects during a mean follow-up of 8-10
13 years [10]. The Sleep Heart Health Study demonstrated that United States subjects sleeping
14 more than 9 hours per night had a 30% higher prevalence of hypertension as compared to
15 those sleeping between 7 and 8 hours per night [7]. The relationship between sleep duration
16 and the occurrence of hypertension in previous studies has been controversial, however, this
17 association among Japanese general population has not been fully investigated. Moreover, the
18 relationship between sleep duration and blood pressure values remain unanswered.

19 The purpose of the present study was to examine cross-sectional associations of sleep

duration with blood pressure values and hypertension occurrence in Japanese male subjects.

Materials and Methods

Study Subjects

The study subjects consisted of 1,887 food and beverage company male employees, aged from 38 from 58 years, who had their annual health examinations during the period from April 2010 to March 2011. A total of 216 subjects were excluded for the following reasons: medication of hypertension (n=201), medication of sleep disturbance (n=12), and medication of depression (n=3). Thus, 1,671 male subjects were enrolled in the present study. The study protocol was approved by the ethical committee of Hokkaido Information University and the written informed consent was obtained from each subjects.

Baseline data measurement

All study subjects were asked to complete a self-questionnaire that included smoking habit, alcohol consumption, the frequency of exercise, family history of hypertension, and medical history. The questionnaire was distributed to the subjects in advance of their annual health check-up and was collected at the examination. The subjects who had never smoked and ex-smokers were classified as “non-smokers”. Drinkers were defined as those who consumed alcohol once per week or more. “Exercise” subjects were defined as those who regularly exercised more than one per week. Hypertension, dyslipidemia, and diabetes mellitus were defined as receiving medical agents at the health examination. Body mass index

(BMI) was calculated as body weight (kilograms) divided by squared height (meters). Blood pressure was measured by a trained nurse using a standard mercury sphygmomanometer, with the study subjects in the sitting position after at least a 5-min rest. A blood sample was obtained from the antecubital vein in the morning after an overnight fast and serum was separated. After precipitation by heparin-manganase, total cholesterol and high-density lipoprotein (HDL)-cholesterol were measured by the phosphotungstate method. Triglyceride was measured by enzymatically. Glucose was enzymatically determined by the hexokinase method.

Definition of hypertension

Incident hypertension subjects were defined as systolic blood pressure ≥ 140 mmHg and/or diastolic blood pressure ≥ 90 mmHg at their health examination.

Statistical Analysis

The study subjects were categorized into four groups according to sleep duration: <6 , 6-, 7-, and ≥ 8 hours. The characteristics of the study subjects were expressed as means $\pm$ SD for continuous variables, median (and interquartile range) for skewed distribution variables, and percentages for categorical variables according to sleep duration. The differences of variables among groups were examined by analysis of variance (ANOVA), Kruskal-Wallis

test, or chi-square test. Next, Bonferroni procedure was used between two groups as a post hoc test. The association between sleep duration and hypertension was assessed by using the multiple logistic regression analysis. The principle model included candidate variables for age, BMI, smoking, alcohol, exercise, dyslipidemia, diabetes mellitus, family history of hypertension, values such as systolic blood pressure, total cholesterol, triglyceride, high density lipoprotein (HDL)-cholesterol, and glucose. A p value of less than 0.05 was considered to indicate statistical significance. All statistical analyses were performed using the SPSS statistical package for Windows version 11.0 (Chicago, IL, USA).

Results

Table 1 shows the characteristics of study subjects according to sleep duration levels.

The mean age and BMI of the study patients was 50 ± 6 years and $23.8 \pm 2.8 \text{ kgm}^{-2}$, respectively.

The numbers (%) of the study subjects with sleep duration of <6, 6-, 7-, and ≥ 8 hours were

201 (12.0), 628 (37.6), 638 (38.2), and 203 (12.2), respectively. Variables such as age, BMI,

alcohol, exercise, family history of hypertension, triglyceride, and high density lipoprotein

(HDL)-cholesterol were significantly different among the groups. Subjects with 7-hours sleep

duration were older and more likely to have non-drinkers.

Figure 1 shows systolic (A) and diastolic (B) blood pressure values according to

sleep duration levels. The mean systolic and diastolic blood pressure values of the study

subjects were 119 ± 15 and 74 ± 11 mmHg, respectively. The mean systolic blood pressure

values with sleep duration <6, 6-, 7-, and ≥ 8 hours were 122 ± 14 , 119 ± 15 , 118 ± 15 , and

122 ± 17 mmHg, respectively (P for trend <0.01). The mean diastolic blood pressure values

with sleep duration <6, 6-, 7-, and ≥ 8 hours were 76 ± 11 , 75 ± 10 , 73 ± 11 , and 78 ± 12

mmHg, respectively (P for trend < 0.001). Compared with subjects with sleep duration of 7-

hours as a referent, those with sleep duration of both <6 and ≥ 8 hours had significantly

greater systolic and diastolic blood pressure values.

Table 2 shows that multivariate predictor of variables for hypertension by multiple

logistic regression analysis. The numbers (%) of the study subjects with hypertension were

139 (8.3%). Risk factors such as age, BMI, smoking, alcohol, family history of hypertension, and glucose were significant and independent predictors for hypertension.

Table 3 shows odds ratio and 95% confidence interval (CI) of hypertension according to sleep duration levels. The numbers (%) of hypertensive subjects with sleep duration <6, 6-, 7-, and ≥8 hours were 27 (13.4), 36 (5.7), 48 (7.5), and 28 (13.8), respectively. On multivariate analysis with sleep duration of 7- hours as the reference, the adjusted odds ratio for subjects with sleep duration <6 hours for hypertension was 2.43 (95%CI: 1.40-4.20, P <0.01). Similarly, that with sleep duration ≥8 hours for hypertension was 2.28 (95%CI: 1.31-3.95, P <0.01).

Discussion

The present study demonstrates that sleep duration of both <6 and ≥ 8 hours are significantly associated with high blood pressure values and hypertension occurrence in Japanese male subjects.

Bonnet et al. reported that nearly one-third of adults had sleep duration less than 6 hours per night and sleep deficiency, long-term sleep deprivation, and alternations in sleep duration have been common in modern society [1]. The quality and sleep duration have been indicated as factors to affect the health condition in general population [11, 12]. In epidemiologic studies, compared with subjects with 7-hours of habitual sleep duration, short and long sleep durations were related to increase body mass index [13, 14], the development of diabetes mellitus [15, 16], and the occurrence of cardiovascular disease [17-19]. Recent epidemiological cohort studies indicated that short and long sleep duration were also closely associated with mortality [20, 21]. Ikehara et al. demonstrated that both shorter and longer sleep duration were associated with increased mortality due to all causes for both gender, yielding a U-shaped relationship with total mortality with a nadir as 7-hours sleep duration from a large-scale prospective study of 98,634 Japanese men and women in the Japan Collaborative Cohort Study [22]. However, the relationship between sleep duration and blood pressure values in Japanese general population has not been fully established.

Previous studies indicated that sleep disorders might play a crucial role in

determining blood pressure values both in the office and over the 24 hours and modulating the day-night blood pressure profile [23, 24], which could affect the prognosis of hypertensive patients [25, 26]. Recent studies demonstrated that both increased and reduced sleep duration may be related to increase the risk of cardiovascular disease and hypertension [22, 27]. These studies clarified that sleep duration defined as acutely induced sleep deprivation was closely associated with blood pressure values, however, the relationship between the habitual sleep duration and blood pressure values has still been controversial [28].

Previous studies reported that the habitual short sleep duration increased blood pressure values in general population [29], however, this relationship might differ among ages and ethnicity [7, 30]. The present study demonstrated that the habitual short sleep duration was significantly associated with high blood pressure values and the occurrence of hypertension in Japanese middle-aged male subjects, which confirmed the previous results. The biological mechanisms mediating the association of the habitual short sleep duration with the development and occurrence of hypertension are uncertain, however, the habitual short sleep duration may disrupt circadian rhythm and autonomic balance [30, 31]. The alternations in these functions might shift of the daily blood pressure profile to higher values, the occurrence of the non-dipping pattern, the increase of blood pressure variability, and the disturbances in the diurnal rhythm of cardiac output [32, 33].

The present study also demonstrated that the habitual long sleep duration was

significantly associated with high blood pressure values and the occurrence of hypertension. The rate of drinkers was more likely to be higher in subjects with sleep duration ≥ 8 than < 6 , 6-, and 7- hours (Table 1) and the covariate of alcohol was a significant and independent predictor for hypertension (Figure 2). Sesso et al. indicated that light-to-moderate alcohol consumption was a linear association with increasing the risk for the development of hypertension [34]. Nakanishi et al. demonstrated that the risk for hypertension increased according to the increment of alcohol consumption doses in Japanese middle-aged male subjects [35]. Thus, these results may be a potential causal basis for the result in the present study. Patel et al. reported that C-reactive peptide and interleukin-6 levels rose with increasing the habitual sleep duration [36]. These inflammatory mediators might also increase the risk of hypertension in subjects with the habitual long sleep [37].

The limitations of the present study are as follows. First, the sleep duration obtained by a self questionnaire might differ from the measured precious sleep duration. However, Lockely et al reported that good agreement was examined between self-reported sleep duration and those obtained through actigraphic monitoring [38]. Second, the present study examined blood pressure values once in the health examination. The accuracy of blood pressure values would likely have a little effect on the results of this analysis. Third, the present study was a cross-sectional observation study. The possibility of unmeasured confounding variables, such as sedentary life style and obstructive sleep apnea, could not be

1 excluded.

2 In conclusion, we demonstrated that habitual both short and long sleep duration were
3 significantly associated with high blood pressure values and the occurrence of hypertension in
4 Japanese male subjects. Finally, it is important to note that adequate sleep duration should be
5 one of the important strategies for hypertension occurrence.

1 **Figure legend**

2 Figure 1

3 Systolic (A) and diastolic (B) blood pressure values according to sleep duration levels.

4 *: P for trend.

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

1 **Conflict of interest**

2 The authors declare that they have no conflict of interest.

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

1 **References**

- 2 (1) Bonnet MH, Arand DL. We are chronically sleep deprived. *Sleep* 1995; 18: 908-11.
- 3 (2) National Sleep Foundadtion: Sleep in America Poll 2003. Washington, DC, National
4 Sleep Founddation. 2003;
- 5 (3) Knutson KL, Spiegel K, Penev P, Van Cauter E. The metabolic consequences of sleep
6 deprivation. *Sleep Med Rev* 2007; 11: 163-78.
- 7 (4) Spiegel K, Tasali E, Leproult R, Van Cauter E. Effects of poor and short sleep on
8 glucose metabolism and obesity risk. *Nat Rev Endocrinol* 2009; 5: 253-61.
- 9 (5) Spiegel K, Leproult R, L'Hermite-Baleriaux M, Copinschi G, Penev PD, Van Cauter E.
10 Leptin levels are dependent on sleep duration: relationships with sympathovagal
11 balance, carbohydrate regulation, cortisol, and thyrotropin. *J Clin Endocrinol Metab*
12 2004; 89: 5762-71.
- 13 (6) Spiegel K, Leproult R, Van Cauter E. Impact of sleep debt on metabolic and endocrine
14 function. *Lancet* 1999; 354: 1435-9.
- 15 (7) Gottlieb DJ, Redline S, Nieto FJ, Baldwin CM, Newman AB, Resnick HE, et al.
16 Association of usual sleep duration with hypertension: the Sleep Heart Health Study.
17 *Sleep* 2006; 29: 1009-14.
- 18 (8) Cappuccio FP, Stranges S, Kandala NB, Miller MA, Taggart FM, Kumari M, et al.
19 Gender-specific associations of short sleep duration with prevalent and incident

hypertension: the Whitehall II Study. *Hypertension* 2007; 50: 693-700.

- (9) Stang A, Moebus S, Mohlenkamp S, Erbel R, Jockel KH. Gender-specific associations of short sleep duration with prevalent hypertension. *Hypertension* 2008; 51: e15-6; author reply e17.

- (10) Staessen JA, Thijs L, Fagard R, O'Brien ET, Clement D, de Leeuw PW, et al. Predicting cardiovascular risk using conventional vs ambulatory blood pressure in older patients with systolic hypertension. Systolic Hypertension in Europe Trial Investigators. *JAMA* 1999; 282: 539-46.

- (11) Kapur VK, Redline S, Nieto FJ, Young TB, Newman AB, Henderson JA. The relationship between chronically disrupted sleep and healthcare use. *Sleep* 2002; 25: 289-96.

- (12) Shankar A, Charumathi S, Kalidindi S. Sleep duration and self-rated health: the national health interview survey 2008. *Sleep* 2011; 34: 1173-7.

- (13) Stranges S, Cappuccio FP, Kandala NB, Miller MA, Taggart FM, Kumari M, et al. Cross-sectional versus prospective associations of sleep duration with changes in relative weight and body fat distribution: the Whitehall II Study. *Am J Epidemiol* 2008; 167: 321-9.

- (14) Patel SR, Hu FB. Short sleep duration and weight gain: a systematic review. *Obesity (Silver Spring)* 2008; 16: 643-53.

- 1 (15) Ayas NT, White DP, Al-Delaimy WK, Manson JE, Stampfer MJ, Speizer FE, et al. A
2 prospective study of self-reported sleep duration and incident diabetes in women.
3 Diabetes Care 2003; 26: 380-4.
- 4 (16) Yaggi HK, Araujo AB, McKinlay JB. Sleep duration as a risk factor for the
5 development of type 2 diabetes. Diabetes Care 2006; 29: 657-61.
- 6 (17) Greenland P, Knoll MD, Stamler J, Neaton JD, Dyer AR, Garside DB, et al. Major risk
7 factors as antecedents of fatal and nonfatal coronary heart disease events. JAMA 2003;
8 290: 891-7.
- 9 (18) Sabanayagam C, Shankar A. Sleep duration and cardiovascular disease: results from
10 the National Health Interview Survey. Sleep 2010; 33: 1037-42.
- 11 (19) Shankar A, Koh WP, Yuan JM, Lee HP, Yu MC. Sleep duration and coronary heart
12 disease mortality among Chinese adults in Singapore: a population-based cohort study.
13 Am J Epidemiol 2008; 168: 1367-73.
- 14 (20) Rod NH, Vahtera J, Westerlund H, Kivimaki M, Zins M, Goldberg M, et al. Sleep
15 disturbances and cause-specific mortality: Results from the GAZEL cohort study. Am
16 J Epidemiol 2011; 173: 300-9.
- 17 (21) Kripke DF, Langer RD, Elliott JA, Klauber MR, Rex KM. Mortality related to
18 actigraphic long and short sleep. Sleep Med 2011; 12: 28-33.
- 19 (22) Ikehara S, Iso H, Date C, Kikuchi S, Watanabe Y, Wada Y, et al. Association of sleep

duration with mortality from cardiovascular disease and other causes for Japanese men and women: the JACC study. *Sleep* 2009; 32: 295-301.

(23) Lored JS, Nelesen R, Ancoli-Israel S, Dimsdale JE. Sleep quality and blood pressure dipping in normal adults. *Sleep* 2004; 27: 1097-103.

(24) Friedman O, Logan AG. Nocturnal blood pressure profiles among normotensive, controlled hypertensive and refractory hypertensive subjects. *Can J Cardiol* 2009; 25: e312-6.

(25) Li Y, Boggia J, Thijs L, Hansen TW, Kikuya M, Bjorklund-Bodegard K, et al. Is blood pressure during the night more predictive of cardiovascular outcome than during the day? *Blood Press Monit* 2008; 13: 145-7.

(26) Fagard RH, Thijs L, Staessen JA, Clement DL, De Buyzere ML, De Bacquer DA. Night-day blood pressure ratio and dipping pattern as predictors of death and cardiovascular events in hypertension. *J Hum Hypertens* 2009; 23: 645-53.

(27) Eguchi K, Pickering TG, Schwartz JE, Hoshida S, Ishikawa J, Ishikawa S, et al. Short sleep duration as an independent predictor of cardiovascular events in Japanese patients with hypertension. *Arch Intern Med* 2008; 168: 2225-31.

(28) Stranges S, Dorn JM, Shipley MJ, Kandala NB, Trevisan M, Miller MA, et al. Correlates of short and long sleep duration: a cross-cultural comparison between the United Kingdom and the United States: the Whitehall II Study and the Western New

- 1 York Health Study. *Am J Epidemiol* 2008; 168: 1353-64.
- 2 (29) Kim J, Jo I. Age-dependent association between sleep duration and hypertension in the
- 3 adult Korean population. *Am J Hypertens* 23: 1286-91.
- 4 (30) Gangwisch JE, Heymsfield SB, Boden-Albala B, Buijs RM, Kreier F, Pickering TG, et
- 5 al. Short sleep duration as a risk factor for hypertension: analyses of the first National
- 6 Health and Nutrition Examination Survey. *Hypertension* 2006; 47: 833-9.
- 7 (31) Folkow B. "Structural factor" in primary and secondary hypertension. *Hypertension*
- 8 1990; 16: 89-101.
- 9 (32) Folkow B. Mental stress and its importance for cardiovascular disorders; physiological
- 10 aspects, "from-mice-to-man". *Scand Cardiovasc J* 2001; 35: 163-72.
- 11 (33) Coca A. Circadian rhythm and blood pressure control: physiological and
- 12 pathophysiological factors. *J Hypertens Suppl* 1994; 12: S13-21.
- 13 (34) Sesso HD, Cook NR, Buring JE, Manson JE, Gaziano JM. Alcohol consumption and
- 14 the risk of hypertension in women and men. *Hypertension* 2008; 51: 1080-7.
- 15 (35) Nakanishi N, Yoshida H, Nakamura K, Suzuki K, Tatara K. Alcohol consumption and
- 16 risk for hypertension in middle-aged Japanese men. *J Hypertens* 2001; 19: 851-5.
- 17 (36) Patel SR, Zhu X, Storfer-Isser A, Mehra R, Jenny NS, Tracy R, et al. Sleep duration
- 18 and biomarkers of inflammation. *Sleep* 2009; 32: 200-4.
- 19 (37) Patel SR. Sleep--an affair of the heart. *Sleep* 2009; 32: 289-90.

- 1 (38) Lockley SW, Skene DJ, Arendt J. Comparison between subjective and actigraphic
- 2 measurement of sleep and sleep rhythms. J Sleep Res 1999; 8: 175-83.

3

Table 1.

Characteristics of the study subjects according to sleep duration levels

	Total (n=1,670)	Sleep duration				P value
		< 6 (n=201)	6- (n=628)	7- (n=638)	≥ 8 (n=203)	
Age (years)	50 ± 6	48 ± 5	49 ± 6	51 ± 6	51 ± 5	<0.001
Body mass index (kg/m ²)	23.8 ± 2.8	24.4 ± 3.1	23.8 ± 2.8	23.6 ± 2.6	23.8 ± 2.8	<0.05
Smoking (%)	52.2	54.7	52.1	51.7	51.5	0.90
Alcohol (%)	74.3	74.1	73.6	71.2	85.2	<0.01
Exercise (%)	28.2	20.9	31.7	27.2	27.9	<0.05
Dyslipidemia (%)	4.0	4.0	3.5	3.8	6.4	0.32
Diabetes mellitus (%)	3.2	1.5	3.8	3.3	2.5	0.38
Family history of hypertension (%)	15.6	21.4	16.9	14.6	9.4	<0.01
Total cholesterol (mg/dL)	208 ± 33	206 ± 36	208 ± 31	208 ± 34	208 ± 33	0.90
Triglyceride (mg/dL)	106 (76-150)	104 (76-150)	127 (87-173)	105 (73-146)	104 (75-152)	<0.05
HDL-cholesterol (mg/dL)	57 ± 14	56 ± 13	57 ± 15	56 ± 14	59 ± 15	<0.05
Glucose (mg/dL)	96 ± 20	94 ± 12	96 ± 19	97 ± 20	97 ± 24	0.13

Values are means ± SD, median (and interquartile range), and percentage.

HDL, high-density lipoprotein.

Table 2.

Multivariate predictors of variables for hypertension by multiple logistic regression analysis

	Odds ratio	95%CI	P value
Age (per one year increase)	1.04	1.01-1.08	<0.05
Body mass index (per one kg/m ² increase)	1.17	1.09-1.25	<0.001
Smoking	1.87	1.26-2.78	<0.01
Alcohol	2.23	1.31-3.77	<0.01
Family history of hypertension	1.78	1.13-2.79	<0.05
Glucose (per 10mg/dL increase)	1.15	1.07-1.23	<0.001

CI, confidence interval.

Table 3.

Odds ratio and 95% CI for hypertension according to sleep duration levels


	Sleep duration (hours)			
	< 6 (n=201)	6- (n=628)	7- (n=638)	8 ≤ (n=203)
Hypertension n (%)	27 (13.4)	36 (5.7)	48 (7.5)	28 (13.8)
Unadjusted				
Odds ratio	2.73	1.33	1.00	2.45
95% CI	1.60-4.64	0.84-2.10	–	1.45-4.15
P value	<0.001	0.23	–	<0.01
Model 1 ^a				
Odds ratio	2.49	1.28	1.00	2.40
95% CI	1.43-4.33	0.80-2.05	–	1.40-4.12
P value	<0.01	0.30	–	<0.01
Model 2 ^b				
Odds ratio	2.43	1.27	1.00	2.28
95% CI	1.40-4.20	0.81-1.99	–	1.31-3.95
P value	<0.01	0.30	–	<0.01

^aadjusted for age, body mass index, smoking, alcohol, exercise, and family history of hypertension.^bModel 1 + adjusted for diabetes mellitus, dyslipidemia, total cholesterol, triglyceride, HDL-cholesterol, and glucose.

CI, confidence interval; HDL high-density lipoprotein.

Figure 1

(A) Systolic blood pressure



(B) Diastolic blood pressure

