



Title	Boundary values of quasiconformal mappings
Author(s)	Zajac, Jozwf
Citation	Hokkaido University technical report series in mathematics, 14, 1
Issue Date	1989-01-01
DOI	https://doi.org/10.14943/5133
Doc URL	https://hdl.handle.net/2115/5448
Type	departmental bulletin paper
File Information	14.pdf



**Boundary values of quasiconformal
mappings**

Jozef Zajac

Series # 14. July, 1989

HOKKAIDO UNIVERSITY
TECHNICAL REPORT SERIES IN MATHEMATICS

- # 1: T. Morimoto, Equivalence Problems of the Geometric Structures admitting Differential Filtrations, 14 pages. 1987.
- # 2: J.L. Heitsch, The Lefschetz Theorem for Foliated Manifolds, 59 pages. 1987.
- # 3: K. Kubota (Ed.), 第 12 回偏微分方程式論札幌シンポジウム予稿集, 77 pages. 1987.
- # 4: J. Tilouine, Kummer's criterion over Λ and Hida's Congruence Module, 85 pages. 1987.
- # 5: Y. Giga (Ed.), Abstracts of Mathematical Analysis Seminar 1987, 17 pages. 1987.
- # 6: T. Yoshida (Ed.), 1987 年度談話会アブストラクト集 Colloquium Lectures, 96 pages. 1988.
- # 7: S. Izumiya, G. Ishikawa (Eds.), “特異点と微分幾何” 研究集会報告集, 1988.
- # 8: K. Kubota (Ed.), 第 13 回偏微分方程式論札幌シンポジウム予稿集, 76 pages. 1988.
- # 9: Y. Okabe (Ed.), ランジュヴァン方程式とその応用予稿集, 64 pages. 1988.
- # 10: I. Nakamura (Ed.), Superstring 理論と K3 曲面, 91 pages. 1988.
- # 11: Y. Kamishima (Ed.), 1988 年度談話会アブストラクト集 Colloquium Lectures, 73 pages. 1989.
- # 12: G. Ishikawa, S. Izumiya and T. Suwa (Eds.), “特異点論とその応用” 研究集会報告集 Proceedings of the Symposium “Singularity Theory and its Applications,” 317 pages. 1989.
- # 13: M. Suzuki (Ed.), “駆け足で有限群を見てみよう” 1987 年 7 月北大での集中講義の記録, 38 pages. 1989.

BOUNDARY VALUES OF QUASICONFORMAL MAPPINGS

by
JOZEF ZAJAC


Notations.

Institute of Mathematics PAN, 90-136 Łódź, ul. Narutowicza 56, POLAND

- \mathbb{E} - closed plane,
- \mathbb{C} - open, complex plane,
- $\bar{\mathbb{C}}$ - complex plane compactified with one point at infinity,
- \mathbb{R} - real line,
- E, E' - sets,
- $\text{fr}E$ - topological boundary of E , ∂E - oriented boundary of E ,
- $\text{cl}E$ - the closure of E ,
- $\text{int}D$ - the interior of D ,
- D - Jordan domain,
- Jordan curves - a homeomorphic image of a circle,
- Jordan arc - a homeomorphic image of a segment, i.e. a connected subset of \mathbb{R} , which does not reduce to a point,
- (a, b) - open segment,
- $\langle a, b \rangle$ - closed segment
- $\Delta = \{z : |z| \leq 1\}$, $\Delta_r = \{z : r \leq |z| \leq 1\}$,
- $\{a_n\}$ - a sequence of numbers, f, g - functions, f^{-1} - inverse funct.
- $\{f_n\}$ - a sequence of functions,
- $f_n \rightarrow f$ - pointwise convergence, $f_n \rightrightarrows f$ - almost uniform convergence.

A sense-preserving homeomorphism.

Suppose that D is a Jordan domain and f is a homeomorphism such that $f: \text{cl}D \rightarrow E$. Then there exists a Jordan domain D' such, that $E = \text{cl}D'$, $f: D \rightarrow D'$, $f|_{\text{fr}D}: \text{fr}D \rightarrow \text{fr}D'$



By the definition of a Jordan domain, there exists a homeomorphism f_1, f_2 such that $f_i: \text{fr } \Delta \rightarrow \text{fr } D$, $i = 1, 2$, and $\arg f_2^{-1} \circ f_1|_{\text{fr } \Delta \setminus \{-1\}}$ is an increasing function of $\arg z$, $z \in \text{fr } \Delta \setminus \{-1\}$.

It is well known that also $g_1 = f_1 \circ f_2^{-1}$ and $g_2 = f_2 \circ f_1^{-1}$ are homeomorphisms such that $\arg g_2^{-1} \circ g_1|_{\text{fr } \Delta \setminus \{-1\}}$ is an increasing function of $\arg z$, $z \in \text{fr } \Delta \setminus \{-1\}$.

Hence f induces a mapping between the orientation of $\text{fr } \Delta$ and the orientation of $\text{fr } D$. If both the orientations are positive or negative with respect to the corresponding domain then $f: D \rightarrow D'$ is said to be sense-preserving.

More general if $f: E \rightarrow E'$ is a homeomorphism of two sets then f is said to be sense-preserving if $f|_D$ is sense-preserving for every Jordan domain D , such that $\text{cl } D \subset E$.

Proposition 1 (Newmann [1])

If $f: E \rightarrow E'$ is a homeomorphism and E is either a domain or the closure of a Jordan domain and there exist a Jordan domain D such that $\text{cl } D \subset E$ and $f|_D$ is sense-preserving, then f is sense-preserving on E .

It is worth-while to note that if f is a sense-preserving homeomorphism, then also is f^{-1} . If f_1 and f_2 are sense-preserving homeomorphisms then also $f_2 \circ f_1$ is the sense-preserving homeomorphism provided the composition makes sense.

Differentiability

Suppose now that E is either an open set or the closure of a Jordan domain. A mapping $f: E \rightarrow E'$ is said to be differentiable at $z_0 \in \text{int } E$ ($z_0, f(z_0) \neq \infty$) if

$$(1) \quad f(z) = f(z_0) + f'_1(z_0)(z_0 - z) + f'_2(z_0)(\bar{z}_0 - \bar{z}) + o(|z - z_0|)$$

where: $f_z = \frac{1}{2}(f_x - if_y)$, $f_{\bar{z}} = \frac{1}{2}(f_x + if_y)$, $z = x + iy$.

Also $\overline{(f_z)} = (\bar{f})_{\bar{z}}$, $\overline{(f_{\bar{z}})} = (\bar{f})_z$,

$$(g \circ f)_z = g_z \circ f \cdot f_z + g_{\bar{z}} \circ f \cdot \bar{f}_z,$$

$$(g \circ f)_{\bar{z}} = g_z \circ f \cdot f_{\bar{z}} + g_{\bar{z}} \circ f \cdot \bar{f}_{\bar{z}}.$$

A mapping f is said to be differentiable at $z = \infty$ if f^* defined by $f^*(z) = f(1/z)$ is differentiable at $z = 0$. A mapping f is said to be differentiable at z_0 such that $f(z_0) = \infty$ if f^{**} defined by $f^{**}(z) = 1/f(z)$ is differentiable at z_0 .

A mapping f is differentiable on a set E if it is differentiable at every point of E .

The directional derivatives $f_{|d}$ are defined by $f_{|d} = e^{-id} [f_t(z + te^{id})]_t$ where d is real and $t \geq 0$.

The Jacobian of f will be denoted by J or J_f . If $z = \infty$, or $f(z_0) = \infty$ for some $z_0 \in E$, we will ^{not} define J_f but only the $\text{sgn } J_f^* = \text{sgn } J_{f^{**}}(z_0)$.

It is easily verified that if $f: E \rightarrow \mathbb{C}$, where $E \subset \mathbb{C}$, is differentiable then

$$(2) \quad f_{|d} = e^{-id} (f_x \cos d + f_y \sin d) = f_z + e^{-2id} f_{\bar{z}},$$

$$(3) \quad \max_{\alpha} |f_{|\alpha}| = |f_z| + |f_{\bar{z}}|, \quad \min_{\alpha} |f_{|\alpha}| = ||f_z| - |f_{\bar{z}}||,$$

$$(4) \quad J_f = |f_z|^2 - |f_{\bar{z}}|^2 = \text{sgn } J_f \max_{\alpha} |f_{|\alpha}| \cdot \min_{\alpha} |f_{|\alpha}| = \frac{1}{2} |f_x \bar{f}_y - f_y \bar{f}_x|$$

If f is differentiable at $z_0 \in \text{int } E$ and $\text{sgn } J_f \neq 0$, then f is called regular at z_0 and z_0 is called a regular point of f . A mapping f is called regular in E if it is regular at every point $z \in \text{int } E$.

A C^1 class regular homeomorphism is called a diffeomorphism. In the case when $z_0 = \infty$ or $f(z_0) = \infty$ the definition of C^1 fund

may be extended similarly to that of differentiability

Suppose now that $f: E \rightarrow E'$ is a homeomorphism and E is either a domain or the closure of a Jordan domain. Then by Newman's result quoted above, if $\text{sgn } J_f(z) = 1$ at some regular point z , then f is sense-preserving and, conversely, if f is sense-preserving then $\text{sgn } J_f(z) = 1$ at any regular $z \in \text{int } E$.

Suppose that f is a diffeomorphism and D is a domain. The ratio

$$(5) \quad p(z) = p_f(z) = \frac{\max_{\alpha} |f_{\alpha}^{\alpha}(z)|}{\min_{\alpha} |f_{\alpha}^{\alpha}(z)|} = \frac{|f_z'(z)| + |f_{\bar{z}}'(z)|}{||f_z'| - |f_{\bar{z}}'|}, \quad z \in D$$

is called the dilatation of f at z . Clearly it is bounded on every compact subset of D and invariant under conformal mappings. I.e. if g_1 and g_2 are conformal such that $f \circ g_1$ and $g_2 \circ f$ makes sense, then $p_{f \circ g_1} = p_f = p_{g_2 \circ f}$.

The last conclusion enables us to extend the definition of p to the cases $z_0 = \infty$ and $f(z_0) = \infty$ analogously to the definition of differentiability was extended.

Definition (Grötzsch)

A sense-preserving diffeomorphism $f: D \rightarrow D'$, where D is a domain and there exist constant number Q , $1 \leq Q < \infty$ such that

$$(6) \quad \sup p_f(z) \leq Q$$

is called a regular Q -quasiconformal mapping (Q -qc).

This very natural definition has the disadvantage that the class of regular Q -qc mappings is not close with respect to almost uniform convergence. Thus

Definition (Gehring, Lehto)

A homeomorphism $f: D \rightarrow D'$, D -being a domain, is said to be a G -qc mapping, if there exist a sequence of regular

of regular quasiconformal mappings $f_n: D \rightarrow D'$ such that $f_n \rightarrow f$, and for a.e. z for which there exists finite partial derivatives $f_z, f_{\bar{z}}$ we have $f_{nz}/f_{n\bar{z}} \rightarrow f_z/f_{\bar{z}}$.

Properties:

- 1° 1-qc mapping $f: D \rightarrow D'$ is identical with a class of conformal mappings,
- 2° if f is a Q -qc mapping then also f^{-1} is Q -qc mapping
- 3° if f_1 is a Q_1 -qc mapping and f_2 is a Q_2 -qc mapping then also $f_1 \circ f_2$ is a $Q_1 Q_2$ -qc mapping
- 4° it is enough to consider classes of qc mapping on special domains like; unit disc Δ , an annulus Δ_N , etc. (canonical domain which are conformally equivalent to a class of domains).

Complex dilatation and Beltrami equation

If f is a sense-preserving homeomorphism differentiable almost everywhere on a set E , where E is either an open set or a closure of a Jordan domain, then the mapping given by its differential taken at a differentiable point $z_0 \in \text{Int} E$ has the form

$$(7) \quad dw = f_z dz + f_{\bar{z}} d\bar{z}$$

and map a unit (infinitesimal) circle $|dz| = ds$ onto the ellipse with half axis $a = |f_z| + |f_{\bar{z}}|$, $b = |f_z| - |f_{\bar{z}}|$ and the center at $f(z_0)$.

Writing down the dilatation ρ_f in the form

$$\rho_f = \frac{|f_z| + |f_{\bar{z}}|}{|f_z| - |f_{\bar{z}}|} = \frac{1 + \left| \frac{f_{\bar{z}}}{f_z} \right|}{1 - \left| \frac{f_{\bar{z}}}{f_z} \right|} = \frac{1 + |\mu|}{1 - |\mu|}$$

where $\mu = f_{\bar{z}}/f_z$, we attain at the complex function μ .

which is called complex dilatation of f . If f is regular Q -qc mapping then $|\mu| < \frac{Q-1}{Q+1}$. Starting from a given complex dilatation we may consider the differential equation $f_{\bar{z}} = \mu f_z$ which is an elliptic type equation known as the Beltrami differential equation. This is when μ satisfies certain conditions. Confining ourselves to the case of the unit disc we present the most general result of Bojarski which in our context is as

Proposition 2

If μ is a measurable function defined in Δ and such that $\|\mu\|_\infty = \inf_E \sup_{z \in \Delta \setminus E} |\mu(z)|$, where the infimum is taken over all set E with Lebesgue measure $|E|=0$, is bounded from 1 i.e. $\|\mu\|_\infty < 1$. Then there exist a unique solution $w=f(z)$ of


$$(B) \quad f_{\bar{z}} = \mu f_z$$

which is a Q -qc selfmapping of Δ with $Q = \frac{1+\|\mu\|_\infty}{1-\|\mu\|_\infty}$ and normalization condition $f(0)=0, f(1)=1$ or, equivalently $f(1)=1, f(i)=i, f(-1)=-1$.

Geometrical characterization of quasiconformal mappings.

A quadrilateral consist of a Jordan domain D and a sequence z_1, z_2, z_3, z_4 of boundary points of D . The points $z_i, i=1,2,3,4$ are called vertices of the quadrilateral. We confine ourselves to the only quadrilaterals $D(z_1, z_2, z_3, z_4)$ whose sequence of vertices agrees with positive orientation with respect to D . The vertices divide the boundary of D into four Jordan arcs, the sides of quadrilateral, the arcs $\widehat{z_1 z_2}$ and $\widehat{z_3 z_4}$ are called the a-sides and the other two arcs, the b-sides of D .

It is well known that image of D under sense-preserving homeomorphism is a quadrilateral.



$$f(D) = D', \quad w_i = f(z_i)$$

It is not, in general, possible to map given quadrilateral onto another, since the image of three boundary points determine the mapping uniquely. All quadrilaterals are therefore divided into several classes which are conformally equivalent.

It follows from Riemann mapping theorem that every quadrilateral $D(z_1, z_2, z_3, z_4)$ can be mapped conformally onto a quadrilateral $H(-\frac{1}{k}, -1, 1, \frac{1}{k})$, where $0 < k < 1$ and H is the upper half-plane. From the classical theory of the elliptic integrals we further obtain, that the function

$$(8) \quad \int(z) = \int_0^z \frac{ds}{\sqrt{(1-s^2)(1-k^2s^2)}}$$

maps $H(-\frac{1}{k}, -1, 1, \frac{1}{k})$ conformally onto a quadrilateral which consist of rectangle and its corners. Composing the above mappings, we can map an arbitrary quadrilateral conformally onto a rectangle. Such a mapping will be called the canonical mapping. All canonical rectangles of a given quadrilateral $D(z_1, z_2, z_3, z_4)$ have therefore the same ratio of sides $M(D) = \frac{a}{b}$ - conformal module of it.

Two quadrilaterals are conformally equivalent if and only if they possess the same ratio of a and b . If one carries on a rearrangement of the vertices of D , the the module behaves as follows

$$M(D(z_1, z_2, z_3, z_4)) = M(D(z_3, z_4, z_1, z_2)) = 1/M(D(z_2, z_3, z_4, z_1)).$$

Definition.

A sense-preserving homeomorphism of E where E is either a domain or a closure of a Jordan domain, is called Q -qc mapping if there exist a finite number $Q \geq 1$ such that

$$(9) \quad \left(\frac{1}{Q}\right) M(D) \leq M(f(D)) \leq Q \cdot M(D)$$

holds for all quadrilaterals $D, dD \subset E$.

Boundary values problem.

This is a problem of existence and continuity of the boundary values under quasiconformal mappings. But the solvability of boundary value problem means the possibility of constructing a quasiconformal mapping with given boundary values.

The simple example of the mapping $f(z) = z - \rho(1 - |z|^2)$, ρ -suff. small shows that the identity mapping $f(e^{i\theta}) = e^{i\theta}$, $-\pi < \theta \leq \pi$ has many solution in the case of the unit disc, when the solution must be qc. It has, of course, exactly one, when we confine ourselves to the conformal automorphism.

Let D and D' be n -tuply connected domains which both have n Jordan curves as boundary components. Then every conformal mapping $f: D \rightarrow D'$ can be extended to a homeomorphism of its closures. If these domains are Jordan domains and the orientation of z_1, z_2, z_3 and w_1, w_2, w_3 are both positive with respect to the corresponding domains, then there exist a unique conformal mapping $f: D \rightarrow D' : f|_{\partial D} : \partial D \rightarrow \partial D'$ - is the homeomorphism and $f(z_i) = w_i, i = 1, 2, 3$. For qc mapping we state the main result as


Proposition 3 If D and D' are Jordan domains then every Q -qc mapping $f: D \rightarrow D'$ can be extended to a homeomorphism of $dD \rightarrow dD'$.

We restrict ourselves to Q -qc mapping between Jordan

domains. It follows from Proposition 3 that every quasiconformal mapping can be extended to a homeomorphism between closure of these domains. By the orientation-preserving theorem (which is in our context as a proposition) the extension is sense-preserving, and the boundary value problem therefore reads as follows.

Let D and D' be two Jordan domains with boundaries C and C' respectively, and φ be a homeomorphism of C onto C' which preserves the positive orientation. It is required to find necessary and sufficient condition on φ such that there exists a qc mapping $f: D \rightarrow D'$ with the boundary values $f(z) = \varphi(z)$.

In view of the invariance of qc mapping under composition with conformal mappings the problem of characterizing the induced homeomorphism φ can be reduced to the case, when $D = D' = H = \{z: \text{Im} z > 0\}$. Then the boundary correspondence is determined by a monotone continuous function φ in this sense that the point $(x, 0)$ corresponds to $(\varphi(x), 0)$. It is sufficiently to consider the case when φ is strictly increasing.



By the definition of Q -quasiconformality based on the moduli of a quadrilateral we see that if F is a Q -qc self-mapping of the upper half-plane H then

$$\frac{1}{Q} M(G) \leq M(F(G)) \leq Q M(G)$$

holds for all quadrilaterals G , $dG \subset H$. Since H is the Jordan domain than we may consider this definition not only for a domain but also for closure of a Jordan domain. Then (9) must also be true when $G=H$. The condition (9) assumes a weaker form, better suited to our applications, if we fix the vertex z_4 of a quadrilateral and choose the other three vertices such that G is conformally equivalent to a square. Then $M(G)=1$ so G' must satisfy the condition

$$(9') \quad \frac{1}{Q} \leq M(G') \leq Q.$$


On the other hand, this necessary condition is also sufficient in the following sense: it guarantees the existence of qc, though not necessarily Q -qc mapping. The order of quasiconformality may increase.

To transform condition (9') to a simple form we set $z_4 = \infty = x_4$ and normalize all our mapping so that they preserve the point at infinity. The quadrilateral $H(x_1, x_2, x_3, \infty)$ is conformally equivalent to a square if $x_3 - x_2 = x_2 - x_1$. Then we write $x_2 = x$, $x_1 = x+t$, $x_3 = x-t$, where $t > 0$ since the orientation is positive with respect to H .

For the moduli of quadrilateral $H' = H(x'_1, x'_2, x'_3, \infty)$ we have

$$(10) \quad M(H') = \frac{2}{\pi} \mu \left(\sqrt{\frac{x'_1 - x'_2}{x'_3 - x'_1}} \right)$$

where $\mu(t)$ denotes the conformal moduli of the unit disc sliced along real line from 0 to t , $0 < t < 1$ and is strictly decreasing with limits ∞ at 0 and 0 at 1



$\mu(t)$ - module of double connected domain, which is $\Delta \setminus \{0, t\}$

The function μ can be also expressed in terms of elliptic integral $K(t) = \int_0^1 \frac{dx}{\sqrt{(1-x^2)(1-t^2x^2)}}$ and $\mu(t) = \frac{\pi}{2} \frac{K(\sqrt{1-t^2})}{K(t)}$.

If we set this value into (9') we obtain for the boundary values of Q -qc mapping $F: H \rightarrow H$, $F(\infty) = \infty$, the double inequality

$$\mu^{-1}\left(\frac{\pi Q}{2}\right) \leq \sqrt{\frac{\varphi(x+t) - \varphi(x)}{\varphi(x+t) - \varphi(x-t)}} \leq \mu^{-1}\left(\frac{\pi}{2Q}\right)$$

To obtain this condition in the symmetric form we write

$$(11) \quad \lambda(Q) = \frac{1}{\left(\mu^{-1}\left(\frac{\pi Q}{2}\right)\right)^2} - 1$$

Then since $\mu\left(\frac{1}{\sqrt{1+\lambda(Q)}}\right) = \frac{\pi Q}{2}$ and $\mu\left(\frac{1}{\sqrt{1+\lambda(1/Q)}}\right) = \frac{\pi}{2Q}$, then $\mu\left(\frac{1}{\sqrt{1+\lambda(Q)}}\right) \cdot \mu\left(\frac{1}{\sqrt{1+\lambda(1/Q)}}\right) = \frac{\pi^2}{4}$ do not depends on Q and we may replace $\lambda(Q)$ by $1/\lambda(1/Q)$ by which $\lambda(1/Q) = 1/\lambda(Q)$.

Thus we obtain

Theorem

The boundary value of a Q -qc self-mapping F of the upper half-plane preserving the point at ∞ satisfy the double inequality

$$(12) \quad 1/\lambda(Q) \leq \frac{\varphi(x+t) - \varphi(x)}{\varphi(x) - \varphi(x-t)} \leq \lambda(Q)$$

for all real x and $t > 0$.

Solution of the boundary values problem

In 1956 Beurling and Ahlfors constructed a quasiconformal self-mapping of the upper half-plane with given boundary values. But the construction is only possible when the boundary values satisfy a condition of (12) form. Then they have show that this condition is also sufficient, although the maximal dilatation of the constructed mapping turns out to be larger than the value arising from (9). Moreover if $Q(\tilde{f})$ is the qc dilatation of f , then $Q(\tilde{f}) \geq 1 + 0.2284 \log s$ for each ϵ extension \tilde{f} of a function satisfying (12) when ϵ is instead of $\lambda(\epsilon)$.

Theorem (Beurling - Ahlfors)

Let φ be a strictly increasing, continuous function of the real which satisfies the condition

$$(13) \quad \frac{1}{\delta} \leq \frac{\varphi(x+t) - \varphi(x)}{\varphi(x) - \varphi(x-t)} \leq \delta$$

for every real $x, t > 0$. Then there exist a qc selfmapping F^* of the upper half-plane, which has the boundary values $F^*_{\mathbb{R}} = \varphi$ and a maximal dilatation less than a bound depending on δ .

Proof.

Let us consider a function of the form $F^*(z) = u + iv$, where $u = \frac{1}{2}(\alpha + \beta)$, $v = \frac{1}{2}(\alpha - \beta)$, where

$$(14) \quad \alpha(x, y) = \int_0^1 \varphi(x + ty) dt, \quad \beta(x, y) = \int_0^1 \varphi(x - ty) dt.$$

This function F^* is defined and continuous in the whole finite \mathbb{C}_z plane. Since φ is strictly increasing, F^* carries every point of the upper half-plane $\text{Im} z > 0$ into the upper half-plane $\text{Im} w > 0$, where $w = u + iv$. On the real line $F^*(x) = \varphi(x)$ and at conjugate points $z = x + iy, \bar{z} = x - iy$ we have $F^*(\bar{z}) = \overline{F^*(z)}$. We have to prove that F^* is a quasiconformal mapping of the whole plane, then the image domain will also consist of the whole plane, and it follows from the above that F^* maps the upper-half plane \mathbb{H} onto itself and has the right boundary values $\varphi(x)$.

We show first that F^* is a homeomorphism of the finite plane. It is enough to prove that F^* assumes every value at no more than one point. It is easy to see that the continuity of the inverse $(F^*)^{-1}$ of F^* will follow from the continuity of one-to-one character and continuity of F^* . Let z_1 and z_2 be two points such that $F^*(z_1) = F^*(z_2)$. It follows from the above that z_1 and z_2 lie in the same half-plane bounded by $v = 0$. Since $F^*(\bar{z}) = \overline{F^*(z)}$ we may

suppose that γ_1 and γ_2 are positive

Writing down (14) in the form

$$(15) \quad \alpha(x, \gamma) = \frac{1}{\gamma} \int_x^{x+\gamma} \varphi(\xi) d\xi, \quad \beta(x, \gamma) = \frac{1}{\gamma} \int_{x-\gamma}^x \varphi(\xi) d\xi$$

we see that the mean values of φ are equal on the intervals $(x_1, x_1 + \gamma_1)$ and $(x_2, x_2 + \gamma_2)$ as well as on $(x_1 - \gamma_1, x_1)$ and $(x_2 - \gamma_2, x_2)$. Since φ is strictly increasing, one of the first mentioned intervals must be contained in the other, and the same holds for the last two. If, for example $x_2 > x_1$ it follows that $x_2 + \gamma_2 \leq x_1 + \gamma_1$ and $x_2 - \gamma_2 \leq x_1 - \gamma_1$. This implies that $z_1 = z_2$, so the mapping F^* is a homeomorphism, because local homeomorphism of the closed \mathbb{C} onto itself is the global one.

Now we show that F^* is quasiconformal in the upper half-plane. From (15) it follows that α and β are continuously differentiable and have the partial derivatives

$$\alpha_x(x, \gamma) = \frac{1}{\gamma} (\varphi(x+\gamma) - \varphi(x)), \quad \beta_x(x, \gamma) = \frac{1}{\gamma} (\varphi(x) - \varphi(x-\gamma)),$$

$$\alpha_\gamma(x, \gamma) = \frac{1}{\gamma} (\varphi(x+\gamma) - \alpha(x, \gamma)), \quad \beta_\gamma(x, \gamma) = \frac{1}{\gamma} (\varphi(x-\gamma) - \beta(x, \gamma))$$

Since φ is strictly increasing then

$$\alpha_x > 0, \quad \beta_x > 0, \quad \alpha_\gamma > 0, \quad \beta_\gamma < 0.$$

The Jacobian $J = \frac{1}{2}(\alpha_\gamma \beta_x - \alpha_x \beta_\gamma)$ of F^* is therefore positive. Hence all points of the upper half-plane are regular for F^* .

To obtain the upper bound for the dilatation quotient $p_{F^*} = (|F'_z| + |F'_{\bar{z}}|) / (|F'_z| - |F'_{\bar{z}}|)$ we write

$$p_{F^*} \leq 2 \frac{|F'_z|^2 + |F'_{\bar{z}}|^2}{|F'_z|^2 - |F'_{\bar{z}}|^2} = \frac{\alpha_x^2 + \beta_x^2 + \alpha_\gamma^2 + \beta_\gamma^2}{\alpha_\gamma \beta_x - \alpha_x \beta_\gamma} = \frac{\alpha_x \beta_x}{-\alpha_\gamma \beta_\gamma} \cdot \frac{\frac{\alpha_x}{\beta_x} + \frac{\alpha_\gamma}{\alpha_x \beta_\gamma} + \frac{\beta_x}{\alpha_x} + \frac{\beta_\gamma^2}{\alpha_x \beta_x}}{-\frac{\beta_x}{\beta_\gamma} + \frac{\alpha_x}{\alpha_\gamma}}$$

To estimate the partial derivatives of α and β we make use of the assumption (13).

From

$$\varphi(x + \frac{1}{2}\gamma) - \varphi(x) \leq g(\varphi(x + \gamma) - \varphi(x + \frac{1}{2}\gamma))$$

it follows that

$$\varphi(x + \gamma) - \varphi(x) \leq (1+g)(\varphi(x + \gamma) - \varphi(x + \frac{1}{2}\gamma)).$$

The the double inequality

$$\begin{aligned} \alpha_x(x, \gamma) > \alpha_\gamma(x, \gamma) &= \frac{1}{\gamma} \int_0^\gamma (\varphi(x + \gamma) - \varphi(x + t\gamma)) dt \geq \\ &\geq \frac{1}{2\gamma} (\varphi(x + \gamma) - \varphi(x + \frac{1}{2}\gamma)) \geq \frac{\alpha_x(x, \gamma)}{2(1+g)}. \end{aligned}$$

In the same manner we obtain

$$\beta_x(x, \gamma) > -\beta_\gamma(x, \gamma) \geq \frac{1}{2\gamma} (\varphi(x - \frac{1}{2}\gamma) - \varphi(x - \gamma)) \geq \frac{\beta_x(x, \gamma)}{2(1+g)}.$$

Finally, it follows from (13) that

$$\frac{\beta_x}{g} \leq \alpha_x \leq g\beta_x$$

and by combining these inequalities we arrive at

$$(16) \quad \rho_{F^*} \leq 8g(1+g)^2.$$

Because of the symmetry property of $F^*(\bar{z}) = \overline{F^*(z)}$ it follows that (16) holds in the lower half-plane also. By this F^* is therefore a quasiconformal mapping of the whole plane with maximal dilatation not exceeding $8g(1+g)^2$.

The bound $8g(1+g)$ was several times improved but the last known as the good (still not the best one) has form $\min\{g^{\frac{3}{2}}, 2g-1\}$ (Zehrfeld).

Quasisymmetric functions.

The condition (13) indicates that φ possesses a degree of approximate symmetry, and for this reason we refer to φ as a quasisymmetric function (mapping), when $\inf \beta =$ is called the dilatation of φ .

Qualitatively, the properties of g -qs functions parallel rather closely to those of quasiconformal mappings. However, they have some shortcomings not shared by qc mappings. We list a number of very basic properties:

- 1° 1-qs function is linear mapping,
- 2° the composition of a g -qs function with a linear function yields a g -qs function
- 3° an unfortunate feature of qs functions is that the dilatation of a function is not necessarily the same as for its inverse, nor is dilatation of a composed function bounded by product of the dilatations whereas both conditions hold for quasiconformal mappings. For example, if $\varphi(x) = x^\alpha$ ($x > 0$), then $g(\varphi) = 2^\alpha - 1$, if $\alpha > 1$ and $g(\varphi) = 1/2^\alpha - 1$ if $0 < \alpha < 1$. Hence, if $\alpha = 2$, then $g(\varphi) = 3$ and $g(\varphi^{-1}) = \sqrt{2} + 1$. Similarly $g(\varphi(\varphi)) = 15$ but $g(\varphi) \cdot g(\varphi) = 9$. But the exact bounds for $g(\varphi^{-1})$ and $g(\varphi \circ \varphi)$ are already known

4. Bojarski has pointed out that there exist $p > 1$ such that the Jacobian of each Q -qc mapping is locally p -integrable. The result is false for quasisymmetric function and says that for each $p > 1$, each $p > 1$, and each compact set E of positive measure, there exists a function φ , g -qs on the entire line, such that

$$\int_E [\varphi'(x)]^p dx = \infty$$

Sapporo, 17 June, 1989