HOKKAIDO UNIVERSITY

Title	ON THE UNIVALENCY OF CERTAIN ANALYTIC FUNCTIONS
Author(s)	Noshiro, Kiyoshi
Citation	Journal of the Faculty of Science Hokkaido Imperial University. Ser. 1 Mathematics, 2(1-2), 089-101
Issue Date	1934
Doc URL	https://hdl.handle.net/2115/55901
Type	departmental bulletin paper
File Information	JFSHIU_02_N1-2_089-101.pdf

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

ON THE UNIVALENCY OF CERTAIN ANALYTIC FUNCTIONS

By

Kiyoshi NOSHIRO

The object of this paper is to give some generalizations on and complements to Mr. J. DIEUDONNÉ's results concerning the univalency (*Schlichtheit*) of a bounded function[1]. First, assuming that $\varphi(z) = c_0 + \ldots\ldots$, $c_0 \neq 0$, is regular and $\varphi(z) < D$ in the unit circle[2], where D is a given connected domain, we obtain a general theorem on the univalency of the function $f(z) \equiv z\varphi(z)$, with the aid of Prof. KAKEYA's principle[3]. Next we state some applications of this theorem, taking some special domains as D. Finally we investigate the univalency of the function $f(z) = c_0 z + c_1 z^2 + \ldots\ldots$, c_0, c_1 given, $c_0 \neq 0$, which is regular and bounded in the unit circle: $|f(z)| < 1$.

§ I.

Let $\varphi(z) = c_0 + \ldots\ldots$, $c_0 \neq 0$, be regular and $\varphi(z) < D$ in the unit circle, where D is a given connected domain. We will here enunciate a theorem on the univalency of the function $f(z) \equiv z\varphi(z)$.

First we consider a function

$$\psi(z) = \alpha + \beta z + \ldots\ldots, \tag{1}$$

which is regular and $\psi(z) < D$ in the unit circle. Let us denote by $g(z)$ a function, which maps D conformally on the unit circle. If we

(1) J. DIEUDONNÉ: Ann. d. l'Éc. Norm. Sup., **48**, 1931. Cf. p. 349-351.

(2) The notation $\varphi(z) < D$ means that the set of values taken by $\varphi(z)$ in the unit circle belongs to the domain D.

(3) S. KAKEYA: Sci. Rep. of Tokyo Bunrika Daigaku, Section A, I, No. 14, 1931, p. 159-165.

take a branch of $g(z)$, we obtain a function $G(z)$ defined by

$$G(z) \equiv g\{\psi(z)\}, \qquad 2)$$

which is uniform, regular in the unit circle and has the following properties:

(i) $|G(z)|<1$ for $|z|<1$, (ii) $G(0)=g(\alpha)$, $G'(0)=\beta g'(\alpha)$.

Hence, applying a theorem of E. LANDAU[4], we have

$$|g(\alpha)|^2+|\beta|\,|g'(\alpha)|\leqq 1$$

or

$$|\beta|\leqq\frac{1-|g(\alpha)|^2}{|g'(\alpha)|}\equiv\Omega(\alpha, D), \qquad (3)$$

since $g'(\alpha)\neq 0$. Here we must remark that the positive quantity $\Omega(\alpha, D)$ depends only on α and D, and does not depend either on the selection of a mapping function $g(z)$ or on the selection of a branch of $g(z)$[5].

Next consider the function

$$\psi(\zeta)=\varphi\left(\frac{z_0-\zeta}{1-\bar{z}_0\zeta}\right)=\varphi(z_0)-(1-|z_0|^2)\varphi'(z_0)\zeta+\cdots, \quad (|z_0|<1). \qquad (4)$$

Since $\psi(\zeta)$ is regular and $\psi(\zeta)<D$ for $|\zeta|<1$, if we put

$$\alpha=\varphi(z_0), \quad \beta=-(1-|z_0|^2)\varphi'(z_0),$$

we obtain, from (3),

$$|\varphi'(z_0)(1-|z_0|^2)|\leqq\Omega(\varphi(z_0), D).$$

---

(4) S. KAKEYA: l. c. p. 161.

(5) Let $g_1(z)$ and $g_2(z)$ be two mapping functions. Then there exists a relation $g_2(z)=L(g_1(z))$, where $L(z)$ is a function of the form $e^{i\theta}\frac{z-\gamma}{\bar{\gamma}z-1}$, $|\gamma|<1$, and so $\frac{1-|g_1(\alpha)|^2}{|g_1'(\alpha)|}=\frac{1-|g_2(\alpha)|^2}{|g_2'(\alpha)|}$. By an entirely similar discussion, we see that $\Omega(\alpha, D)$ does not also depend on the selection of a branch of $g(z)$. Cf. BIEBERBACH: Lehrbuch der Funktionentheorie, II, 2. Aufl., p. 45–46.

Consequently, if $\varphi(z_0) \neq 0$, we have

$$\left| z_0 \frac{\varphi'(z_0)}{\varphi(z_0)} \right| \leqq \frac{|z_0|}{1-|z_0|^2} \frac{\Omega(\varphi(z_0), D)}{|\varphi(z_0)|}. \tag{5}$$

On the other hand, it is sufficient that

$$\left| z \frac{\varphi'(z)}{\varphi(z)} \right| < 1 \quad \text{for} \quad |z| < R, \tag{6}$$

to show that $f(z) \equiv z\varphi(z)$ is univalent and starshaped for $|z| < R \leqq 1$[6].

Thus we obtain a general theorem:

Theorem 1. *Let $\varphi(z) = c_0 + \ldots$, $c_0 \neq 0$, be regular and $\varphi(z) < D$ in the unit circle, where D is a given connected domain. Let $g(z)$ be an arbitrary branch of a function mapping D conformally on the unit circle, and put $\Omega(a, D) = \dfrac{1-|g(a)|^2}{|g'(a)|}$, where a is a point in D. Then $f(z) \equiv z\varphi(z)$ is univalent and starshaped with respect to the origin for $|z| < R$, provided that*

$$\frac{|z|}{1-|z|^2} \frac{\Omega(\varphi(z), D)}{|\varphi(z)|} < 1 \quad \text{for} \quad |z| < R. \tag{7}$$

§ II.

We consider a circular domain $D: |z| < M$. Since $g(z) = \dfrac{z}{M}$ is a mapping function, it is easy to see that

$$\Omega(a, D) = \frac{M^2 - |a|^2}{M} \quad \text{and} \quad \left| \frac{z\varphi'(z)}{\varphi(z)} \right| \leqq \frac{r}{1-r^2} \frac{1 - \left| \dfrac{\varphi(z)}{M} \right|^2}{\left| \dfrac{\varphi(z)}{M} \right|}$$

$$\text{for} \quad |z| = r[7], \tag{8}$$

---

(6) The following theorem is known: Suppose that $f(z) = c_0 z + \ldots$, $c_0 \neq 0$, is regular in the unit circle. Then $f(z)$ is univalent and starshaped with respect to the origin for $|z| < R$, provided that

$$\Re\left[z \frac{f'(z)}{f(z)} \right] > 0 \quad \text{for} \quad |z| < R \leqq 1.$$

Cf. A. KOBORI: Memoirs of the College of Science, Kyoto Imperial University, s. A, 1932, p. 279–291.

(7) Hereafter we often denote $|z|$ by r.

if $\varphi(z)$ does not vanish at a point z within the unit circle. Thus

Theorem 2. *Let $\varphi(z) = c_0 + \ldots$, $c_0 \neq 0$, be regular and bounded in the unit circle: $|\varphi(z)| < M$. Then $f(z) \equiv z\varphi(z)$ is univalent and starshaped with respect to the origin for $|z| < R$, if*

$$\left|\frac{\varphi(z)}{M}\right| > |z| \quad \text{for} \quad |z| < R.$$

Particularly, if we put $c_0 = 1$ in theorem 2, then we easily get

$$\frac{1-Mr}{M-r} \leqq \left|\frac{\varphi(z)}{M}\right| \leqq \frac{1+Mr}{M+r} \quad \text{for} \quad |z| = r \tag{9}$$

and so it is sufficient that r should be less than $R = M - \sqrt{M^2-1}$ in order that $\left|\frac{\varphi(z)}{M}\right| > |z|$. Thus we obtain DIEUDONNÉ's theorem:

Theorem 3. *Suppose that $f(z) = z + \ldots$. is regular and bounded in the unit circle: $|f(z)| < M$. Then $f(z)$ is univalent and starshaped with respect to the origin for $|z| < M - \sqrt{M^2-1}$. This limit can be attained by the function $f_0(z) = \dfrac{Mz(1-Mz)}{M-z}$.*

Next consider a half-plane D: $\Re[z] > 0$. Taking a mapping function $g(z) = \dfrac{1-z}{1+z}$, we have

$$\Omega(a, D) = 2\Re[a] \quad \text{and} \quad \left|z\frac{\varphi'(z)}{\varphi(z)}\right| \leqq \frac{2r}{1-r^2}\frac{\Re[\varphi(z)]}{|\varphi(z)|} \leqq \frac{2r}{1-r^2}$$

$$\text{for} \quad |z| = r. \tag{10}$$

From this we get

Theorem 4. *Let $\varphi(z)$ be regular and $\Re[\varphi(z)] > 0$ for $|z| < 1$. Then $f(z) \equiv z\varphi(z)$ is univalent and starshaped for $|z| < \sqrt{2}-1$. This limit can be attained by the function $f_0(z) = cz\dfrac{1-z}{1+z}$, $(c > 0)$.*

§ III.

We now consider a pricked (*punktiert*) circular domain D: $0<|z|<M$. A function which maps D on the unit circle is

$$g(z)=\frac{1+\log\frac{z}{M}}{1-\log\frac{z}{M}}.$$

By a short calculation, we have

$$\Omega(a, D)=2|a|\log\frac{M}{|a|} \quad \text{and} \quad \left|z\frac{\varphi'(z)}{\varphi(z)}\right|\leqq\frac{2r}{1-r^2}\log\left|\frac{M}{\varphi(z)}\right| \quad \text{for} \quad |z|=r. \tag{11}$$

Hence we easily obtain

Theorem 5. *Let* $\varphi(z)=c_0+\ldots.$ *be regular for* $|z|<1$ *and* $0<|\varphi(z)|<M$ *for* $0<|z|<1$. *Then* $f(z)\equiv z\varphi(z)$ *is univalent and starshaped with respect to the origin for* $|z|<r$, *if*

$$\left|\frac{\varphi(z)}{M}\right|\geqq e^{-\frac{1-r^2}{2r}} \quad \text{for} \quad |z|=r.$$

Put $c_0=1$ in theorem 5 and consider the function

$$F(z)=\frac{\log\frac{\varphi(z)}{M}-\log\frac{1}{M}}{\log\frac{\varphi(z)}{M}+\log\frac{1}{M}},$$

taking a branch of $\log z$ such that $\log 1=0$. Since $F(z)$ is regular and $|F(z)|<1$ in the unit circle, $F(0)$ being equal to 0, SCHWARZ's lemma gives $|F(z)|\leqq|z|$, whence we have

$$M^{-\frac{1+r}{1-r}}\leqq\left|\frac{\varphi(z)}{M}\right|\leqq M^{-\frac{1-r}{1+r}} \quad \text{for} \quad |z|=r. \tag{12}$$

Thus we can enunciate a theorem analogous to DIEUDONNÉ's.

Theorem 6. *Let* $f(z)=z+\ldots.$ *be regular for* $|z|<1$ *and* $0<|f(z)|<M$ *for* $0<|z|<1$. *Then*

(1) $f(z)$ *maps* $|z|<R=\log eM-\sqrt{(\log eM)^2-1}$ *on a starshaped domain* Δ *with respect to the origin.*

(2) *If we denote by* d *the shortest distance of the origin from the boundary of* Δ, *then* $d\geqq\left\{\log eM-\sqrt{(\log eM)^2-1}\right\}M^{-\frac{2}{\log M+\sqrt{\log M(2+\log M)}}}$.

(3) *These limits* (1) *and* (2) *are attained by the function* $f_0(z)=zM^{\frac{2e^{i\theta}z}{1+e^{i\theta}z}}$.

Proof. From (11) and (12) we have

$$\left|z\frac{\varphi'(z)}{\varphi(z)}\right|\leqq\frac{2r}{(1-r)^2}\log M \quad \text{for} \quad |z|=r. \tag{13}$$

Since the right hand side is an increasing function of r for $0<r<1$, the equation $\frac{2r}{(1-r)^2}\log M=1$ has just one root $R=\log eM-\sqrt{(\log eM)^2-1}$ there. From (12) we have $|f(Re^{i\theta})|\geqq RM^{-\frac{2R}{1-R}}$. Our theorem is completely proved, considering the function

$$f_0(z)=zM^{\frac{2e^{i\theta}z}{1+e^{i\theta}z}},$$

which has

$$f_0(-Re^{-i\theta})=-Re^{-i\theta}M^{-\frac{2R}{1-R}},\quad f_0'(-Re^{-i\theta})=\left(1-\frac{2R}{(1-R)^2}\log M\right)M^{-\frac{2R}{1-R}}=0.$$

Lastly we consider a ring-domain D: $m<|z|<M$, which can be mapped on the unit circle by a function

$$g(z)=\left(e^{i\frac{\pi}{2}\frac{\log\frac{z}{\sqrt{mM}}}{\log\sqrt{\frac{M}{m}}}}-1\right):\left(e^{i\frac{\pi}{2}\frac{\log\frac{z}{\sqrt{mM}}}{\log\sqrt{\frac{M}{m}}}}+1\right).$$

In this case our discussion gives

$$\left|z\frac{\varphi'(z)}{\varphi(z)}\right| \leqq \frac{2}{\pi}\frac{r}{1-r^2}\left(\log\frac{M}{m}\right)\cos\frac{\pi\log\frac{|\varphi(z)|}{\sqrt{mM}}}{\log\frac{M}{m}} \quad \text{for} \quad |z|=r\,.$$

Consequently we obtain

Theorem 7. *Let $\varphi(z)$ be regular and $m<|\varphi(z)|<M$ for $|z|<1$. Then $f(z)\equiv z\varphi(z)$ is univalent and starshaped for $|z|<r$ if*

$$\cos\frac{\pi}{2}\frac{\log\frac{|\varphi(z)|^2}{mM}}{\log\frac{M}{m}} \leqq \frac{\pi}{2}\frac{\frac{1-r^2}{r}}{\log\frac{M}{m}} \quad \textit{for} \quad |z|=r\,.$$

§ IV.

DIEUDONNÉ's theorem (see theorem 3) can be enunciated, with a slight modification, as follows: let $f(z)=c_0z+\ldots$, c_0 given and $\neq 0$, be regular and $|f(z)|<1$ for $|z|<1$. Then $f(z)$ is univalent and starshaped with respect to the origin for

$$|z|<\frac{1}{|c_0|}-\sqrt{\frac{1}{|c_0|^2}-1}\,.$$

It seems to me very difficult to generalize this theorem for the case where n initial coefficients $c_0, c_1, \ldots, c_{n-1}$ ($c_0\neq 0$) are given. In this paragraph we will treat the case where only two initial coefficients c_0, c_1 ($c_0\neq 0$) are given. First suppose that

$$\varphi(z)=c_0+c_1z+\ldots. \qquad (c_0,\ c_1 \text{ given},\ c_0\neq 0) \qquad (14)$$

is regular and $|\varphi(z)|<1$ for $|z|<1$. Then the function

$$g(z)\equiv\frac{\varphi(z)-c_0}{1-\bar{c}_0\varphi(z)}=\frac{c_1}{1-|c_0|^2}z+\ldots. \qquad (15)$$

is regular and $|g(z)|<1$ for $|z|<1$. Hereafter, for brevity, we put

$$\alpha_0 = \frac{c_1}{1-|c_0|^2}. \quad (8) \tag{16}$$

From (15) we have

$$\varphi(z) = \frac{c_0+g(z)}{1+\bar{c}_0 g(z)}. \tag{17}$$

Considering $|g(z)|<1$ for $|z|<1$, we have

$$\frac{|c_0|-|g(z)|}{1-|c_0||g(z)|} \leqq |\varphi(z)| \leqq \frac{|c_0|+|g(z)|}{1+|c_0||g(z)|} \tag{18}$$

for $|z|<1$. Since the expression on the left hand side decreases and that on the right increases as $|g(z)|$ increases, using a known inequality

$$|g(z)| \leqq r\frac{|\alpha_0|+r}{1+|\alpha_0| r}, \tag{19}$$

we obtain the following inequality

$$\frac{|c_0|-|\alpha_0|(1-|c_0|)r-r^2}{1+|\alpha_0|(1-|c_0|)r-|c_0| r^2} \leqq |\varphi(z)| \leqq \frac{|c_0|+|\alpha_0|(1+|c_0|)r+r^2}{1+|\alpha_0|(1+|c_0|)r+|c_0| r^2} \tag{20}$$

for $|z|=r$. In order that the left expression of (20) be positive, we have only to take r less than

$$\rho = \frac{1}{2}\left\{-|\alpha_0|(1-|c_0|)+\sqrt{|\alpha_0|^2(1-|c_0|)^2+4|c_0|}\right\}, \tag{21}$$

where ρ is the root between 0 and 1 of the equation

$$|c_0|-|\alpha_0|(1-|c_0|)r-r^2=0. \tag{22}$$

(8) Of course $|\alpha_0| \leqq 1$ and the equality holds if and only if $\varphi(z) = \dfrac{c_0+e^{i\theta}z}{1+\bar{c}_0 e^{i\theta}z}$.

Thus we see that $\varphi(z)$ does not vanish for $|z|<\rho$[9]. This limit cannot be replaced by any greater number, for the lower bound of $|\varphi(z)|$ of (20) can be attained by the function

$$\varphi_0(z)=\frac{c_0+(c_0\bar{a}_0e^{i\theta}+a_0)z+e^{i\theta}z^2}{1+(\bar{a}_0e^{i\theta}+\bar{c}_0a_0)z+\bar{c}_0e^{i\theta}z^2}$$

$$(\theta\equiv 2\operatorname{amp}c_1-\operatorname{amp}c_0+\pi\ (\text{mod. }2\pi)) \tag{23}$$

at a point $z_0=re^{i\varphi_0}$, where $\varphi_0\equiv\operatorname{amp}c_0-\operatorname{amp}c_1+\pi$ (mod. 2π), and $\varphi_0(\rho e^{i\varphi_0})=0$. From (17)

$$|\varphi'(z)|=(1-|c_0|^2)\left|\frac{g'(z)}{(1+\bar{c}_0g(z))^2}\right|. \tag{24}$$

From (8)

$$\left|\frac{g'(z)}{z}-\frac{g(z)}{z^2}\right|\leqq\frac{1-\left|\frac{g(z)}{z}\right|^2}{1-r^2}, \tag{25}$$

or

$$|g'(z)|\leqq\frac{(1-|g(z)|)(r^2+|g(z)|)}{r(1-r^2)}. \tag{26}$$

From

$$\left|\frac{z\varphi'(z)}{\varphi(z)}\right|=(1-|c_0|^2)\left|\frac{zg'(z)}{(c_0+g(z))(1+\bar{c}_0g(z))}\right|,$$

using (26), we have

$$\left|\frac{z\varphi'(z)}{\varphi(z)}\right|\leqq\frac{1-|c_0|^2}{1-r^2}\frac{(1-|g(z)|)(r^2+|g(z)|)}{(|c_0|-|g(z)|)(1-|c_0||g(z)|)}, \tag{27}$$

provided that $|c_0|>|g(z)|$.

---

(9) This result has been obtained by M. KÖSSLER: Cf. Ueber die α-Stellen von beschränkten Potenzreihen. Mémoires de la Société Royale des Sciences de Bohème, Année 1930.

Since $\frac{1-t}{|c_0|-t}\frac{r^2+t}{1-|c_0|t}$ is an increasing function of t for $0<t<|c_0|$, using the inequality (19), we obtain

$$\left|\frac{z\varphi'(z)}{\varphi(z)}\right| \leqq (1-|c_0|^2)\frac{r(|\alpha_0|+2r+|\alpha_0|r^2)}{(|c_0|-|\alpha_0|(1-|c_0|)r-r^2)(1+|\alpha_0|(1-|c_0|)r-|c_0|r^2)} \tag{28}$$

for $|z|<\rho=\frac{1}{2}\left\{-|\alpha_0|(1-|c_0|)+\sqrt{|\alpha_0|^2(1-|c_0|)^2+4|c_0|}\right\}$. (10)

Since the right hand side of (28) is an increasing function of r in the interval $(0, \rho)$, the equation

$$(1-|c_0|^2)\frac{r(|\alpha_0|+2r+|\alpha_0|r^2)}{(|c_0|-|\alpha_0|(1-|c_0|)r-r^2)(1+|\alpha_0|(1-|c_0|)r-|c_0|r^2)}=1 \tag{29}$$

or

$$c-2a(1-c)r-\left\{3-c^2+a^2(1-c)^2\right\}r^2-2a(1-c)r^3+cr^4=0, \tag{30}$$

where $c=|c_0|$, $a=|\alpha_0|=\frac{|c_1|}{1-|c_0|^2}$,

has only one root R between 0 and ρ(11).

Thus we obtain the following

Theorem 8. *Suppose that*

$$f(z)=c_0z+c_1z^2+\cdots \quad (c_0,\ c_1 \text{ given},\ c_0\neq 0)$$

is regular and $|f(z)|<1$ *for* $|z|<1$. *Then*

(10) By (19), we see that

$$|c_0|>r\frac{|\alpha_0|+r}{1+|\alpha_0|r}\geqq|g(z)| \text{ for } |z|<\rho.$$

(11) The equation (30) has 4 real roots, of which two are positive and the others negative. If we denote a negative root by τ, all roots can be written as follows: $R, \frac{1}{R}, \tau, \frac{1}{\tau}$.

(1) *$f(z)$ is univalent for $|z|<R$ and maps the circle $|z|<R$ on a starshaped domain Δ with respect to the origin, if R is the root between 0 and 1 of the equation*

$$c-2\alpha(1-c)r-\left\{3-c^2+\alpha^2(1-c)^2\right\}r^2-2\alpha(1-c)r^3+cr^4=0\,,$$

where

$$c=|c_0|\,,\qquad \alpha=\frac{|c_1|}{1-|c_0|^2}\,.$$

(2) *If we denote by d the shortest distance of the origin from the boundary of Δ, then*

$$d\geqq R\frac{c-\alpha(1-c)R-R^2}{1+\alpha(1-c)R-cR^2}\,.$$

(3) *These limits* (1) *and* (2) *can be attained by the function*

$$f_0(z)=z\frac{c_0+(c_0\bar{\alpha}_0e^{i\theta}+\alpha_0)z+e^{i\theta}z^2}{1+(\bar{\alpha}_0e^{i\theta}+\bar{c}_0\alpha_0)z+\bar{c}_0e^{i\theta}z^2}\,,$$

where $\theta\equiv 2\ \mathrm{amp}\ c_1-\mathrm{amp}\ c_0+\pi\ (\mathrm{mod.}\ 2\pi)$, $\alpha_0=\dfrac{c_1}{1-|c_0|^2}$.

Proof. The first assertion is proved above and the second follows directly from (20). Finally, by an elementary calculation, we see that

$$f_0'(Re^{i\varphi_0})=0\qquad\text{and}\qquad |f_0(Re^{i\varphi_0})|=R\frac{c-\alpha(1-c)R-R^2}{1+\alpha(1-c)R-cR^2}\,,$$

where $\varphi_0\equiv\mathrm{amp}\ c_0-\mathrm{amp}\ c_1+\pi\ (\mathrm{mod.}\ 2\pi)$.

Remark. The root R of the equation (30) can be written explicitly:

$$\begin{aligned}R=\frac{1}{2c}\Big[&\alpha(1-c)+\sqrt{(1+c)(1-c)^2\alpha^2+c(3-c)(1+c)}\\&-\Big\{(2+c)(1-c)^2\alpha^2+c(3+c)(1-c)\\&+2\alpha(1-c)\sqrt{(1+c)(1-c)^2\alpha^2+c(3-c)(1+c)}\Big\}^{\frac{1}{2}}\Big]\,.\end{aligned}$$

When c is fixed, R is a decreasing function of α in the interval (0,1) (see footnote (8)) and especially

$$R = \frac{1}{c} - \sqrt{\frac{1}{c^2} - 1}, \qquad \text{if} \quad \alpha = 1,$$

$$R = \frac{1}{2}\left\{\sqrt{\frac{(3-c)(1+c)}{c}} - \sqrt{\frac{(3+c)(1-c)}{c}}\right\}, \quad \text{if} \quad \alpha = 0.$$

§ V.

In this paragraph we shall make some remarks on the univalency of certain analytic functions. Suppose that $f(z) = \alpha + \beta z + \ldots$. ($\alpha$, β given and $\beta \neq 0$) is regular and $f(z) < D$ in the unit circle, where D is a given *simply* connected domain. It is well known that there exists one and only one function $\varphi(z)$ which maps D in a one-to-one and conformal manner on a circle under the condition: $\varphi(\alpha) = 0$, $\varphi'(\alpha) = \frac{1}{\beta}$. We denote the radius of the mapped circle by M. Next consider the function $F(z) = \varphi(f(z))$. Then $F(z) = z + \ldots$. is regular and bounded in the unit circle: $|F(z)| < M$. Here we can, without difficulty, assert the following:

1°. The existence of $F(z)$ follows from that of $f(z)$, and conversely. Hence, it is necessary and sufficient that $M \geqq 1$[12] for the existence of $f(z)$.

2°. If $f(z)$ takes the same value at n given points $z_1, z_2, \ldots, z_n$ within the unit circle, then $F(z)$ also takes the same value with the same multiplicity, its converse being also true.

Hence, as an immediate result from a theorem due to DIEUDONNÉ[13], we can state the following theorem: *$f(z)$ assumes at most p times*

(12) The equality holds if and only if $f(z) = \varphi^{-1}(z)$ where φ^{-1} is an inverse function of φ.

(13) DIEUDONNÉ has proved that $F(z)$ assumes at most p times the same value for $|z| < \rho_p$ and ρ_p can be attained. Cf. l.c. p. 348. Or my paper: Proc. of Phys.-Math. Soc. of Japan, 3rd Ser., Vol. 14, 1932, p. 304-309. Cf. p. 308.

the same value for $|z|<\rho_p$, *where* ρ_p *is the root between* 0 *and* 1 *of the equation* $1+r^2+r^4+\ldots+r^{2p}-(p+1)Mr^p=0$ *and moreover* ρ_p *cannot be replaced by any greater number.*

For example, we consider a half-plane D: $\Re[z]>0$. Without difficulty, we have

$$\varphi(z)=\frac{2\Re[\alpha]}{\beta}\,\frac{z-\alpha}{z+\bar{\alpha}} \qquad \text{and} \qquad M=\frac{2\Re[\alpha]}{|\beta|}.$$

Hence, it is necessary and sufficient that $2\Re[\alpha]\geqq|\beta|$ for the existence of $f(z)$. By the theorem above we get at once

Theorem 9.[14] *Let* $f(z)=\alpha+\beta z+\ldots.$ (α, β *given and* $\beta\neq 0$) *be regular and* $\Re[f(z)]>0$ *for* $|z|<1$. *Then* $f(z)$ *is univalent for*

$$|z|<\rho_1=\frac{2\Re[\alpha]}{|\beta|}-\sqrt{\left(\frac{2\Re[\alpha]}{|\beta|}\right)^2-1},$$

and the limiting case can be attained by the function

$$f_0(z)=\frac{Mae^{-i\theta}+(\bar{\alpha}-e^{-i\theta}\alpha)z-\bar{\alpha}Mz^2}{Me^{-i\theta}-(1+e^{-i\theta})z+Mz^2} \qquad \left(\theta=\operatorname{amp}\beta,\quad M=\frac{2\Re[\alpha]}{|\beta|}\right),$$

which has a derivative vanishing at a point on the circle $|z|=\rho_1$.

March, 1933.
Mathematical Institute
Hokkaido Imperial University.

(14) Compare with theorem 4 in KOBORI's paper: Memoirs of the College of Science, Kyoto Imperial University, Ser. A. Vol. XIV, 1931, p. 251–255.