HOKKAIDO UNIVERSITY

Title	ON *-MODULAR RIGHT IDEALS OF AN ALTERNATIVE RING
Author(s)	Hashimoto, Hiroyoshi
Citation	Journal of the Faculty of Science Hokkaido University. Ser. 1 Mathematics, 15(1-2), 131-133
Issue Date	1960
Doc URL	https://hdl.handle.net/2115/56013
Type	departmental bulletin paper
File Information	JFSHIU_15_N1-2_131-133.pdf

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

ON $*$-MODULAR RIGHT IDEALS OF AN ALTERNATIVE RING

By

Hiroyoshi HASHIMOTO

It is well known that, in any alternative ring A, the Smiley radical SR(A) is contained in every modular maximal right ideal M. E. Kleinfeld has shown that every primitive alternative, non-associative ring is a Cayley-Dickson algebra.

Now we introduce the notion of $*$-modularity as follows: a right ideal I of an alternative ring A is called $*$-modular if there exist two elements a, $u \in A$ such that

$$x+ax+(a, x, u) \in I \tag{1}$$

for all $x \in A$, where (a, x, u) denotes the associator $ax \cdot u - a \cdot xu$ of a, x, u, and in this case we call a a *left $*$-modulo unit of I*. Clearly, modularity implies $*$-modularity.

In this note, we shall show that the above results are also true if we replace modular ideals by $*$-modular ideals.

If a ring A is assumed to be alternative, then (a, b, c) becomes a skew-symmetric function of its three variables.

The Smiley radical SR(A) of an alternative ring A is defined as the totality of elements $z \in A$ for which each element of $(z)_r$ is right quasi-regular.

In the next lemma we develop an important property of $*$-modular right ideals.

Lemma 1. *Let I^* be a $*$-modular right ideal of an alternative ring A, and suppose that a left $*$-modulo unit a of I^* is right quasi-regular. Then $I^*=A$.*

Proof. Let b be a right quasi-inverse of a:

$$a+b+ab=0\,. \tag{2}$$

Since a is a left $*$-modulo unit of I^* and since $(a, a, u)=0$, we have $a+a^2 \in I^*$ by putting $x=a$ in (1), while $(a+a^2)b-(a, b, u)=ab+a^2b-(a, a, u)-(a, b, u)=ab+a^2b-(a, a+b, u)=ab+a^2b+(a, ab, u) \in I^*$ by (2). Hence it follows that $(a, b, u) \in I^*$. On the other hand, if we put $x=b$ in (1), we

have $b+ab+(a, b, u)\in I^*$, whence $b+ab\in I^*$. Thus $a\in I^*$, which implies together with (1) that every $x\in A$ is in I^*, that is, $I^*=A$.

As any modular maximal right ideal is a member of a set of $*$-modular maximal right ideals, the intersection of all the modular maximal right ideals is contained in the intersection of modular maximal right right ideals.

Now, we show a connection between the intersection of all the $*$-modular maximal right ideals and the radical $\mathrm{SR}(A)$ in an alternative ring A.

Theorem 1. *Let A be an alternative ring. Then the Smiley radical $SR(A)$ is contained in the intersection of all the $*$-modular maximal right ideals M^*:*

$$\mathrm{SR}(A)\subseteq \bigcap M^*.$$

Proof. Let $z\in A$ be an element not contained in the intersection $\bigcap M^*$. Then there exists a $*$-modular maximal right ideal M^* does not contain z. And we have $A=M^*+(z)_r$. Let a, u be elements such that $x+ax+(a, x, u)\in M^*$ for all $x\in A$, and let m^* and z' be elements of M^* and $(z)_r$ respectively such that $a=m^*+z'$. Then $x+z'x+(z', x, u)=x+(a-m^*)x+(a-m^*, x, u)=x+ax+(a, x, u)-m^*x+(m^*, x, u)\in M^*$ for all $x\in A$. Thus z' is also a left $*$-modulo unit of M^*. But, since $M^*\neq A$, z' is not right quasi-regular by Lemma 1, and so z' is not in $\mathrm{SR}(A)$. This proves our theorem.

Next we refer to the structure of $*$-primitive alternative ring.

An alternative ring is defined to be $*$-primitive in case it contains a $*$-modular maximal right ideal whose quotient is zero.

Lemma 2. *The quotient $(I^*: A)=\{x\in A;\ Ax\subseteq I^*\}$ of a $*$-modular right ideal I^* is an ideal of A.*

Proof. The $*$-modularity of I^* assures the existence of $a, u\in A$ with the property that $x+ax+(a, x, u)\in I^*$ for every $x\in A$. Since $ax\in I^*$ for $x\in(I^*: A)$, we have $x+(a, x, u)=x+a\cdot ux-au\cdot x\in I^*$. And further $ux+a\cdot ux+(a, ux, u)\in I^*$. Combining this with $ux\in I^*$, we obtain $a\cdot ux\in I^*$ and eventually $x\in I^*$. Hence, for any $y\in A$, $A\cdot xy$ and $A\cdot yx$ are both in I^*.

By the light of this lemma, it is clear that A is $*$-primitive if and only if A has a $*$-modular maximal right ideal which contains no nonzero two-sided ideals.

An alternative ring is called simple if it has no nonzero proper two-sided ideals and is not a nil ring.

The following lemma is due to E. Kleinfeld [1].

Lemma 3. *A simple alternative ring is either a Cayley-Dickson algebra or associative.*

Most results in primitive alternative rings which were stated in [2] are also true in our ∗-primitive case under a slight modification of the modularity.

We obtain the following:

Theorem 2. *Every ∗-primitive, alternative, non-associative ring A is a Cayley-Dickson algebra.*

Proof. We may prove, with the help of the proof in primitive case [2], that every ∗-modular maximal right ideal M^* of A is zero. And so, it is enough only to show that A is a simple alternative ring. For any left ∗-modulo unit a of M^*, $a+a^2=0$ and then $a^n=\pm a$ for every integer $n\geq 1$. It shows that a is not nilpotent, and hence A is simple by Lemma 3. Therefore, A is a Cayley-Dickson algebra.

Bibliography

[1] E. KLEINFELD, Simple alternative ring, Ann. of Math. vol. 58 (1953), pp. 544–547.
[2] ———, Primitive alternative ring and semi-simplicity, Amer. J. of Math. vol. 77 (1955), pp. 725–730.
[3] M. F. SMILEY, The radical of an alternative ring, Ann. of Math. vol. 49 (1948), pp. 702–709.

Hokkaido University