HOKKAIDO UNIVERSITY

Title	ON A SIMPLE RING WITH A GALOIS GROUP OF ORDER p^{e}
Author(s)	Takazawa, Takao; Tominaga, Hisao
Citation	Journal of the Faculty of Science Hokkaido University. Ser. 1 Mathematics, 15(3-4), 198-201
Issue Date	1961
Doc URL	https://hdl.handle.net/2115/56016
Type	departmental bulletin paper
File Information	JFSHIU_15_N3-4_198-201.pdf

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

ON A SIMPLE RING WITH A GALOIS GROUP OF ORDER p^e

By

Takao TAKAZAWA and Hisao TOMINAGA

Recently in [2, §3],[1] the next was obtained: *Let R be a simple ring (with minimum condition) of characteristic $p\neq 0$, and $\mathfrak{G}$ a DF-group of order p^e. If $S=J(\mathfrak{G}, R)$, then $[R:S]$ divides p^e, and $V_R(S)$ coincides with the composite of the center of R and that of S.* More recently, in [1], M. Moriya has proved the following: *Let R be a division ring, $\mathfrak{G}$ an automorphism group[2] of order p^e (p a prime), and $S=J(\mathfrak{G}, R)$. If the center of S contains no primitive p-th roots of 1, then $[R:S]$ divides p^e, and $V_R(S)$ coincides with the composite of the center of R and that of S. And moreover, $[R:S]$ is equal to p^e provided R is not of characteristic p.* The purpose of this note is to extend these facts to simple rings in such a way that our extension contains also the fact cited at the beginning.

In what follows, we shall use the following conventions: R is a simple ring with the center C, and $\mathfrak{G}$ a DF-group of order p^e where p is a prime number. We set $S=J(\mathfrak{G}, R)$, which is a simple ring by [2, Lemma 2]. And by Z and V we shall denote the center of S and the centralizer $V_R(S)$ of S in R respectively. Finally, as to notations and terminologies used here, we follow [2].

Now, we shall begin our study with the following theorem.

Theorem 1. *If Z contains no primitive p-th roots of 1, then $[R:S]$ diviees p^e.*

Proof. Firstly, in case $e=1$, $\mathfrak{G}$ is either outer or inner. If $\mathfrak{G}$ is outer, then it is well-known that there holds $[R:S]=p$. Thus, we may, and shall, assume that $\mathfrak{G}$ is inner, and set $\mathfrak{G}=\{1, \tilde{v}, \cdots, \tilde{v}^{p-1}\}$. Then, to be easily seen, v is contained in $Z(\supseteq C)$, and $v^p=c$ for some $c\in C$. If the polynomial $X^p-c\in C[X]$ is reducible, then it possesses a linear factor, that is, there exists an element $c_0\in C$ such that $c_0^p=c$, whence it follows that

1) Numbers in brackets refer to the references cited at the end of this note.

2) One may remark here that in case R is a division ring any automorphism group of finite order becomes naturally a DF-group.

$(vc_0^{-1})^p=1$. Recalling here $vc_0^{-1}\in Z$, we obtain $vc_0^{-1}=1$. But this contradicts $\tilde{v}\neq 1$. Consequently, we see that X^p-c is irreducible in $C[X]$, and so $V=C[v]$ yields at once $p=[V:C]=[R:S]$. Now we proceed with induction for e, and assume $e>1$. Take a subgroup $\mathfrak{P}$ of order p which is contained in the center of $\mathfrak{G}$, and set $P=J(\mathfrak{P}, R)$. Then, by [2, Lemma 3], $\mathfrak{P}$ is also a DF-group and $V_P(S)$ is a division ring of finite dimension over $V_P(P)$. Hence, $\mathfrak{G}\mid P$(=the restriction of $\mathfrak{G}$ to P) is a DF-group whose order is a divisor of p^{e-1}. And so, by our induction hypothesis, $[P:S]$ is a divisor of p^{e-1}. Further, noting that $J(\mathfrak{G}\mid V_P(S), V_P(S))=Z$ and the order of $\mathfrak{G}\mid V_P(S)$ is a divisor of p^{e-1}, we see that $[V_P(S):Z]$ is a divisor of p^{e-1} again by our induction hypothesis. Accordingly, it follows that $V_P(S)$, so that $V_P(P)$ contains no primitive p-th roots of 1. Combining this with the fact that $\mathfrak{P}$ is a DF-group of order p, we obtain $[R:P]=p$. Hence, $[R:S]=[R:P]\cdot[P:S]$ is a divisor of p^e.

Lemma 1. *If Z contains no primitive p-th roots of 1, then $S\neq C$ provided $e>0$.*

Proof. If, on the contrary, $S=C$ then R is a division ring necessarily and $\mathfrak{G}$ is inner. Now, choose a subgroup $\mathfrak{P}=\{1, \tilde{v}, \cdots, \tilde{v}^{p-1}\}$ of order p contained in the center of $\mathfrak{G}$. Then, for each $\sigma=\tilde{u}\in\mathfrak{G}$, $\tilde{v}\sigma=\sigma\tilde{v}$ implies $v\sigma=vc_\sigma$ with some $c_\sigma\in C\subseteq Z$. And $v^p=uv^pu^{-1}=(v\sigma)^p=v^pc_\sigma^p$ yields $c_\sigma^p=1$, i. e. $c_\sigma=1$. This means evidently $v\in S=C$. But this is a contradiction.

Theorem 2. *If Z contains no primitive p-th roots of 1, then V is the conposite $C[Z]$ of C and Z.*

Proof. Since the order of $\mathfrak{G}\mid V$ is a divisor of p^e and $J(\mathfrak{G}\mid V, V)=Z$, $[V:Z]$ divides p^e by Theorem 1. We see therefore that V contains no primitive p-th roots of 1. For the subgroup $\mathfrak{H}=\tilde{V}$ of $\mathfrak{G}$, the order of $\mathfrak{H}\mid V$ is a divisor of p^e and $J(\mathfrak{H}\mid V, V)$ coincides with the center Z_0 of V. And so, by Lemma 1, $\mathfrak{H}\mid V=1$, that is, V is a field. (If $e=0$, then $V=C$ evidently.) Finally, suppose $V\supsetneq C[Z]$. Since $V=V(\mathfrak{G})$ (=the subring generated by all regular elements $v\in R$ with $\tilde{v}\in\mathfrak{G}$), $\mathfrak{G}$ contains an inner automorphism determined by an element v not contained in $C[Z]$. Then evidently $v^{p^d}=c$ for some $d>0$ and $c\in C$. Since V is Galois and finite over $C[Z]$, and so, since the field V is normal and separable over the subfield $C[Z]$, there exists an element $u\in V$ different from v such that $u^{p^d}=v^{p^d}$, i. e. $(vu^{-1})^{p^d}=1$. Recalling here V does not contain primitive p-th roots of 1, we have $vu^{-1}=1$, i. e. $u=v$. But this is a contradiction. We have proved therefore $V=C[Z]$.

Now, combining Theorem 2 with [3, Theorem 1.1] and [3, Theorem 3.1], we obtain the next at once.

Corollary 1. *If Z contains no primitive p-th roots of 1, then each intermediate ring T of R/S is a simple ring and $T=S[t]$ with some t.*

Theorem 3. *If Z contains no primitive p-th roots of 1, and S is not of characteristic p, then $[R:S]$ coincides with p^e.*

Proof. At first, it may be noted that the characteristic of S is different from 2. If $e=1$, then our assertion has been shown in the proof of Theorem 1. We shall proceed again by induction for e. Take a subgroup $\mathfrak{P}$ of order p which is contained in the center of $\mathfrak{G}$, and set $P=J(\mathfrak{P}, R)$. Then, as is cited in the proof of Theorem 1, $\mathfrak{P}$ and $\mathfrak{G}\,|\,P$ are DF-groups of R and P respectively, and $V_P(P)$ contains no primitive p-th roots of 1. Thus, by our induction hypothesis, it follows that $[R:S]=[R:P]\cdot[P:S]=p\cdot$(order of $\mathfrak{G}\,|\,P$). In what follows, we shall prove that $\mathfrak{G}(P)=\{\sigma\in\mathfrak{G}\,;\,x\sigma=x \text{ for all } x\in P\}$ coincides with $\mathfrak{P}$, which enables us evidently to complete our proof. Since in case $\mathfrak{P}$ is outer there is nothing to prove, we shall restrict our proof to the case where $\mathfrak{P}$ is inner: $\mathfrak{P}=\{1, \bar{v}, \cdots, \bar{v}^{p-1}\}$. Since R/P is evidently inner Galois, each element of $\mathfrak{G}(P)$ is an inner automarphism. If $\bar{u}\neq 1$ is in $\mathfrak{G}(P)$, then $u^{p^d}=c'$ with some $d>0$ and $c'\in C$. Recalling that the field $V_R(P)=C[v]$ is of dimension p over C, u possesses a minimal polynomial $f(x)=X^p+\cdots+c_p\in C[X]$. If ζ is a primitive p^d-th root of 1 (contained in a suitable extension field of V), then $\{u\zeta^i\,;\,i=0, \cdots, p^d-1\}$ exhausts the roots of $X^{p^d}-c'=0$. Hence, noting that $f(X)$ divides $X^{p^d}-c'$ in $C[X]$, we obtain $-c_p=u^p\zeta^j$ with some j. Since, as is noted in the proof of Theorem 2, $V_R(P)(\subseteq V)$ contains no primitive p-th roots of 1, $\zeta^i=-c_pu^{-p}\in V_R(P)$ yields at once $u^p=-c_p\in C$. Consequently, by [1, Hilfssatz 4], it will be seen that $u=v^kc$ with some integer k and $c\in C$, which shows that $\bar{u}=\bar{v}^k\in\mathfrak{P}$.

As a direct consequence of Theorem 3 and [2, Theorem 4], we obtain the following:

Corollary 2. *If Z contains no primitive p-th roots of 1, and S is not of characteristic p, then R/S possesses a $\mathfrak{G}$-normal basis element, that is, there exists an element $r\in R$ such that $R=\sum_{\sigma\in\mathfrak{G}}\oplus(r\sigma)S$.*

References

[1] M. MORIYA: Zur Galoisschen Theorie der Schiefkörper, Math. J. Okayama Univ., 9 (1959), 49–62.

[2] T. NAGAHARA, T. ONODERA and H. TOMINAGA: On normal basis theorem and strictly Galois extensions, Math. J. Okayama Univ., 8 (1958), 133–142.

[3] T. NAGAHARA and H. TOMINAGA: On Galois and locally Galois extensions of simple rings, Math. J. Okayama Univ., 10 (1961) to appear.

Department of Mathematics,
Hokkaido University

(Received September 19, 1960)