HOKKAIDO UNIVERSITY

Title	NOTE ON DECOMPOSITION SETS OF SEMI-PRIME RINGS
Author(s)	Kishimoto, Kazuo; Kurata, Yoshiki
Citation	Journal of the Faculty of Science Hokkaido University. Ser. 1 Mathematics, 16(3-4), 214-220
Issue Date	1962
Doc URL	https://hdl.handle.net/2115/56029
Type	departmental bulletin paper
File Information	JFSHIU_16_N3-4_214-220.pdf

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

NOTE ON DECOMPOSITION SETS OF SEMI-PRIME RINGS

By

Kazuo KISHIMOTO and Yoshiki KURATA

Introduction. As has been observed by Jacobson the set $\mathfrak{P}=\mathfrak{P}(A)$ of all primitive ideals of a ring A may be made into a topological space endowed with Stone's topology, and recently, concerning topological properties of the structure space, Suliński [8] obtained some structure theorems of a semi-simple ring which is represented as a subdirect sum of simple rings with unity.

In this note, we shall extend his results to semi-prime rings and give necessary and sufficient conditions for a semi-prime ring to have a minimal decomposition set.

§ 1. First of all, we shall prove the following extension of [1, Theorem 1].

Lemma 1. *Let T be an ideal of a ring A.*

(1) *If p is a prime ideal of A then $T\frown p$ is a prime ideal of the ring T and if moreover p does not contain T then $(p\frown T: T)$*[1] $=p$.

(2)[2] *If p_1 is a prime ideal of the ring T, then there exists a prime ideal p of A such that $p\frown T=p_1$ and, if $p_1\neq T$, then $(p_1: T)=p$.*

Proof. (1) By [6, Lemma 2], $T\frown p$ is a prime ideal of the ring T. Assume that p does not contain T. Then $T\cdot(p\frown T: T)\subseteqq p$ implies $(p\frown T: T)\subseteqq p$ and hence we have $(p\frown T: T)=p$.

(2) Let B be the ideal of A generated by p_1 and let x be an arbitrary element of $B\frown T$. Since $xTxTx\subseteqq TBT\subseteqq p_1$ and p_1 is a prime ideal in T, x belongs to p_1, and hence $T\frown B=p_1$. The complement C of p_1 in T is an *m-system* (in T whence) in A and does not meet B. By Zorn's lemma, there exists a prime ideal p of A containing B such that p does not meet C and satisfies $T\frown p=p_1$. Moreover, if $p_1\neq T$ then p can not contain T, and hence, by (1), we have $(p_1: T)=p$.

A ring A is called *a semi-prime ring* if it is isomorphic to a subdirect sum of prime rings, i.e., if there exist prime ideals p_α $(\alpha\in\Lambda)$ of

1) We shall denote by $(p\frown T: T)$ the set $\{a\in A;\ Ta\subseteqq p\frown T\}$.

2) Cf. [3] and [7].

A such that $\bigcap_{\alpha\in\Lambda} p_\alpha=0$.

As is easily seen, the annihilator[3] of a non-zero ideal in a semi-prime ring is always represented as the intersection of all prime ideals which contain the annihilator. However, we have

Corollary 1. *A non-zero ideal T of a semi-prime ring A is a prime ring if and only if the annihilator $(0: T)$ is a prime ideal in A.*

Let A be an arbitrary ring and let $\mathfrak{Q}=\mathfrak{Q}(A)$ be the set of all prime ideals of A other than A. For any non-empty subset $\mathfrak{N}$ of $\mathfrak{Q}$, we define the closure $\overline{\mathfrak{N}}$ of $\mathfrak{N}$ as the totality of those prime ideals p in $\mathfrak{Q}$ which contains $I(\mathfrak{N})$, where $I(\mathfrak{N})$ denotes the intersection of all prime ideals belonging to $\mathfrak{N}$. $\mathfrak{Q}$ becomes a topological space relative to this closure operation $\mathfrak{N}\to\overline{\mathfrak{N}}$, and is called the *structure space* of the ring A.

For the lower radical $R=I(\mathfrak{Q})$ of A, we set $T^*=(R: T)$ for any ideal T of A. If A is semi-prime, then the lower radical R of A is equal to 0 and hence T^* coincides with the right annihilator $r(T)$ of T as well as the left annihilator $l(T)$ of T.

Lemma 2. *Let A be a ring. Then, for any subset $\mathfrak{N}$ of $\mathfrak{Q}$, we have $I(\mathfrak{N})^*=I(\mathfrak{Q}-\overline{\mathfrak{N}})$*[4].

In particular, we have $I(\mathfrak{N})^*\frown I(\mathfrak{N})=R$.

Proof. $I(\mathfrak{N})\cdot I(\mathfrak{Q}-\overline{\mathfrak{N}})\subseteqq I(\mathfrak{N})\frown I(\mathfrak{Q}-\overline{\mathfrak{N}})=I(\overline{\mathfrak{N}})\frown I(\mathfrak{Q}-\overline{\mathfrak{N}})=I(\mathfrak{Q})=R$. Conversely, for any prime ideal $p\in\mathfrak{Q}-\overline{\mathfrak{N}}$, we have $I(\mathfrak{N})\cdot I(\mathfrak{N})^*\subseteqq R\subseteqq p$ and hence $I(\mathfrak{N})^*\subseteqq p$, thus $I(\mathfrak{N})^*\subseteqq I(\overline{\mathfrak{Q}}-\mathfrak{N})$.

Lemma 3. *Let A be a ring and let p be in $\mathfrak{Q}$. Then the following conditions are equivalent:*

(1) $p^*\neq R$.

(2) $p^{**}=p$.

(3) $\overline{\{p\}}$ *contains a non-empty open subset $\mathfrak{N}$ of $\mathfrak{Q}$.*

Moreover, if this is the case, p is a minimal prime ideal of A.

Proof. (1)$\rightleftarrows$(2). Assume that $p^*\neq R$. Then $p^*\nsubseteqq p$ and, since $p^*p^{**}\subseteqq R\subseteqq p$, we have $p^{**}\subseteqq p$ and hence $p^{**}=p$. Conversely, assume that $p^{**}=p$ and $p^*=R$. Then $p=A$, a contradiction.

(1)$\rightleftarrows$(3). $p^*\neq R$ means $\overline{\mathfrak{Q}-\{p\}}\neq\mathfrak{Q}$. Thus, $\mathfrak{N}=\mathfrak{Q}-\overline{\mathfrak{Q}-\{p\}}$ $(\subseteqq\overline{\{p\}})$ is

3) In a semi-prime ring, the right annihilator $r(T)$ of any ideal T coincides with its left annihilator $l(T)$.

4) We shall denote by $\mathfrak{Q}-\overline{\mathfrak{N}}$ the set theoretical complement of $\overline{\mathfrak{N}}$ in $\mathfrak{Q}$.

open. Conversely, let $\mathfrak{N}$ be a non-empty open subset of $\overline{\{p\}}$. Then $p^* = I(\Omega - \overline{\{p\}}) \supseteq I(\Omega - \mathfrak{N}) \supsetneqq R$ because $\Omega - \mathfrak{N}$ is closed. Thus $p^* \neq R$.

Now assume that $p^* \neq R$ and let p_1 be a prime ideal of A such that $p_1 \subsetneqq p$. Since $p_1 \notin \overline{\{p\}}$, $p^* = I(\Omega - \overline{\{p\}}) \subseteq p_1$, and hence $p^* \subseteq p$, which is a contradiction.

Corollary 2. *Let A be a semi-prime ring and let p be a prime ideal in A such that $p^* \neq 0$. Then p^* is a prime ring, and is maximal in the set of those ideals of A which are prime as ring.*

Proof. From Lemma 3 and Corollary 1, p^* is a prime ring. Let T be an ideal in A which is prime as a ring and $T \supsetneqq p^*$. Then $T \frown p$ and p^* are non zero ideals in the prime ring T and $(T \frown p) \cdot p^* = 0$. This is a contradiction.

Lemma 4. *Let A be a ring and let $\mathfrak{N} = \{p_\alpha\}_{\alpha \in \Lambda'}$ be a set of different minimal prime ideals in A. If $I(\mathfrak{N}) = 0$ then $r(p_\alpha) = l(p_\alpha) = I(\mathfrak{N} - \{p_\alpha\})$ for each $\alpha \in \Lambda'$.*

Proof. Let p_α be in $\mathfrak{N}$. Then for each p_β in $\mathfrak{N}$, we have either $p_\alpha \subseteq p_\beta$ or $r(p_\alpha) \subseteq p_\beta$. Since p_β is a minimal prime ideal in A, $r(p_\alpha) \subseteq p_\beta$ for all p_β with $\beta \neq \alpha$. Therefore $r(p_\alpha)$ ($\subseteq$whence) $= I(\mathfrak{N} - \{p_\alpha\})$. Similarly, we have $l(p_\alpha) = I(\mathfrak{N} - \{p_\alpha\})$.

§ 2.

Definition 1. *Let A be a ring. We shall denote by $\mathfrak{D}$ the set of all prime ideals $p \in \Omega$ such that $p^* \neq R$, and call it the decomposition set for A.*

Definition 2. *Let A be a semi-prime ring. A subset $\mathfrak{N}$ of Ω will be called a minimal decomposition set for A if $I(\mathfrak{N}) = 0$ and $I(\mathfrak{N} - \{p\}) \neq 0$ for all p in $\mathfrak{N}$* (Goldie [1]).

In [4, Theorem 3], one of the present authors proved that a semi-prime ring has at most one minimal decomposition set for A, and, if it exists, it should coincide with $\mathfrak{D}$.

Now we shall give necessary and sufficient conditions for a semi-prime ring to have a minimal decomposition set.

Theorem 1. *If A is a semi-prime ring, then the following conditions are equivalent:*

(1) *There exists a minimal decomposition set $\mathfrak{M}$ for A.*

(2) *Every non-zero ideal T of A contains a non-zero ideal B of the ring T which is prime as a ring.*

(3) *The annihilator of the ideal generated by all those non-zero*

ideals of A which are prime as ring is zero.

(4) *There exists a subset $\mathfrak{N}$ of $\mathfrak{D}$ such that $I(\mathfrak{N})=0$.*

Proof. (1)$\rightarrow$(2). Let T be any non-zero ideal of A. There exists a prime ideal p in $\mathfrak{M}$ such that $T \not\subseteq p$. Then $T \cdot p^*$ is a non-zero ideal of the ring T. For otherwise, $p^* \neq R$ and so $p^{**}=p$ by Lemma 3, which would imply $T \subseteq p$. Besides, $T \cdot p^*$ is prime as a ring by Lemma 1 (1) because of $(T \cdot p^*) \cap p \subseteq (T \cap p^*) \cap p = T \cap (p^* \cap p) = 0$.

(2)$\rightarrow$(3). It is easily seen, by Corollaries 1 and 2, that the ideal generated by all those non-zero ideals of A which are prime as ring coincides with the ideal $\sum p^*$ generated by all p^* with $p \in \mathfrak{D}$. Now $(\sum p^*)^* = \bigcap p^{**} = \bigcap p = I(\mathfrak{D})$ by Lemma 3.

Next, suppose that $I(\mathfrak{D}) \neq 0$. Then, by our assumption, there exists a non-zero ideal B of the ring $I(\mathfrak{D})$ which is prime as a ring. By Lemma 1 (2), there exists a prime ideal $p \notin \mathfrak{D}$ such that $0=(p \cap I(\mathfrak{D})) \cap B = p \cap B$. B contains a non-zero ideal B' of A by [2, Proposition IV. 3.2]. Since $B' \cap p = 0$, we have $p^* \supseteq B' \neq 0$, which contradicts $p \notin \mathfrak{D}$.

(3)$\rightarrow$(4). This is clear by the proof of (2)$\rightarrow$(3).

(4)$\rightarrow$(1). Since every prime ideal p belonging to $\mathfrak{D}$ is a minimal prime ideal by Lemma 3, Lemma 4 yields our implication.

Corresponding to [8, Theorem 5], we have

Theorem 2. *Let A be a semi-prime ring and let T be a non-zero ideal of A. Then we have $I(\mathfrak{D}_T)=I(\mathfrak{D}) \cap T$, where $\mathfrak{D}_T$ denotes the decomposition set for the ring T.*

Proof. Let $\mathfrak{N}$ be the set of all p in $\mathfrak{Q}$ such that $p \supseteq T$. Then we have $I(\mathfrak{D}) \cap T = I(\mathfrak{D}-(\mathfrak{D} \cap \mathfrak{N})) \cap I(\mathfrak{D} \cap \mathfrak{N}) \cap T = I(\mathfrak{D}-(\mathfrak{D} \cap \mathfrak{N})) \cap T$. Now assume that $p' \in \mathfrak{D}-(\mathfrak{D} \cap \mathfrak{N})$ and $(T \cap p')^* \cap T = 0$. Then $(T \cap p')^* \subseteq p'$ because $p' \not\supseteq T$ contradicting $p' \in \mathfrak{D}$. Hence, $(T \cap p')^* \cap T \neq 0$ and we have $T \cap p' \in \mathfrak{D}_T$. Thus $I(\mathfrak{D}) \cap T = I(\mathfrak{D}-(\mathfrak{D} \cap \mathfrak{N})) \cap T \supseteq I(\mathfrak{D}_T)$.

Conversely, let p_1 be in $\mathfrak{D}_T$. Then there exists, by Lemma 1 (2), a prime ideal p in $\mathfrak{Q}$ such that $T \cap p = p_1$ and $(T \cap p)^* \cap T \neq 0$. Since $(((T \cap p)^* \cap T) \cap p)^2 = ((T \cap p)^* \cap (T \cap p))^2 \subseteq (T \cap p)^* \cdot (T \cap p) = 0$, $((T \cap p)^* \cap T) \cap p = 0$ and hence $(T \cap p)^* \cap T \subseteq p^*$. Thus $p^* \neq 0$, showing that $I(\mathfrak{D}_T)$ contains $T \cap I(\mathfrak{D})$. This completes our proof.

As a corollary of Theorem 2, we have the following second necessary and sufficient condition for a semi-prime ring to have a minimal decomposition set.

Corollary 3. *A semi-prime ring A has a minimal decomposition set*

if and only if A has an ideal T such that $T^*=0$ and T has a minimal decomposition set.

Proof. Let $\mathfrak{M}$ be a minimal decomposition set for A and let $T=\sum p_\alpha^*$ with $p_\alpha\in\mathfrak{M}$. Then $T^*=0$ by Theorem 1 and for $\alpha\neq\beta$, $p_\alpha^*\frown p_\beta^*\subseteqq p_\alpha^*\frown p_\alpha=0$ since $p_\beta^*=I(\mathfrak{M}-\{p_\beta\})\supseteqq p_\alpha$ by Lemma 4. Thus for each α, $T=p_\alpha^*\oplus T_\alpha$ with $T_\alpha=\sum_{\beta\neq\alpha}p_\beta^*$. Moreover, $\bigcap T_\alpha\subseteqq\bigcap p_\alpha=I(\mathfrak{M})=0$. Hence, T is isomorphic to a special subdirect sum of p_α^* with $p_\alpha\in\mathfrak{M}$, by [5, Theorem 15]. Therefore, T has a minimal decomposition set for T by [4, Corollary to Theorem 4].

Conversely, let T be an ideal of A such that $T^*=0$ and $I(\mathfrak{D}_T)=0$. By Theorem 2, $I(\mathfrak{D})\frown T=I(\mathfrak{D}_T)=0$ and hence $I(\mathfrak{D})=0$ because $T^*=0$. By Theorem 1 this completes our proof.

Definition 3. *Let A be a ring. We shall denote by $\mathfrak{D}_0$ the intersection of all dense subsets of $\mathfrak{Q}$ and call it the minimal set for A.* (Suliński [8]).

Lemma 5. *Let A be a ring. Then $p\in\mathfrak{D}_0$ if and only if $\{p\}$ is open in $\mathfrak{Q}$.*

Proof. If we assume that $\{p\}$ is not open, then $\overline{\mathfrak{Q}-\{p\}}=\mathfrak{Q}$, and hence $\mathfrak{Q}-\{p\}\supseteqq\mathfrak{D}_0$. Thus $p\notin\mathfrak{D}_0$.

Conversely, assume that $\{p\}$ is open in $\mathfrak{Q}$ and $p\notin\mathfrak{D}_0$. Then there exists a dense subset $\mathfrak{N}$ of $\mathfrak{Q}$ such that $\mathfrak{N}\notin p$. Accordingly $\mathfrak{N}\subseteqq\mathfrak{Q}-\{p\}$ and $\mathfrak{Q}=\overline{\mathfrak{N}}\subseteqq\overline{\mathfrak{Q}-\{p\}}=\mathfrak{Q}-\{p\}$. This contradiction shows $p\in\mathfrak{D}_0$.

In general, the minimal set $\mathfrak{D}_0$ is contained in the decomposition set $\mathfrak{D}$ by Lemmas 3 and 5. However, in case the structure space of A is a T_1-space, $\mathfrak{D}$ coincides with $\mathfrak{D}_0$.

The following is an extension of [8, Theorem 7].

Theorem 3. *Let A be a semi-prime ring. Then the following conditions are equivalent:*

(1) *$\mathfrak{D}$ is empty.*

(2) *A has no non-zero ideal which is prime as a ring.*

Proof. Assume that $\mathfrak{D}$ is empty and there exists a non-zero ideal T of A which is prime as a ring. Then, by Lemma 1 (2), there is a prime ideal p of A such that $T\frown p=0$. Hence $p^*\supseteqq T\neq 0$, a contradiction.

The converse is easy from Corollary 2.

§ 3. Finally, we shall consider the case where $I(\mathfrak{D})\neq 0$ and $\mathfrak{D}\neq\phi$, that is, the case where A is neither *special* nor *completely non-special* in

Suliński's sense [8].

Lemma 6. *Let A be a semi-prime ring and let T be a non-zero ideal of A such that $T^* \neq 0$.*

(1) *If the ring T has a minimal decomposition set, then the semi-prime ring A/T^*[5] has a minimal decomposition set too.*

(2) *If both T and T^* have minimal decomposition sets, then the ring A has a minimal decomposition set too.*

Proof. Let $\mathfrak{N}$ and $\mathfrak{N}'$ be the sets of all prime ideals $p \in \Omega$ such that $p \supseteq T$ and $p \supseteq T^*$ respectively. Since $T \neq 0$ and $T^* \neq 0$, both $\mathfrak{N}$ and $\mathfrak{N}'$ are not empty, and $\mathfrak{N} \smile \mathfrak{N}' = \Omega$ and $(\mathfrak{D} \frown \mathfrak{N}) \frown (\mathfrak{D} \frown \mathfrak{N}') = \phi$.

Let $\tilde{p}$ be a prime ideal in the ring A/T^*. Then there exists a prime ideal $p \in \mathfrak{N}'$ such that $p/T^* = \tilde{p}$, and $\tilde{p}^*$[6] $= 0$ if and only if $(T^* : p) = T^*$.

(1) Suppose that T has a minimal decomposition set. Then, by Theorems 1 and 2, $0 = I(\mathfrak{D}_T) = I(\mathfrak{D}) \frown T = I(\mathfrak{D} - (\mathfrak{D} \frown \mathfrak{N})) \frown T$ as was seen in the proof of Theorem 2 and these are equal to $I(\mathfrak{D} \frown \mathfrak{N}') \frown T$. Hence $I(\mathfrak{D} \frown \mathfrak{N}')$ ($\subseteq T^*$ whence) $= T^*$. Now, let p be in $\mathfrak{D} \frown \mathfrak{N}'$. Then $\tilde{p}^* \neq 0$. For otherwise, we would have $p^* \subseteq (T^* : p) = T^* \subseteq p$. Thus $\tilde{p}$ is contained in the decomposition set of the ring A/T^*. Therefore $I(\mathfrak{D}_{A/T^*}) \subseteq I(\mathfrak{D} \frown \mathfrak{N}')/T^*$ and hence we have $I(\mathfrak{D}_{A/T^*}) = 0$.

(2) Suppose that both T and T^* have minimal decomposition sets. Then, $0 = I(\mathfrak{D}_{T^*}) = I(\mathfrak{D}) \frown T^* = I(\mathfrak{D}) \frown I(\mathfrak{D} \frown \mathfrak{N}') = I(\mathfrak{D})$ since $T^* = I(\mathfrak{D} \frown \mathfrak{N}')$ as was seen above. Thus, A has a minimal decomposition set.

Combining Lemma 6 (1) with Theorem 2, we obtain a generalization of [1, Theorem 6].

Lemma 7. *Let A be a semi-prime ring and let T be a non-zero ideal of A such that $T^* \neq 0$.*

(1) *If the decomposition set of the ring T is empty, then that of the semi-prime ring A/T^* is also empty.*

(2) *If the decomposition sets of both T and T^* are empty, then that of the ring A is also empty.*

Proof. (1) Suppose that $\mathfrak{D}_T$ is empty. Then by Theorem 2 $I(\mathfrak{D}) \supseteq I(\mathfrak{D}) \frown T = I(\mathfrak{D}_T) = T$ and hence $I(\mathfrak{D})^* \subseteq T^*$. Let $\mathfrak{N}'$ be as in the proof of Lemma 6. Then $\mathfrak{D} \frown \mathfrak{N}' = \phi$. For otherwise, there would exist a prime ideal p such that $p \in \mathfrak{D}$ and $p \supseteq T^*$. Then $p \supseteq T^* \supseteq I(\mathfrak{D})^*$, and hence p^*

5) As is remarked in § 1, $T^* = I(\mathfrak{N})$, $\mathfrak{N} = \{p \in \Omega : p \supseteq T^*\}$, and hence the ring A/T^* is semi-prime.

6) Since no confusion can arise, we shall use this notation in the residue class ring A/T^*.

$\subseteqq I(\mathfrak{D})\subseteqq p$, because $I(\mathfrak{D})^{**}=I(\mathfrak{D})$, which is a contradiction. Let p be in $\mathfrak{N}'$. Then $p\cdot(T^*: p)\subseteqq T^*$, $p\cdot(T^*: p)\cdot T=0$, $(T^*: p)\cdot T\subseteqq p^*=0$ and hence $(T^*: p)$ ($\subseteqq$whence)$=T^*$. This completes our proof.

(2) Suppose that both $\mathfrak{D}_T$ and $\mathfrak{D}_{T^*}$ are empty. Then we have, by Theorem 2, $T=I(\mathfrak{D}_T)=I(\mathfrak{D})\frown T\subseteqq I(\mathfrak{D})$ and $T^*=I(\mathfrak{D}_{T^*})=I(\mathfrak{D})\frown T^*\subseteqq I(\mathfrak{D})$. Therefore $I(\mathfrak{D})\supseteqq T^*\supseteqq I(\mathfrak{D})^*$, $(I(\mathfrak{D})^*)^2=0$, and hence $I(\mathfrak{D})^*=0$. Thus $I(\mathfrak{D})=I(\mathfrak{D})^{**}=A$. This completes our proof.

As an easy consequence of Lemmas 6 and 7, we have the following

Theorem 4. *Let A be a semi-prime ring and let $I(\mathfrak{D})\neq 0$ and $\neq A$.*

(1) *The ring $I(\mathfrak{D})^*$ has a minimal decomposition set.*

(2) *The decomposition set of the ring $I(\mathfrak{D})$ is empty.*

(3)[7] *The semi-prime ring $A/I(\mathfrak{D})$ has a minimal decomposition set.*

(4)[7] *The decomposition set of the semi-prime ring $A/I(\mathfrak{D})^*$ is empty.*

Proof. (1), (2) and (3), (4) follow from Theorem 2 and Lemmas 6 (1) and 7 (1) respectively.

References

[1] A.W. GOLDIE: Decompositions of Semi-Simple Rings, J. London Math. Soc., Vol. 31 (1956), 40-48.

[2] N. JACOBSON: Structure of Rings, Providence (1956).

[3] C.W. KOHLS: The Space of Prime Ideals of a Ring, Fundamenta Math., Vol. 45 (1957), 17-27.

[4] Y. KURATA: Decompositions of Semi-Prime Rings and Jordan Isomorphisms, Osaka Math. J., Vol. 9 (1957), 189-193.

[5] N.H. MCCOY: Subdirect Sums of Rings, Bull. Amer. Math. Soc., Vol. 53 (1947), 856-877.

[6] N.H. MCCOY: Prime Ideals in General Rings, Amer. J. Math., Vol. 71 (1949), 823-833.

[7] M. NAGATA: On the Theory of Radicals in a Ring, J. Math. Soc. Japan, Vol. 3 (1951), 330-334.

[8] A. SULIŃSKI: On Subdirect Sums of Simple Rings with Unity I, Bull. Acad. Polonaise Sci., Ser. Sci., Math., Astr. et Phys., Vol. 8 (1960), 223-228.

[9] A. SULIŃSKI: On the Classification of Semi-Simple Rings, Bull. Acad. Polonaise Sci., Ser. Sci., Math., Astr. et Phys., Vol. 9 (1961), 1-6.

Department of Mathematics,
Hokkaido University

(Received April 30, 1962)

7) Cf. [9].