HOKKAIDO UNIVERSITY

Title	Synthesis of 1-Alkenylboronic Esters via Palladium-Catalyzed Cross-Coupling Reaction of Bis(pinacolato)diboron with 1-Alkenyl Halides and Triflates
Author(s)	Takahashi, Kou; Takagi, Jun; Ishiyama, Tatsuo et al.
Issue Date	2000-02-05
Doc URL	https://hdl.handle.net/2115/56184
Type	journal article
File Information	(20) B-B + X-Vinyl (Com).pdf

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

Synthesis of 1-Alkenylboronic Esters *via* Palladium-Catalyzed Cross-Coupling Reaction of Bis(pinacolato)diboron with 1-Alkenyl Halides and Triflates


Kou Takahashi, Jun Takagi, Tatsuo Ishiyama and Norio Miyaura*

Division of Molecular Chemistry, Graduate School of Engineering, Hokkaido University, Sapporo 060-8628
(Received XXXX)

The synthesis of 1-alkenylboronic acid pinacol esters *via* the palladium-catalyzed cross-coupling reaction of 1-alkenyl halides or triflates with bis(pinacolato)diboron [$(Me_4C_2O_2)B\text{-}B(O_2C_2Me_4)$] was carried out in toluene at 50 °C in the presence of potassium phenoxide (1.5 equivalents) and $PdCl_2(PPh_3)_2 \bullet 2PPh_3$ (3 mol%).


The transition metal-catalyzed cross-coupling reaction of disilanes and distannanes is an elegant method for the synthses of organosilicone[1] and -tin[2] compounds directly from organic electrophiles, but the lack of suitable boron nucleophiles has limited this protocol. We have recently demonstrated the synthesis of arylboronates from aryl halides[3] or triflates[4], and allylboronates from allyl acetates[5] *via* a palladium-catalyzed cross-coupling reaction of tetra(alkoxo)diborons. Very recently, Matsuda and Murata[6] found that pinacolborane $(Me_4C_2O_2)BH$ is an excellent boron-nucleophile for the palladium-catalyzed coupling reaction in the presence of triethylamine. Thus, tetra(alkoxo)diboron and pinacolborane are two nucleophiles now available for the boron cross-coupling reaction.

In this paper, we report the palladium-catalyzed coupling reaction of bis(pinacolato)diboron (**1**) and 1-alkenyl halides or triflates, which provides a one-step procedure for the synthesis of 1-alkenylboronic esters from vinyl electrophiles (Eq. 1)[7].



(1)

Our initial studies (Eq. 2 and Table 1) were focused on the reaction conditions selectively yielding a vinyl boronate (**3**) because the previous reaction reported for the coupling with aryl halides[3] resulted in the formation of an inseparable mixture of several by-products (**4**-**6**) (entry 1). Thus, the reaction involved the Heck coupling[8] between **2a** and **3** yielding a small amount of **5**, the homocoupling between **2a** and **3** giving dimer **6**[9], and an unusual coupling giving a positional isomer **4** together with the desired coupling to afford **3**. Although the formation of positional isomer **4** is not well understood, the Heck product and dimer (**5** and **6**) can be eliminated by changing KOAc or K_2CO_3 (entries 1 and 2) to a strong base because those side reactions are due to the slow transmetalation of **1** to the vinyl-Pd-Br intermediate. Thus, the selective coupling to give **3** was finally achieved by carrying out the reaction at 50 °C in the presence of a solid PhOK suspended in toluene (entries 10 and 11), whereas other solid bases such as K_3PO_4•nH_2O and NaOEt were not effective. The palladium-triphenylphosphine catalyst gave adequately better results than the palladium-dppf complex (entries 3-11).




(2)

The synthesis of 1-alkenylboronates from the representative haloalkenes is summarized in Table 2. The

reaction with 2-iododecene **2b** and *t*-butyl derivative **2c** still suffered from the formation of a positional isomer (entries 1 and 2), but other 2-bromoalkenes (**2d**-**2g**) selectively gave alkenylboronates in high yields, which are not available by conventional hydroboration of terminal alkynes[10] (entries 2-6). The use of an insoluble base suspended in toluene can tolerate various functional groups in haloalkenes (entries 4-6). Although (*E*)-1-alkenylboronates have been synthesized by hydroboration of terminal alkynes and (*Z*)-derivatives by a two-step procedure from 1-halo-1-alkynes[11], the coupling reaction of diboron provided an alternative and stereoselective method for synthesizing both isomers with high stereoselectivity over 99% (entries 7 and 8). 2,2-Disubstituted 1-alkenylboronate and cyclic 1-alkenylboronate were obtained from the corresponding bromides without any difficulty because the side-reactions observed in Eq. 2 were very slow for these substrates (entries 9 and 10). However, all attempts at the couplings with 4-iodo-3-penten-2-one and methyl 3-bromo-2-methylpropenoate were unsuccessful.

Preliminary results for the cross-coupling of diboron **1** with 1-alkenyl triflate are shown in Eq. 3. Under similar reaction conditions to those used for halides, various triflates provided the corresponding 1-alkenylboronates, which results will be reported elsewhere.



(3)

References and notes

1 Y. Hatanaka, T. Hiyama, *Tetrahedron Lett*., **28**, 4715 (1987).; H. Matsumoto, K. Shono, Y. Nagai, *J. Organomet. Chem*., **208**, 145 (1981).

2 H. Azizian, C. Eaborn, A. Pidcock, *J. Organomet. Chem*., **215**, 49-58 (1981); D. Azarian, S. S. Dua, C. Eaborn, D. R. M. Walton, *J. Organomet. Chem*., **117**, C55 (1976).

3 T. Ishiyama, M. Murata, N. Miyaura, *J. Org. Chem*., **60**, 7508 (1995). S. R. Piettre, S. Baltzer, *Tetrahedron Lett*. **38**, 1197 (1997). A. Giroux, Y. Han, P. Prasit, *Tetrahedron Lett*. **38**, 3841 (1997).

4 T. Ishiyama, Y. Itoh, T. Kitano, N. Miyaura, *Tetrahedron Lett*. **38**, 3447 (1997).

5 T. Ishiyama, T.-a. Ahiko, N. Miyaura, *Tetrahedron Lett*. **37**,





6889 (1996). T.-a. Ahiko, T. Ishiyama, N. Miyaura, *Chem. Lett.*, 811 (1997).
6 M. Murata, S. Watanabe, Y. Matsuda, *J. Org. Chem.*, **62**, 6458 (1997);
7 During the course of our study, a similar cross-coupling reaction of haloalkenes with diboron **1** or pinacolborane was reported: S. M. Marcuccio, M. Rodopoulos, H. Weigold, *10th International Conference on Boron Chemistry*, Durham, England, July, 1999, PB-35; M. Murata, T. Oyama, S. Watanabe, Y. Matsuda, *76th Annual Meeting of Chemical Society of Japan*, March, 1999, 2B712.
8 R. F. Heck, in "*Palldium Reagents in Organic Syntheses*", Academic Press, New York (1985), pp 214-242.
9 Dimers included 2-octyl-1,3-dodecadiene and 2,3-dioctyl-1,3-butadiene, but the reaction shown in entry 1 selectively gave the former diene *via* the head to tail coupling between **2a** and **3**. N. Miyaura, A. Suzuki, *J. Organomet. Chem.*, **213**, C53 (1981).
10 I. Rivera and J. A. Soderquist, *Tetrahedron Lett.*, **32**, 2311 (1991). Y. Yamamoto, R. Fujikawa, A. Yamada, N. Miyaura, *Chem. Lett.*, 1069 (1999).
11 D. S. Matteson, in "*Stereodirected Synthesis with Organo-boranes*", Springer, New York (1995), pp 31-41.

Table 1. Reaction conditions[a]

Entry	Catalyst/Base/Solvent	Yield/% (**3**/**4**/**5**/**6**))[b]	
1	$PdCl_2$(dppf)/AcOK/DMSO	53	(51/11/ 1/37)[c]
2	$PdCl_2$(dppf)/K_2CO_3/DMSO	56	(52/ 2/ 5/41)[d]
3	$PdCl_2$(dppf)/PhOK/DMSO	67	(70/ 2/ 4/24)[d]
4	$PdCl_2$(dppf)/PhOK/DMF	77	(71/ 1/ 4/24)[d]
5	$PdCl_2$(dppf)/PhOK/EtOH	69	(80/ 1/ 9/10)[d]
6	$PdCl_2$(dppf)/PhOK/dioxane	86	(91/ 1/ 4/ 4)[d]
7	$PdCl_2$(dppf)/PhOK/toluene	74	(89/ 7/ 3/ 1)[d]
8	$PdCl_2(PPh_3)_2$/PhOK/DMSO	81	(90/ 1/ 1/ 8)[d]
9	$PdCl_2(PPh_3)_2$/PhOK/DMF	89	(92/ 1/ 1/ 6)[e]
10	$PdCl_2(PPh_3)_2$/PhOK/toluene	78	(97/ 1/ 1/ 1)[e]
11	$PdCl_2(PPh_3)_2$•$2PPh_3$/PhOK/toluene	96	(96/ 1/ 2/ 1)[e]

[a]A mixture of 2-bromodecene **2a** (1.0 mmol), diboron **1** (1.1 mmol), palladium catalyst (0.03 mmol) and base (1.5 mmol) in solvent (6 ml) was stirred at 50 °C or 80 °C.
[b]GC yields and their compositions were shown in the parentheses.
[c]At 80 °C for 16 h.
[d]At 50 °C for 16 h.
[e]At 50 °C for 5 h.

Table 2. Synthesis of 1-alkenylboronates (Eq. 1)[a]

entry	haloalkene	yield/%[b]
		
1	**2b**: R=C_8H_{17}, X=I	65[d]
2	**2c**: R=t-C_4H_9, X=Br	69 [e]
3	**2d**: R=Ph, X=Br	88
4	**2e**: R=$(CH_2)_3Cl$, X=Br	85
5	**2f**: R=$(CH_2)_3CN$, X=Br	85
6	**2g**: R=$CH_2CH(OSi^tBuMe_2)CH(CH_3)_2$, X=Br	70
7		47[f]
8		74
9		57
10		99

[a]A mixture of haloalkene (1.0 mmol), **1** (1.1 mmol), $PdCl_2(PPh_3)_2$ (0.03 mmol) plus PPh_3 (0.06 mmol), and PhOK (1.5 mmol) in toluene (5 ml) was stirred at 50 °C for 5 h.
[b]GC yields. The products were isolated by Kugelrohr distillation or chromatography over silica gel.
[d]The reaction accompanied with (*Z*)-1-dodecenylboronate (3%).
[e](*E*)-1-(2-*t*-butylethenyl)boronate (7%) was also produced.
[f](1-octylethenyl)boronate (6%).

Synthesis of 1-Alkenylboronic Esters *via*Palladium-Catalyzed Cross-Coupling Reaction of Bis-(pinacolato)diboron with 1-Alkenyl Halides and Triflates

Koh Takahashi, Atsushi Takagi, Tatsuo Ishiyama, and Norio Miyaura*

