



# HOKKAIDO UNIVERSITY

|                  |                                                                                                                                      |
|------------------|--------------------------------------------------------------------------------------------------------------------------------------|
| Title            | Increased Levels of Plasma p3-Alc alpha 35, a Major Fragment of Alcadein alpha by gamma-Secretase Cleavage, in Alzheimer's Disease   |
| Author(s)        | Omori, Chiori; Kaneko, Madoka; Nakajima, Etsuko et al.                                                                               |
| Citation         | Journal of alzheimers disease, 39(4), 861-870<br><a href="https://doi.org/10.3233/JAD-131610">https://doi.org/10.3233/JAD-131610</a> |
| Issue Date       | 2014                                                                                                                                 |
| Doc URL          | <a href="https://hdl.handle.net/2115/56638">https://hdl.handle.net/2115/56638</a>                                                    |
| Type             | journal article                                                                                                                      |
| File Information | WoS_64980_Hata.pdf                                                                                                                   |



## **Increased levels of plasma p3-Alc $\alpha$ 35, a major fragment of Alcadein $\alpha$ by $\gamma$ -secretase cleavage, in Alzheimer's disease**

**Chiori Omori<sup>1</sup>, Madoka Kaneko<sup>1</sup>, Etsuko Nakajima<sup>2</sup>, Hiroyasu Akatsu<sup>3</sup>, Masaaki Waragai<sup>4</sup>, Masahiro Maeda<sup>5</sup>, Maho Morishima-Kawashima<sup>6</sup>, Yuhki Saito<sup>1</sup>, Tadashi Nakaya<sup>1</sup>, Hidenori Taru<sup>1</sup>, Tohru Yamamoto<sup>7</sup>, Takashi Asada<sup>2</sup>, Saori Hata<sup>1\*</sup> and Toshiharu Suzuki<sup>1\*</sup>, for the Japanese Alzheimer's Disease Neuroimaging Initiative**

1. Laboratory of Neuroscience, Graduate School of Pharmaceutical Sciences, Hokkaido University, Sapporo 060-0812, Japan
2. Department of Neuropsychiatry, Institute of Clinical Medicine, University of Tsukuba, Tsukuba 305-8575, Japan.
3. Choju Medical Institute, Fukushima Hospital, Toyohashi 441-8124, Japan.
4. Department of Neurology, Higashi Matsudo Municipal Hospital, Matsudo 270-2222, Japan
5. Immuno-Biological Laboratories Co., Ltd. (IBL), Fujioka 375-0005, Japan.
6. Department of Molecular Neuropathology, Faculty of Pharmaceutical Sciences, Hokkaido University, Sapporo 060-0812, Japan
7. Department of Molecular Neurobiology, Faculty of Medicine, Kagawa University, Miki-cho 761-0793, Japan

\* Correspondence to: Saori Hata and Toshiharu Suzuki. Laboratory of Neuroscience, Graduate School of Pharmaceutical Sciences, Hokkaido University, Kita12-Nishi6, Kita-ku, Sapporo 060-0812, Japan.

Phone: +81-11-706-3250, Fax: +81-11-706-4991,

E-mail: shata@pharm.hokudai.ac.jp, tsuzuki@pharm.hokudai.ac.jp

Running title: Plasma p3-Alc $\alpha$ 35 levels in AD

Key words: Alzheimer's disease, Alcadein, p3-Alc,  $\gamma$ -secretase, diagnosis, biomarker

## **Abstract**

The p3-Alc $\alpha$  is a metabolic fragment of Alcadin $\alpha$  (Alc $\alpha$ ). Like Alzheimer's amyloid  $\beta$ -protein precursor (A $\beta$ PP) to generate p3 fragment, Alc $\alpha$  is cleaved by  $\alpha$ - and  $\gamma$ -secretases to secrete p3-Alc $\alpha$  peptides in the cerebrospinal fluid (CSF). The p3-Alc $\alpha$  are also detected in the plasma as is amyloid- $\beta$  (A $\beta$ ) which is a metabolic fragment of A $\beta$ PP cleaved by an amyloidogenic  $\beta$ - and  $\gamma$ -secretases. Because p3-Alc $\alpha$  is a non-aggregatable and stable peptide, unlike aggregatable A $\beta$  and metabolically labile p3 of A $\beta$ PP, the changes of p3-Alc $\alpha$  in quality and/or quantity in CSF and plasma are expected as a marker for assessing alteration of substrate cleavage by  $\gamma$ -secretase such as A $\beta$  generation from A $\beta$ PP. The present study describes a sandwich enzyme-linked immunosorbent assay (sELISA) for quantifying levels of p3-Alc $\alpha$ 35, the major form of the p3-Alc $\alpha$  species, and examines levels of p3-Alc $\alpha$ 35 in the plasma of three independent Japanese cohorts. In two of the three cohorts, the p3-Alc $\alpha$ 35 levels were significantly increased with a concomitant decrease in the mini mental state examination (MMSE) score, or in clinically diagnosed AD patients, when compared with age-matched non-demented subjects. The values were significantly lower in AD subjects who were administered donepezil, when compared to AD subjects without donepezil treatment. The increase in plasma p3-Alc $\alpha$ 35 levels may indicate an endophenotype in subjects in whom AD is due to a progressing cognitive impairment in subjects with a  $\gamma$ -secretase malfunction, or a disorder of the clearance of peptides.

**Keywords:** Alcadin, p3-Alc,  $\gamma$ -secretase, Alzheimer's disease, plasma biomarker, donepezil.

## Introduction

Alcadeins (Alc $\alpha$ , Alc $\beta$  and Alc $\gamma$ , also called calsynenin or XB31) constitute a family of neural type I transmembrane proteins, all of which are encoded by their respective genes, and are highly conserved among mammals [1, 2]. Both Alc $\alpha$  and the amyloid- $\beta$  protein precursor (A $\beta$ PP), which is involved in Alzheimer's disease (AD), function as cargo receptors for the kinesin-1 motor that transports membrane vesicles anterogradely in neurons [3-6]. Alc is subject to proteolytic processing by a combination of A $\beta$ PP  $\alpha$ - and  $\gamma$ -secretases, but not an amyloidogenic combination of  $\beta$ - and  $\gamma$ -secretases [7]. This processing of Alc secretes a large amino-terminal extracellular domain and small p3-Alc peptide, along with the intracellular release of the cytoplasmic domain fragments, AlcICDs [8]. In neurons, Alc and A $\beta$ PP form a complex mediated by the cytoplasmic interaction of X11-like (X11L), a neural adaptor protein which stabilizes proteolytic cleavage of both proteins, and facilitate intracellular colocalization of both membrane proteins in the neuron [2, 9, 10]. X11L (also called X11 $\beta$ , Mint2 or APBA2) was recently reported as a possible candidate of ApoE4-related late-onset AD effector [11]. The integrated genomic approach of late-onset/sporadic-type AD suggests that the ApoE4 variant is involved in the intracellular trafficking of A $\beta$ PP, in which X11L plays an important role [12-14].

In transporting membrane vesicles in the late secretory pathway, plasma membrane, or in the endocytotic recycling pathway, Alc and A $\beta$ PP are likely to be cleaved by primary secretases. A $\beta$ PP is cleaved by  $\alpha$ -secretase ADAM 10 and ADAM 17, or  $\beta$ -secretase BACE, to leave the membrane associated A $\beta$ PP carboxy-terminal fragment (A $\beta$ PP CTF $\alpha$  or CTF $\beta$ ) [15], while Alc is cleaved by only  $\alpha$ -secretase to leave Alc CTF [7]. All CTFs are further cleaved by  $\gamma$ -secretase to secrete a p3 peptide from A $\beta$ PP CTF $\alpha$ , A $\beta$  peptide from A $\beta$ PP CTF $\beta$ , and p3-Alc from Alc CTF into the extracellular milieu [7, 8]. Therefore, Alc and A $\beta$ PP perform similar functions [5], show large colocalization in the neuron [2], and are subject to almost the same regulation of proteolytic processing [9]. This suggests that some alteration in the processing systems of the substrates in specific regions of the brain, or the malfunction of the clearance system for secreted short peptides, may appear in the qualitative and/or quantitative alteration of metabolic products derived from A $\beta$ PP and Alc almost equivalently [16]. In fact, recent reports describe that  $\gamma$ -secretase dysfunction and/or malfunction of the

A $\beta$  clearance system are observed in AD patients [16-19], suggesting that some AD pathogenesis is due to an altered membrane substrate cleavage, or a clearance failure of the cleaved products.

The p3 of A $\beta$ PP is metabolically labile to detect in the CSF and plasma, and A $\beta$ , especially the more AD-pathogenic A $\beta$ 42, is progressively aggregatable to detect quantitative or qualitative changes in the plasma. A $\beta$  is a causative metabolic peptide of AD, which detects qualitative and quantitative alterations in the CSF and/or plasma, and is very important in diagnosing AD patients [20]. However, the aggregatable form of this peptide, and various aggregated soluble A $\beta$  oligomers, make it difficult to investigate the alteration of A $\beta$  levels in body fluids precisely. Instead of A $\beta$ , non-aggregatable p3-Alc can be available as a surrogate marker for the detection of changes in the quality and quantity of the  $\gamma$ -cleavage of substrates.

In the adult brain, the expression of Alc $\alpha$  and Alc $\beta$  is more prominent than Alc $\gamma$ , and p3-Alc $\alpha$  and p3-Alc $\beta$  are also more prominent in the CSF than p3-Alc $\gamma$  [7]. In human CSF, p3-Alc $\alpha$ 35 is the major peptide among several p3-Alc $\alpha$  species, while p3-Alc $\beta$ 37 and p3-Alc $\beta$ 40 are the major products of Alc $\beta$  [7, 16]. We previously developed a sandwich ELISA (sELISA) system for quantifying the total amount of p3-Alc $\alpha$  [21]. This sELISA was constructed with pan-p3-Alc $\alpha$  antibodies that can quantify the total amount of all of the p3-Alc $\alpha$  species, including the major p3-Alc $\alpha$ 35, and the minor p3-Alc $\alpha$  species in the CSF and plasma. However, the sELISA cannot selectively quantify specific species such as p3-Alc $\alpha$ 35 or p3-Alc $\alpha$ 38. Despite the restricted conditions of the sELISA, several trials using patients' samples have provided significant information about p3-Alc $\alpha$ : (a) the total p3-Alc $\alpha$  level in the plasma correlates with the level in the CSF of individuals, (b) the total p3-Alc $\alpha$  level correlates with the levels of A $\beta$ 40, an A $\beta$  species that is less aggregatable than A $\beta$ 42, in the CSF and plasma samples of several cohorts, and (c) the CSF and total plasma p3-Alc $\alpha$  levels of several cohorts increases in AD patients, when compared to age-matched control subjects [21-23]. These observations suggest that changes in p3-Alc $\alpha$  levels in body fluids may be able to diagnose AD status. The present study develops a novel monoclonal antibody which specifically recognizes p3-Alc $\alpha$ 35, and establishes a new sELISA system for quantifying p3-Alc $\alpha$ 35 levels. The plasma p3-Alc $\alpha$ 35 levels of AD patients, mild cognitive impaired (MCI) and non-demented subjects of three independent Japanese cohorts will be investigated, in addition to the fluctuation of

p3-Alc $\alpha$ 35 levels in AD subjects treated with donepezil.

## **Methods**

### ***Antibodies and the ELISA system***

p3-Alc $\alpha$ 35 is a peptide that includes the sequence from Ala817 to Thr851 of the human Alc $\alpha$ 1. The monoclonal mouse antibody was raised against an antigen peptide containing the sequence between positions Asn839 and Thr851. The antibody reacts with the antigen peptide specifically, but not with another peptide containing the sequence between positions Asn839 and Ile854 (for amino acid sequence of p3-Alc $\alpha$ , see [7]). Clone 63A1 was selected among several clones showing specific reactivity to p3-Alc $\alpha$ 35. The affinity-purified antibody 63A1 was used to capture p3-Alc $\alpha$ 35. The horseradish peroxidase-conjugated pan-p3-Alc $\alpha$  rabbit polyclonal antibody 817, which was raised against a peptide containing the sequence between positions Ala817 and Val822 [21], and tetramethyl benzidine, were used to detect the captured p3-Alc $\alpha$ 35.

Total amount of p3-Alc $\alpha$  was quantified with an ELISA kit of pan-p3-Alc $\alpha$  monoclonal antibody to capture all p3-Alc $\alpha$  species, which was supplied from Immuno-Biological Laboratories Co., Ltd. This ELISA kit is different from our previous assay system in which pan-p3-Alc $\alpha$  polyclonal antibody 839 is used to capture all p3-Alc $\alpha$  species [21].

Blood samples were collected from the subjects into tubes containing EDTA and centrifuged. Two hundred microliters of plasma was used per duplicate assay. p3-Alc $\alpha$  was extracted from the plasma, as described for the total extraction of p3-Alc $\alpha$  for ELISA [21], and quantified for p3-Alc $\alpha$ 35 using the new sELISA system, in duplicate.

### ***Cohort information***

First cohort (Cohort 1, n=135) is largely composed of MCI and AD patients, many of whom are not hospitalized and living in the countryside together with non-demented control subjects, the second cohort (Cohort 2, n=252) is largely composed of inpatients with normal controls, MCI, AD and other neurological diseases (OND), and the third cohort (Cohort 3, n=91) is a mixture of inpatients and non-hospitalized subjects living in the city. Detailed descriptions of all subjects are shown in supplementary Tables S1 to S3. The cohorts are different from them of previous studies for an analysis of total p3-Alc $\alpha$  [21, 23].

## Results

### *Characterization of the ELISA system with the monoclonal p3-Alc $\alpha$ 35 C-terminal end-specific antibody*

To develop the new sELISA, the 63A1 antibody was used to capture p3-Alc $\alpha$ 35 specifically, instead of the polyclonal pan-p3-Alc $\alpha$  antibody 839 used in previous studies, to examine the total amount of p3-Alc $\alpha$  [21]. The new ELISA recognized p3-Alc $\alpha$ 35 specifically, but not p3-Alc $\alpha$ 39 (**Fig. 1A**), and does not react with the p3-Alc $\beta$  species (**data not shown**), indicating the establishment of a new sELISA system specific for p3-Alc $\alpha$ 35 analysis. Previously, plasma samples were extracted, and total p3-Alc $\alpha$  was quantified using the sELISA, because the treatment of plasma samples with a standard organic extraction protocol removed factor(s) that interfered with the immuno-detection of p3-Alc $\alpha$  [21]. This extraction procedure was also included in the new ELISA to allow significant recovery of p3-Alc $\alpha$  from the plasma, and therefore accurate quantification of p3-Alc $\alpha$ 35 levels in the plasma. The ELISA system was performed to quantify the amount of synthetic p3-Alc $\alpha$ 35 dissolved in the human plasma (**Fig. 1B**). After the extraction process, the slope of the standard curve was almost identical to that of the standard curve in which an identical amount of the synthetic p3-Alc $\alpha$ 35 was dissolved in assay buffer. This procedure results in a yield of over 90% (compare closed square with open circle in Fig. 1B), which is sufficient to be considered quantitative, while the recovery of synthetic p3-Alc $\alpha$ 35 from the plasma was approximately 50% to 60% when plasma samples were assayed without the extraction process (compare closed square with closed triangle in Fig. 1B). Therefore, plasma p3-Alc $\alpha$ 35 quantification was carried out by sELISA after extraction of the endogenous p3-Alc $\alpha$  peptides, as described previously [21].

### **Plasma p3-Alc $\alpha$ 35 levels of subjects in three independent Japanese cohorts**

The levels of p3-Alc $\alpha$ 35 in the plasma of subjects from three cohorts were investigated (**Table 1**) .

The correlation between p3-Alc $\alpha$ 35 and total amount of p3-Alc $\alpha$  levels was first examined. The p3-Alc $\alpha$ 35 levels were significantly correlated with the total amount of p3-Alc $\alpha$  in three cohorts (**Fig. 2 left**). Furthermore, in all cohorts, plasma p3-Alc $\alpha$ 35 levels significantly increased in an age-dependent manner (**Fig. 2 right**).

The relationship between p3-Alc $\alpha$ 35 levels and MMSE scores was then investigated (**Fig. 3**). The increase in plasma p3-Alc $\alpha$ 35 levels correlates with the decrease of the score significantly in two of the three cohorts (**Fig. 3A and B**). These analyses suggest that p3-Alc $\alpha$ 35 levels increase during the aging process, and that subjects appearing to have cognitive impairment show higher levels of p3-Alc $\alpha$ 35 in their plasma.

The p3-Alc $\alpha$ 35 levels of AD and MCI patients were then compared with those of non-demented control subjects or patients of OND patients, in an age-matched population (**Fig. 4 and Table 2**). In all cohorts, AD and MCI were clinically diagnosed based on two major criteria: the Diagnostic and Statistical Manual of Mental Disorders: 4th Edition (DSM-IV) and the National Institute of Neurological and Communicational Disorders and Stroke-Alzheimer's Disease and Related Disorders Association (NINCDS-ADRDA), although we realize that there are many different definitions of MCI [24].

In the subjects of cohort 1 (**Fig. 4A and Table 2A**), the p3-Alc $\alpha$ 35 levels of AD patients were significantly higher than the values of the age-matched non-demented controls. The age-matched MCI subjects showed higher levels of p3-Alc $\alpha$ 35 than non-demented controls, although this finding was not statistically significant.

In cohort 2 (**Fig. 4B and Table 2B**), the subjects of OND was distinguished into two subgroups with (MMSE score  $\leq 22$ ) or without (MMSE score  $\geq 23$ ) remarkable cognitive impairment [25]. We examined the p3-Alc $\alpha$ 35 levels of age-matched control, MCI, AD, OND (MMSE score  $\geq 23$ ) and OND (MMSE score  $\leq 22$ ) subjects. The p3-Alc $\alpha$ 35 level in AD subjects was significantly high when compared to the values of the control subjects. OND patients who show cognitive impairment (MMSE score  $\leq 22$ ) also presented higher p3-Alc $\alpha$ 35 levels than subjects of OND without severe cognitive impairment (MMSE score  $\geq 23$ ) and control subjects, although these were not significant. The present findings suggest that the increase in the p3-Alc $\alpha$ 35 levels may be involved in neurodegeneration and cognitive impairment.

We also analyzed the p3-Alc $\alpha$ 35 levels of 67 age-matched subjects between 63 and 83 years old in cohort 3 (**Fig. 4C and Table 2C**). The levels of p3-Alc $\alpha$ 35 are statistically identical between non-demented controls, MCI, AD and OND subjects.

The subjects of cohorts 2 and 3 were also classified by other criteria for dementia,

the clinical dementia rating (CDR) scale [26]. The p3-Alc $\alpha$ 35 levels were compared in age-matched populations with a CDR score of CDR 0, CDR 0.5 and CDR 1, 2, 3 (**Fig. 5**). CDR 0 subjects were selected from clinically diagnosed MCI and OND patients by criteria of non-demented subjects, and CDR 1, 2, 3 subjects were selected from OND and AD subjects. CDR score of some subjects does not agree with the clinical diagnosis. Thus, we removed subjects, from analysis, who have remarkably different diagnosis such as a subject of MCI with CDR 2. In the present analysis of cohort 2 (**Fig. 5A**), CDR 1, 2, 3 patients (n=197, 198.0  $\pm$  60.5 pg/mL) presented significantly high p3-Alc $\alpha$ 35 levels when compared to those of CDR 0 subjects (n=17, 152.8  $\pm$  44.1 pg/mL) and CDR 0.5 subjects (n=6, 153.1  $\pm$  15.7 pg/mL). In cohort 3 (**Fig. 5B**), there are no significant differences between the respective CDR subjects. Based on at least two criteria (**Figs. 4 and 5**), the p3-Alc $\alpha$ 35 levels showed a tendency of increase in AD and/or demented (CDR 1, 2, 3) subjects.

We also investigated the p3-Alc $\alpha$ 35 levels of AD patients with or without donepezil hydrochloride (Aricept) administration (**Fig. 6**). In 31 age-matched AD subjects with CDR 1 and CDR 2 in cohort 2, non-treated subjects (n=18, average age 80.4  $\pm$  9.2) showed significantly higher levels of plasma p3-Alc $\alpha$ 35 (182.7  $\pm$  35.9 pg/mL) when compared to the levels detected in subjects who were treated with the drug (157.1  $\pm$  24.7 pg/mL; n=13, average age 81.5  $\pm$  5.3), suggesting that the increase in p3-Alc $\alpha$ 35 levels may be slowed by the suppression of cognitive impairment by donepezil administration.

## Discussion

We previously showed that p3-Alc $\alpha$ 35 is the major p3-Alc $\alpha$  species in human CSF by MALDI-TOF/MS spectrometric analysis of p3-Alc $\alpha$  peptides immuno-isolated with a pan-p3-Alc $\alpha$  antibody. By semi-quantitative estimation with the mass spectrometric analysis, approximately 70% or more of the p3-Alc $\alpha$  species is p3-Alc $\alpha$ 35 [7]. Furthermore, using an ELISA system with the pan-p3-Alc $\alpha$  antibody, which can detect all of the p3-Alc $\alpha$  species and quantify total amounts of p3-Alc $\alpha$ , we estimated a total of 5,000–15,000 pg/mL of p3-Alc $\alpha$  in the CSF of human subjects. The levels are comparable to the A $\beta$ 40 levels, and increase in AD patients with a significant correlation to A $\beta$ 40 levels in the CSF [22]. p3-Alc $\alpha$  is detectable in the plasma, but the total p3-Alc $\alpha$  content in the plasma is less than that detected in the CSF, and

approximately 50–300 pg/mL of total p3-Alc $\alpha$  is quantified by the pan-p3-Alc $\alpha$  sELISA system [21]. Although the total plasma p3-Alc $\alpha$  levels also increase in AD patients, it remains unclear whether the p3-Alc $\alpha$ 35 levels increase in AD patients.

In the present study, we analyzed levels of plasma p3-Alc $\alpha$ 35, a major species of the p3-Alc $\alpha$  peptides generated from the cleavage of Alc $\alpha$  by  $\gamma$ -secretase, by using a newly established ELISA system with a C-terminal end-specific monoclonal antibody. First, we found that p3-Alc $\alpha$ 35 is a major p3-Alc $\alpha$  species in the plasma, and we can estimate approximately 60% or more of total p3-Alc $\alpha$  species is p3-Alc $\alpha$ 35 in the plasma, which is a comparable ratio to that of the CSF. We also found that the p3-Alc $\alpha$ 35 levels in the plasma increased in subjects with a lower MMSE score in two of three of the cohorts, although the plasma levels appeared to increase in an age-dependent manner. In age-matched subjects, the p3-Alc $\alpha$ 35 levels increased in AD patients of two of the cohorts. One cohort showed a significant increase in p3-Alc $\alpha$ 35 levels in a CDR-dependent manner among age-matched subjects. Taken together, the plasma p3-Alc $\alpha$ 35 levels showed an increase in subjects who appear to have cognitive impairment, as demonstrated by the total p3-Alc $\alpha$  levels [21, 23]. However, the increased magnitude of plasma p3-Alc $\alpha$ 35 in AD was small compared to the total amount of p3-Alc $\alpha$  detected in our previous studies using the sELISA system with the polyclonal pan-p3-Alc $\alpha$  antibody [21, 23]. Therefore, we cannot rule out the possibility that the p3-Alc $\alpha$ 35 specific monoclonal antibody may lack some component(s) of p3-Alc $\alpha$  that are greatly increased in the blood of AD patients.

Unfortunately, we could not detect a significant increase of p3-Alc $\alpha$ 35 levels of AD patients in cohort 3. This may be due to a cause of heterogeneity of sporadic AD patients. We previously suggested that causes of sporadic AD may be various [22], and a recent publication for the different A $\beta$  fibrils formation in individual AD patients supports this idea [27]. Alternatively, in blood examination, the quantification of p3-Alc $\alpha$ 35 alone may be difficult to classify AD patients clearly. Combination assay with another blood marker may be more effective for blood sample.

The age-matched AD subject population of CDR 1 and CDR 2 treated with donepezil showed lower plasma p3-Alc $\alpha$ 35 values than the population without treatment. Although it remains unclear how donepezil lowers the level of p3-Alc $\alpha$ 35 and this is a result of limited number of subjects, the improvement of neuronal activity may contribute to the suppression of the increase in p3-Alc $\alpha$ 35 levels, or facilitating the

removal of p3-Alc $\alpha$ 35. More convincing study, such as a study to compare the levels of p3-Alc $\alpha$ 35 in patients with AD before and after donepezil treatment, will be needed to confirm this effect. The plasma p3-Alc $\alpha$ 35 levels can be reflected by the levels detected in the CSF of patients with cognitive impairment. If so, the level of plasma p3-Alc $\alpha$ 35 may be an indicator of cognitive ability of aged subjects. Another peptide derived from  $\gamma$ -secretase cleavage of the A $\beta$ PP-like protein 1 (APLP1) is also reported to reflect the amyloidogenic state of A $\beta$ PP metabolism in the brain [17]. The concentration of this peptide in the CSF (4.5 nM) is comparable to the level of p3-Alc $\alpha$  in the CSF [22]; however, the concentration of the APLP1 peptide in the plasma has not been determined. A $\beta$  and p3-Alc $\alpha$  are the main  $\gamma$ -secretase peptide products of the type I membrane proteins expressed largely in the brain that are detectable in the plasma. A $\beta$ PP695 shows neuron-specific expression, while other isoforms such as A $\beta$ PP770/751 are moderately expressed in non-neuronal tissues, including immunocompetent cells in the blood [10, 28, 29]. p3-Alc $\alpha$  is largely derived from the brain because Alc $\alpha$  expression is mostly observed in the brain [2], and increased plasma p3-Alc $\alpha$  levels are detected in Alc $\alpha$ -CTF transgenic mice with a neuron-specific promoter (unpublished observation). Moreover, it should be noted that A $\beta$ PP and Alc express and colocalize largely in neurons, appear to have the same function as kinesin-1 cargo receptors, and are subject to similar proteolytic metabolism [2, 5, 7, 9]. Therefore, changes in the quality and quantity of the plasma p3-Alc $\alpha$  levels may provide a glimpse into the metabolic state of  $\gamma$ -secretase substrates, such as A $\beta$ PP, in a specific brain region. Although physiological functions of p3-Alc $\alpha$  in brain and blood remain unclear, it is interesting to know the function for understanding the means of changes of p3-Alc $\alpha$  levels in aging and loss of cognitive functions. Analysis of more cohorts should be performed to evaluate the changes of plasma p3-Alc $\alpha$ 35 in subjects as an indicator of brain cognitive impairment such as that involved in AD.

## Acknowledgements

This work was supported in part by Grant-in-Aid for Scientific Research (SH and TS) from Ministry of Education, Culture, Sports, Science and Technology (MEXT), and a grant from the New Energy and Industrial Technology Development Organization (NEDO) in Japan. SH was supported by Astellas Foundation for Research on Metabolic

Disorders and The Cell Science Research Foundation. We thank Drs. Yasuo Ihara and Ryozi Kuwano in J-ADNI for their critical discussion.

## References

- [1] Hintsch G, Zurlinden A, Meskenaite V, Steuble M, Fink-Widmer K, Kinter J, Sonderegger P (2002) The calsyntenins a family of postsynaptic membrane proteins with distinct neuronal expression patterns. *Mol Cell Neurosci* **21**, 393-409.
- [2] Araki Y, Tomita S, Yamaguchi H, Miyagi N, Sumioka A, Kirino Y, Suzuki T (2003) Novel cadherin-related membrane proteins, Alcadeins, enhance the X11-like protein mediated stabilization of amyloid  $\beta$ -protein precursor metabolism. *J Biol Chem* **278**, 49448-49458.
- [3] Kamal A, Stokin GB, Yang Z, Xia C-H, Goldstein LB (2000) Axonal transport of amyloid precursor protein is mediated by direct binding to the kinesin light chain subunit of kinesin-1. *Neuron* **28**, 449-459.
- [4] Konecna A, Frischknecht R, Kinter J, Ludwig A, Steuble M, Meskenaite V, Indermuhle M, Engel M, Cen C, Mateos JM, Streit P, Sonderegger P (2006) Calsynenin-1 docks vesicular cargo to kinesin-1. *Mol Biol Cell* **17**, 3651-3663.
- [5] Araki Y, Kawano T, Taru H, Saito Y, Wada S, Miyamoto K, Kobayashi H, Ishikawa OH, Ohsugi Y, Yamamoto T, Matsuno K, Kinjo M, Suzuki T (2007) The novel cargo receptor Alcadein induces vesicle association of kinesin-1 motor components and activates axonal transport. *EMBO J* **26**, 1475-1486.
- [6] Kawano T, Araseki M, Araki Y, Kinjo M, Yamamoto T, Suzuki T (2012) A small peptide sequence is sufficient for initiating kinesin-1 activation through part of TPR region of KLC1. *Traffic* **13**, 834-848.
- [7] Hata S, Fujishige S, Araki Y, Kato N, Araseki M, Nishimura M, Hartmann D, Saftig P, Fahrenholz F, Taniguchi M, Urakami K, Akatsu H, Martins RN, Yamamoto K, Maeda M, Yamamoto T, Nakaya T, Gandy S, Suzuki T (2009) Alcadein cleavages by APP  $\alpha$ - and  $\gamma$ -secretases generate small peptides p3-Alcs indicating Alzheimer disease-related  $\gamma$ -secretase dysfunction. *J Biol Chem* **284**, 36024-36033.

- [8] Piao Y, Kimura A, Urano S, Saito Y, Taru H, Yamamoto T, Hata S, Suzuki T (2013) Mechanism of intramembrane cleavage of Alcadeins by  $\gamma$ -secretase. *PLoS One* **8**, e62431.
- [9] Araki Y, Miyagi N, Kato N, Yoshida T, Wada S, Nishimura M, Komano H, Yamamoto T, De Strooper B, Yamamoto K, Suzuki T (2004) Coordinated metabolism of Alcadein and amyloid  $\beta$ -protein precursor regulates FE65-dependent gene transactivation. *J Biol Chem* **279**, 24343-24354.
- [10] Suzuki T, Nakaya T (2008) Regulation of amyloid  $\beta$ -protein precursor by phosphorylation and protein interactions. *J Biol Chem* **283**, 29633-29637.
- [11] Rhinn H, Fujita R, Qiang L, Cheng R, Lee J H, Abeliovich A (2013) Integrative genomics identifies APOE $\epsilon$ 4 effectors in Alzheimer's disease. *Nature* **500**, 45-50.
- [12] He X, Cooley K, Chung CH, Dashti N, Tang J (2007) Apolipoprotein receptor 2 and X11 $\alpha/\beta$  mediate apolipoprotein E-induced endocytosis of amyloid- $\beta$  precursor protein and  $\beta$ -secretase, leading to amyloid- $\beta$  production. *J Neurosci* **27**, 4052-4060.
- [13] Saito Y, Sano Y, Vassar R, Gandy S, Nakaya T, Yamamoto T, Suzuki T (2008) X11 proteins regulate the translocation of amyloid  $\beta$ -protein precursor (APP) into detergent-resistant membrane and suppress the amyloidogenic cleavage of APP by  $\beta$ -site-cleaving enzyme in brain. *J Biol Chem* **283**, 35763-35771.
- [14] Saito Y, Akiyama M, Araki Y, Sumioka A, Shiono M, Taru H, Nakaya Y, Yamamoto T, Suzuki T (2011) Intracellular trafficking of the amyloid  $\beta$ -protein precursor (APP) regulated by novel function of X11-like. *PLoS One* **6**, e22108.
- [15] Thinakaran G, Koo EH (2008) Amyloid precursor protein trafficking, processing, and function. *J Biol Chem* **283**, 29615-29619.

- [16] Hata S, Fujishige S, Araki Y, Taniguchi M, Urakami K, Peskind E, Akatsu H, Araseki M, Yamamoto K, Martins N R, Maeda M, Nishimura M, Levey A, Chung KA, Montine T, Leverenz J, Fagan A, Goate A, Bateman R, Holtzman DM, Yamamoto T, Nakaya T, Gandy S, Suzuki T (2011) Alternative  $\gamma$ -secretase processing of  $\gamma$ -secretase substrates in common forms of mild cognitive impairment and Alzheimer disease: Evidence for  $\gamma$ -secretase dysfunction. *Ann Neurol* **69**, 1026-1031.
- [17] Yanagida K, Okochi M, Tagami S, Nakayama T, Kodama TS, Nishitomi K, Jiang J, Mori K, Tatsumi S, Arai T, Ikeuchi T, Kasuga K, Tokuda T, Kondo M, Ikeda M, Deguchi K, Kazui H, Tanaka T, Morihara T, Hashimoto R, Kudo T, Steiner H, Haass C, Tsuchiya K, Akiyama H, Kuwano R, Takeda M (2009) The 28-amino acid form of an APLP1-derived Abeta-like peptide is a surrogate marker for Abeta42 production in the central nervous system. *EMBO Mol Med* **1**, 223-235.
- [18] Mawuenyega KG, Sigurdson W, Ovod V, Munsell L, Kasten T, Morris JC, Yarasheski KE, Bateman RJ (2010) Decreased clearance of CNS  $\beta$ -amyloid in Alzheimer's disease. *Science* **330**, 1774.
- [19] Kakuda N, Shoji M, Arai H, Furukawa K, Ikeuchi T, Akazawa K, Takami M, Hatsuta H, Murayama S, Hashimoto Y, Miyajima M, Arai H, Nagashima Y, Yamaguchi H, Kuwano R, Nagaike K, Ihara Y, Japanese Alzheimer's Disease Neuroimaging Initiative (2012) Altered  $\gamma$ -secretase activity in mild cognitive impairment and Alzheimer's disease. *EMBO Mol Med* **4**, 344-352.
- [20] Benilova I, Karran E, De Strooper B (2012) The toxic A $\beta$  oligomer and Alzheimer's disease: an emperor in need of clothes. *Nat Neurosci* **15**, 349-357.
- [21] Konno T, Hata S, Hamada Y, Horikoshi Y, Nakaya T, Saito Y, Yamamoto T, Yamamoto T, Maeda M, Gandy S, Akatsu H, Suzuki T, Japanese Alzheimer's Disease Neuroimaging Initiative (2011) Coordinate increase of  $\gamma$ -secretase reaction products in the plasma of some female Japanese sporadic Alzheimer's disease patients: quantitative analysis with a new ELISA system. *Mol Neurodegen* **6**, 76.

- [22] Hata S, Taniguchi M, Piao Y, Ikeuchi T, Fagan AM, Holzman DM, Bateman R, Sohrabi HR, Martins RN, Gandy S, Urakami K, Suzuki T, Japanese Alzheimer's Disease Neuroimaging Initiative (2012) Multiple  $\gamma$ -secretase product peptides are coordinately increased in concentration in the CSF of a subpopulation of sporadic Alzheimer's disease subjects. *Mol Neurodegen* **7**, 16.
- [23] Kamogawa K, Kohara K, Tabara Y, Takita R, Miki T, Konno T, Hata S, Suzuki T (2012) Utility of plasma levels of soluble p3-Alcadein $\alpha$  as a biomarker for sporadic Alzheimer's disease. *J Alzheimer Dis* **31**, 421-428.
- [24] Stephan BC, Brayne C, McKeith IG, Bond J, Matthews FE. (2008) Mild cognitive impairment in the older population: Who is missed and does it matter? *Int J Geriatr Psychiatry* **23**, 863-871.
- [25] MRC CFAS (1998) Cognitive function and dementia in six areas of England and Wales: the distribution of MMSE and prevalence of GMS organicity level in the MRC CFA Study. *Psychol Med* **28**, 319-335.
- [26] Morris JC (1993) The clinical dementia rating (CDR): current version and scoring rules. *Neurology* **43**, 2412-2414.
- [27] Lu J-X, Qiang W, Yau W-M, Schwieters CD, Meredith SC, Tycko R (2013) Molecular structure of  $\beta$ -amyloid fibrils in Alzheimer's disease brain tissue. *Cell* **154**, 1257-1268.
- [28] Rohan de Silva HA, Jen A, Wickenden C, Jen LS, Wilkinson SL, Patel AJ (1997) Cell-specific expression of  $\beta$ -amyloidprecursor protein isoform mRNAs and proteins in neurons and astrocytes. *Mol Brain Res* **47**, 147-156.
- [29] Monning U, Konig G, Banati RB, Mechler H, Czech C, Gehrman J, Schreiter-Gasser U, Masters CL, Beyreuther K (1992) Alzheimer  $\beta$ A4-amyloid protein precursor in immunocompetent cells. *J Biol Chem* **267**, 23950-23956.

**Table 1. Summary of subjects' data analyzed in Figure 2.**

|                       | A. Cohort 1  | B. Cohort 2  | C. Cohort 3  |
|-----------------------|--------------|--------------|--------------|
| N                     | 135          | 252          | 91           |
| Age (years)           | 74.9 ± 6.50  | 81.3 ± 10.1  | 76.6 ± 8.37  |
| p3-Alcα35 (pg/mL)     | 164.6 ± 46.9 | 192.1 ± 0.1  | 140.5 ± 43.8 |
| p3-Alcα total (pg/mL) | 241.6 ± 45.0 | 240.0 ± 74.4 | 172.5 ± 51.5 |

Average age and average values of p3-Alcα35 and p3-Alcα total in three cohorts are summarized. Numbers indicate means ± standard deviation. Details of individual subjects are shown in Supplemental Tables S1 – S3.

**Table 2. Summary of subjects' data analyzed in Figure 4.**

A.Cohort 1

|                   | Control      | MCI          | AD           |
|-------------------|--------------|--------------|--------------|
| N                 | 15           | 70           | 17           |
| p3-Alcα35 (pg/mL) | 140.1 ± 37.8 | 166.3 ± 45.1 | 178.7 ± 38.5 |
| Age (years)       | 74.3 ± 4.1   | 75.5 ± 3.7   | 77.1 ± 4.1   |

B.Cohort 2

|                   | Control      | MCI          | AD           | OND<br>(MMSE <sub>≥</sub> 23) | OND<br>(MMSE <sub>≤</sub> 22) |
|-------------------|--------------|--------------|--------------|-------------------------------|-------------------------------|
| N                 | 10           | 12           | 116          | 7                             | 85                            |
| p3-Alcα35 (pg/mL) | 148.0 ± 40.3 | 166.4 ± 32.6 | 196.5 ± 55.7 | 159.2 ± 31.7                  | 194.8 ± 64.0                  |
| Age (years)       | 74.6 ± 10.4  | 81.5 ± 4.4   | 82.8 ± 7.7   | 79.3 ± 11.1                   | 78.9 ± 10.1                   |

C.Cohort 3

|                   | Control      | MCI          | AD           | OND          |
|-------------------|--------------|--------------|--------------|--------------|
| N                 | 19           | 22           | 17           | 9            |
| p3-Alcα35 (pg/mL) | 128.6 ± 26.8 | 146.0 ± 49.5 | 130.8 ± 32.7 | 122.9 ± 52.3 |
| Age (years)       | 74.5 ± 4.4   | 76.2 ± 6.4   | 75.3 ± 6.6   | 74.4 ± 5.7   |

Average age and average values of p3-Alcα35 in three cohorts are summarized. Numbers indicate means ± standard deviation. MCI, mild cognitive impairment. AD, Alzheimer's disease. OND, other neurological diseases.

## Figure legends

### Figure 1. Specificity of the sELISA.

**A. Specific reactivity of the sELISA system to p3-Alcα35.** The indicated amount of synthetic p3-Alcα35 (closed square) and p3-Alcα39 (open diamond) were dissolved in buffer A (PBS containing 1% (w/v) bovine serum albumin and 0.05% (v/v) Tween-20), and assayed with the ELISA.

**B. Quantification of the p3-Alcα35 peptide in human plasma with or without an extraction process.** The indicated amounts of synthetic p3-Alcα35 were dissolved in human plasma (prepared from a non-AD healthy volunteer; open circle and closed triangle) or buffer A (closed square). The plasma was subject to extraction with (open circle) or without (closed triangle) a standard organic extraction protocol prior to analysis by ELISA, as described [21]. The horseradish peroxidase-conjugated antibody and tetramethyl benzidine were used to detect the captured p3-Alcα35 colorimetrically at OD<sub>450</sub>.

### Figure 2. Correlation of p3-Alcα35 levels with total p3-Alcα levels in plasma and age-dependency of plasma p3-Alcα35 levels.

**(Left)** The correlation between p3-Alcα35 and total p3-Alcα levels are shown. **A.** Cohort 1 ( $r^2=0.1009$ ,  $p<0.001$ ), **B.** Cohort 2 ( $r^2=0.5362$ ,  $p<0.001$ ), **C.** Cohort 3 ( $r^2=0.5790$ ,  $p<0.001$ ). **(Right)** The correlation between p3-Alcα35 levels and age are shown. **A.** Cohort 1 ( $r^2=0.1512$ ,  $p<0.001$ ), **B.** Cohort 2 ( $r^2=0.055912$ ,  $p<0.001$ ), **C.** Cohort 3 ( $r^2=0.2005$ ,  $p<0.001$ ). Statistical analysis was performed by using the Pearson's correlation coefficient test. Subject numbers (N), average age and p3-Alcα35 amounts of each cohort are summarized in Table 1.

### Figure 3. The relationship between MMSE scores and plasma p3-Alcα35 levels.

The correlation between mini mental state examination (MMSE) scores and p3-Alcα35 levels in the plasma of subjects is shown. In the graphs, statistical analysis was performed by using the Pearson's correlation coefficient test. **A.** Cohort 1 ( $r^2=0.05303$ ,  $p<0.01$ ), **B.** Cohort 2 ( $r^2=0.03525$ ,  $p<0.01$ ), **C.** Cohort 3 ( $r^2=0.02934$ ,  $p=0.1045$ ).

**Figure 4. Levels of plasma p3-Alcα35 following subgrouping into non-demented, MCI, AD and OND.**

Subjects from three cohorts were clinically divided into four groups: non-demented subject (control), subjects with mild cognitive impairment (MCI), Alzheimer's disease (AD), and patients with other neurological diseases (OND). In cohort 2 (**B**), OND are further distinguished into subjects with (MMSE score  $\leq 22$ ) or without (MMSE score  $\geq 23$ ) remarkable cognitive impairment. Plasma p3-Alcα35 levels of age-matched subjects in these subgroups were compared within the respective cohorts. **A.** Cohort 1 (subjects between 70 and 83 years old), **B.** Cohort 2 (subjects between 73 and 94 years old), **C.** Cohort 3 (subjects between 63 and 83 years old). Statistical analysis was performed using the Dunn's multiple comparison test following the Kruskal-Wallis test. \*,  $P < 0.05$ . Subject numbers (N), average age and p3-Alcα35 amounts are summarized in Table 2.

**Figure 5. Levels of plasma p3-Alcα35 of subjects divided into subgroups using the clinical dementia rating (CDR) scale.**

Age-matched subjects of cohort 2 (**A**) and cohort 3 (**B**) were divided into three groups based on CDR scales. Statistical analysis was performed using the Dunn's multiple comparison test following the Kruskal-Wallis test. \*,  $P < 0.05$ ; \*\*,  $P < 0.01$ .

**Figure 6. Influence of donepezil hydrochloride administration on plasma p3-Alcα35 levels.**


Age-matched AD (CDR 1 and CDR 2) patients treated with Aricept (donepezil hydrochloride) were selected from cohort 2, and their plasma p3-Alcα35 levels were compared to those of age-matched subjects who were not treated with Aricept (non-treated). Statistical analysis was performed using the Mann-Whitney test. \*,  $P < 0.05$ .

**Supplementary Table S1. Information of subjects in cohort 1.**

**Supplementary Table S2. Information of subjects in cohort 2.**



**Supplementary Table S3. Information of subjects in cohort 3.**

# Fig.1





# Fig.2



**A**




**B**



**C**




# Fig.3




# Fig.4

**A**



**B**



**C**




Fig.5




Fig.6

