



HOKKAIDO UNIVERSITY

Title	Glaciological observations in 2012 and 2013 at SIGMA-A site, Northwest Greenland
Author(s)	YAMAGUCHI, Satoru; MATOBA, Sumito; YAMAZAKI, Tetsuhide et al.
Citation	Bulletin of Glaciological Research, 32, 95-105 https://doi.org/10.5331/bgr.32.95
Issue Date	2014-08
Doc URL	https://hdl.handle.net/2115/56974
Type	journal article
File Information	yamaguchi_bgr_2014_fieldrepo.pdf



Glaciological observations in 2012 and 2013 at SIGMA-A site, Northwest Greenland

Satoru YAMAGUCHI¹, Sumito MATOBA², Tetsuhide YAMAZAKI³, Akane TSUSHIMA⁴,
Masashi NIWANO⁵, Tomonori TANIKAWA⁶ and Teruo AOKI⁵

¹ Snow and Ice Research Center, National Research Institute for Earth Science and Disaster Prevention, Nagaoka, 940-0821, Japan
* yamasan@bosai.go.jp

² Institute of Low Temperature Science, Hokkaido University, Sapporo, 060-0819, Japan

³ Avangnaq, Takatsuki, 569-0094, Japan

⁴ Graduate School of Environmental Science, Hokkaido University, Sapporo, 060-0810, Japan

⁵ Meteorological Research Institute, Tsukuba, 305-0052, Japan

⁶ Japan Aerospace Exploration Agency, Tsukuba, 305-8505, Japan

(Received May 7, 2014; Revised manuscript received July 17, 2014)

Abstract

Glaciological observations were conducted in 2012 and 2013 at the SIGMA-A site on the northwest Greenland ice sheet (78° 03'06"N, 67° 37'42"W, 1490 m a.s.l.) as part of the Snow Impurity and Glacial Microbe effects on abrupt warming in the Arctic (SIGMA) project. The meteorological conditions during the two observations were quite different. The meteorological condition during the 2012 observation period was warm, and heavy rainfall occurred during the observation period, thus the snow was very wet. In contrast, the meteorological condition during the observation period in 2013 was cold, with a blowing snow event, thus the snow was quite dry.

The glaciological observations in 2012 consisted of 1) snow-stake measurements, 2) snow pit observations, 3) grain size observations for validation of satellite-derived snow products, 4) snow specific surface area measurements using a near-infrared camera, 5) snow sampling for chemical analyses, and 6) drilling of firn cores with a hand auger. The glaciological observations in 2013 consisted of 1) snow-stake measurements, 2) snow pit observations, and 3) snow sampling for chemical analyses.

Key words: SIGMA project, SIGMA-A, glaciological observation, snow property, accumulation area in north-west Greenland

1. Introduction

The Greenland ice sheet is presently undergoing drastic changes (Steffensen *et al.*, 2008; van den Broeke *et al.*, 2009; Rignot *et al.*, 2011). One of the possible drastic changes in the accumulation areas in Greenland is “positive albedo feedback (Box *et al.*, 2012),” namely, an increase in snow grain size due to acceleration of snow metamorphism, thus reducing the albedo when the air temperature increases. To elucidate the snow/ice albedo feedback effect caused by snow grain growth and light-absorbing snow/ice impurities, including glacial microbes, the Snow Impurity and Glacial Microbe effects on abrupt warming in the Arctic (SIGMA) project was launched in 2011 (Aoki *et al.*, 2014a). As part of the SIGMA project, an automatic weather station (AWS) was established in the accumulation area on the northwest Greenland ice sheet (78°03'06"N, 67°37'42"W, 1490 m a.s.l.), which was named “SIGMA-A site” (Fig. 1), and field campaigns for glaciological observations at SIGMA-A site

were conducted in 2012 and 2013 (Aoki *et al.*, 2014a).

The 2012 glaciological observation was conducted at SIGMA-A site from June 26 to July 16, 2012. The meteorological conditions during the first half of this observation period (June 26 to July 9, 2012) were relatively warm (average air temperature was -0.2°C), while heavy rainfall (60 to 100 mm) occurred from July 10 to July 13, 2012, in the second half (average air temperature was 1.3°C) (Aoki *et al.*, 2014a). During the same period of observation in 2012, a record melting event of surface snow/ice was observed over the Greenland ice sheet (Nghiem *et al.*, 2012; Tedesco *et al.*, 2013).

The 2013 glaciological observations were conducted at SIGMA-A site from July 23 to July 30, 2013. The monthly average air temperature in July 2013 was approximately 4°C colder than that in July 2012, and heavy blowing snow occurred during the observation period in 2013. There was thus a large difference between the meteorological conditions during the glaciological observations in 2012 and 2013. The aim of this paper is to present the preliminary results of




Fig. 1. Map showing the location of SIGMA-A site (78°03'06"N, 67°37'42"W, 1490 m a.s.l.).




Fig. 2. Location of observation sites in 2012 at SIGMA-A site.

glaciological observations in 2012 and 2013 at SIGMA-A site for further studies.

2. Glaciological observations in 2012

2.1 Snow-stake measurements

A snow-stake network was established on June 30, 2012 (Fig. 2) consisting of 16 snow-stakes from Japan, made of bamboo, approximately 2 m in length. The snow-stakes were inserted into the snow cover until they reached a thick ice layer at approximately 80 cm below the snow surface (the average value was 82.4 cm with a standard deviation of ± 27.1 cm). Each height from the snow surface to the top of the snow-stake (h) was measured each day at 9:00 a.m. from June 30 to July 13, 2012. Then, the surface lowering Δh (cm), which is the difference between the h value of the previous day and the present day, was calculated.

Fig. 3a shows the values of Δh_{ave} , which is the Δh averaged over all the snow-stake data at each measurement. Because the Δh values showed a small fluctuation at each stake, the dispersion of the data is also shown by bars. Data from July 13, which is shown in a different color, had different measurement intervals due to lack of data on July 11 and 12 because of heavy rainfall (Aoki *et al.*, 2014b); namely, the value consists of the cumulative data from three days, July 11 to July 13. The fluctuation




Fig. 3. Results of snow-stake observations in 2012.

a. Results of daily surface lowering and mean air temperature. Blue circles indicate the averaged surface lowering over all measurements from 9:00 a.m. of the previous day to 9:00 a.m. of the present day. Red circle indicates the averaged cumulative surface lowering from 9:00 a.m. on July 10 to 9:00 a.m. on July 13. Bars indicate the dispersion of the data. Black rhombuses indicate the mean air temperature during each observation interval.

b. The relationship between positive degree hours and cumulative surface lowering or cumulative melting. Blue circles indicate the cumulative value of surface lowering since June 30, 2012. Black rhombuses indicate the cumulative value of melting estimated using the measured surface density.

of mean air temperature from 10:00 a.m. of the previous day to 9:00 a.m. of the present day, which was measured at 3 m height at the AWS of SIGMA-A (Aoki *et al.*, 2014a), is also shown in Fig. 3a. Although the mean air temperature were sometimes negative, there was a good correlation between the surface lowering and mean air temperature; large surface lowering occurred when the mean air temperature was high.

The surface lowering should include the following three pieces of information: 1) erosion by wind, 2) melting, and 3) settling of the surface due to compression by snow cover. We considered that the effect of the erosion should be neglected because of the wet snow surface condition. Although it was difficult to estimate surface settling from the observation data, and the numerical snow cover simulation should be examined to evaluate the amount of surface settling, we deemed that the influence of surface settling on Δh was small because no snowfall occurred, and the densities of the snow cover were relatively high during the observation period.

In general, the amount of melt showed good correlation with the cumulative positive daily mean temperature (Positive Degree Day, PDD) (*e.g.*, Ohmura, 2001).

Because the period of our data was limited, we introduced the idea of positive degree hours (ΣT_h^+), which is essentially the same as PDD, namely, the sum of positive hourly air temperatures during the observation period ($^{\circ}\text{C h}$). Fig. 3b shows the relationship between the cumulative Δh_{ave} ($\Sigma \Delta h_{ave}$) (cm) and ΣT_h^+ during the observation period. There is a high linear correlation between $\Sigma \Delta h_{ave}$ and ΣT_h^+ , as shown by the following equation:

$$\Sigma \Delta h_{ave} = 0.16 \Sigma T_h^+ \quad (R=0.99), \quad (1)$$

where R is the correlation coefficient.

The values of the average melting (M_{ave}) at each Δh_{ave} were estimated using the measured surface density data, then the cumulative M_{ave} ($\Sigma \Delta M_{ave}$) (mm for water equivalent, W.E.) during the observation period was also calculated (Fig. 3b). There is also a high linear correlation between $\Sigma \Delta M_{ave}$ and ΣT_h^+ , as shown by the following equation:

$$\Sigma \Delta M_{ave} = 0.59 \Sigma T_h^+ \quad (R=0.99). \quad (2)$$

Here, it must be noted that M_{ave} may not be equal to ablation because there is the possibility of internal accumulation of meltwater, such as superimposed ice formation, which we will discuss in section 2.2, “Snow pit observations.”

2.2 Snow pit observations

Standard snow pit observations were carried out at approximately 10:00 a.m. from June 30 to July 13, 2012. The observation items were snow stratigraphy (grain shape F and grain size E), snow density ρ , snow temperature T , mass liquid water content (LWC) calculated from a dielectric measurement (Denoth *et al.*, 1984) and ρ , and water equivalent of all layers above a thick ice layer, which will be discussed later, using a snow sampler. The snow pit observations followed the methodology outlined by the Nihon Seppyo Gakkai (2010).

A thick ice layer, which was too hard to penetrate, was found 88 cm below the surface during the first snow pit observation on June 30. The layers above the thick ice layer were essentially dry, except for a few layers near the surface which were probably caused by a melting event in very recent days. This result implies that melting events did not occur after accumulation of snow above the thick ice layer, except for recently on June 30. On the other hand, the pit observation data on July 30, 2011 at 1469 m a.s.l., which is almost the same altitude as SIGMA-A, showed the existence of thick melt forms layers (85 cm thick) due to large melting events (Aoki *et al.*, 2014a). In addition to this observation result, the layer just above the thick ice layer consisted of a developed depth hoar, which indicated a record high temperature gradient after accumulation of this layer (Akitaya, 1974). Based on previous studies of development of depth hoars overlying wet snow (*e.g.* Fukuzawa and Akitaya, 1993; Jamieson and Fierz, 2004), we considered that this depth hoar layer developed from a layer overlying the snow surface of the previous summer, which




Fig. 4. Snow pit observations in 2012.

The colors and symbols representing snow classifications are in accordance with the snow classification system of Fierz *et al.* (2009). F is grain shape, E is grain size, LWC is mass liquid water content, T is snow temperature, and ρ is snow density.

must have been wet. For these reasons, we finally concluded that this thick ice layer was the reference surface of the previous summer surface, and all snow pit observations were conducted for the snow cover above this thick ice layer. Our assumption that IF_{ref} was the previous summer surface was supported by the result of chemical analyses of the snow pit described in section 2.5, “Snow sampling for chemical analyses.” An approximately 1.5 m wide pit was dug on June 30, and the pit was expanded forward at least 1 m from the previous pit wall at each observation (Fig. 2).

Fig. 4 shows the measurement results at each snow pit observation, in which snow heights (H) are defined as the height from IF_{ref} . The distributions of snow temperature and grain shape in the first half-period (June 30 to July 5), except for the data from July 2 and 3, indicated that melting did not dominate during this period because of low temperatures (Fig. 3a). The data from July 2 and 3 showed that the snow temperature during the most




Fig. 4. Continued.

snow cover was 0°C, and the dominant grain shapes were melt forms, which should have been affected by the existence of a water pass because the snow stake observations showed that remarkable melting (surface lowering) did not occur during this period (Fig. 3a). After July 6, the snow temperature was 0°C from the top, and the 0°C position reached the bottom on July 10 because the mean air temperature during this period became positive (Fig. 3a).

Fig. 5 shows the ratios of melt forms, including ice layers and dry grain shapes. In this analysis, layers that had two grain shapes, melt forms and dry grain shapes, were classified as melt forms layers. The ratio of melt forms increased with time, and all layers became melt forms after the heavy rainfall event on July 10–13 (Aoki *et al.*, 2014b).

Fig. 6 shows the values of the water equivalent (mm W. E.) of snow cover above the IF_{ref} measured using a snow sampler and estimated using the distribution of snow density measured during each snow pit observation. Essentially, the value measured using the snow sampler was smaller than that estimated from the distribution of

snow density during the snow pit observation. It is not clear why the values from the snow sampler were smaller than those estimated from the distribution of snow density, but one possibility is that the snow sampler measurement did not catch a part of the depth hoar layer on the IF_{ref}, which was very fragile. It is notable that both estimation methods showed a gradual decrease of the water equivalent of snow cover with time. Two possible explanations for the decrease of the water equivalent of snow cover with time should be considered. The first possible explanation is the discharge of meltwater from the area. We considered this possibility to be small because the complete impermeable layers that are needed to make aquifers for large horizontal movement of water, such as thick ice layers, were not observed except for IF_{ref}, although some discontinuous ice layers with melt forms were observed. Moreover, the snow cover was so cold that a large amount of water could not have moved away from the area before it was refrozen. The second possibility is that meltwater reached the IF_{ref} by preferred flow through water passes and refroze there (superimposed ice information). The existence of

Fig. 4. Continued.




Fig. 4. Continued.




Fig. 5. The ratios of melt forms and dry grain types in 2012. Red areas are the ratios of melt forms and blue areas are dry grain shapes.

water passes was sometimes observed, although these areas were usually avoided in order to carry out snow pit observations, and the existence of water passes was not reflected in the data from the daily snow pit observations. One indication of a water pass can be seen in the data from July 2 and 3, in which the ratios of melt forms were




Fig. 6. Measurement results of the water equivalent of snow cover in 2012.

Gray bars indicate measured data using the snow sampler, and black bars indicate the water equivalent of snow cover using the distribution of density of the snow pit observations.




Fig. 7. Measurement results of 4 m wide snow pit on July 4, 2012.

a. Photo of 4 m wide snow pit.

b. Horizontal variations of snow height (red line), depth hoar layer (green area), and ice layers (blue lines).

larger (Fig. 5), and the IF_{ref} temperature was higher than on subsequent days (Fig. 4).

The temperature of IF_{ref} gradually increased with time and reached 0°C. This trend may have resulted from the heat generated by refreezing water on the IF_{ref}. For these reasons, we considered that the cause of the decrease of the water equivalent of snow cover with time is the surface position change of IF_{ref} due to superimposed ice formation on the IF_{ref}, which was not taken into account in the snow sampling and snow pit observations. However, there is no direct evidence of superimposed ice formation, thus our discussion calls for further analyses, such as using a numerical snow cover model.

To examine the horizontal variation of snow structure, a 4 m wide snow pit observation was conducted on July 4, 2012 (Fig. 7). In the snow pit observation, the distributions of snow surface height from the IF_{ref} and the height of the top of the depth hoar layer on the IF_{ref} were measured with a 0.5 m horizontal resolution, and the

Table 1. Horizontal variations of measured snow densities from a 4 m wide snow pit observed on July 4, 2012. Values in parentheses indicate standard deviations.

Position	Near surface of snow cover (Position measured from the surface)			Inner part of snow cover (Position measured from the top of IF _{ref})		
	0–3 cm	2–5 cm	7–10 cm	62 cm	40 cm	17 cm
Density (kg m ⁻³)	362 (±40)	425 (±9)	426 (±14)	410 (±25)	386 (±17)	369 (±21)

horizontal distributions of each ice layer were also measured (Fig. 7). Here, it must be noted that the following discussion is based on the assumption that IF_{ref} has a flat level throughout the 4 m wide snow pit. We regard this assumption to be reasonable, although the exact level of IF_{ref} was not measured. The snow height averaged over a width of 4 m was 84.8 cm, with a standard deviation of ± 1.9 cm, and the height of the top of the depth hoar averaged over 4 m was 11.4 cm, with a standard deviation of ± 1.1 cm. These horizontal variations in the snow pit observations were negligible. Some ice layers persisted throughout the 4 m width, but other ice layers appeared and disappeared in the 4 m interval. Moreover, the positions of the ice layers fluctuated up and down, even when they were persistent. These results imply that the distributions of ice layers were strongly affected by the local degree of water penetration. Thus, interpretation of the distribution of ice layers in the snow pit observation data needs to consider local influences.

Table 1 shows the variations of snow densities at several heights, measured with 1 m horizontal resolution. The snow density at 0–3 cm from the surface showed large variations, with a standard deviation of 11% for the averaged value, while other snow densities showed small variations, with a standard deviation of less than 6% of the averaged values. These results indicate that the surface conditions varied widely.

2.3 Grain size observations for validation of satellite-derived snow products

In addition to traditional grain size (Fierz *et al.*, 2009), detailed grain size observations near the surface were conducted because one of the aims of the glaciological observations at SIGMA-A site was the validation of satellite-derived snow products (Aoki *et al.*, 2007; Hori *et al.*, 2007; Stamnes *et al.*, 2007). To compare the optical properties of snow and measured grain size in the field, Aoki *et al.* (2007) proposed three definitions of grain size in field observations: d_1 , the length of the major axis of crystals or dendrites; d_2 , the branch width of dendrites or the width of a narrower portion of broken crystals; and d_3 , the thickness of plate-like crystals. Aoki *et al.* (2007) also reported that d_2 generally corresponds to the optical grain size (d_{opt}). In the detailed grain size observations, maximum, minimum and medium d_2 values were measured at three positions (0–2 cm, 2–5 cm, and 5–10 cm




Fig. 8. Measurement results of d_2 near the surface in 2012. Black circles indicate the medium values at each measurement position. Bars indicate the range of d_2 from maximum to minimum.
a. Data at 0–2 cm depth from the surface.
b. Data at 2–5 cm depth from the surface.
c. Data at 5–10 cm depth from the surface.

from the surface) using a hand held lens.

Fig. 8 shows the fluctuation of d_2 during the observation period. In the figure, the maximum to minimum ranges of d_2 are shown by a bar. The values of d_2 at all depths fluctuated around 1.0 mm, except for the measurement at 0–2 cm on July 4 and 5. The data on July 4 and 5 showed a smaller value of d_2 than other data because of the developing surface hoar (Hori *et al.*, 2014).

2.4 Snow specific surface area measurement using a near-infrared (NIR) camera

Recently, optical methods based on the near-infrared reflectance value (*Ref*) have been used to measure the specific surface area (SSA) of snow in the field (*e.g.* Matzl and Schneebeli, 2006; Gallet *et al.*, 2009; Arnaud *et al.*, 2011; Zuanon, 2013). During the glaciological observation in 2012, one of the optical methods, namely, the NIR photometry method using around 900 nm wavelength light (Matzl and Schneebeli, 2006), was introduced for measuring the SSA of snow in the field. Detailed descriptions of NIR photometry have been provided by Matzl and Schneebeli (2006) and Yamaguchi *et al.* (2014).




Fig. 9. Continued.

Warren, 1999; Mitchell, 2002; Langlois *et al.*, 2010).

2.5 Snow sampling for chemical analyses

The stable isotope composition of precipitation is controlled by variations in the origin of the water and the condensation history during atmospheric transport, and the chemical composition of precipitation is controlled by its source and the history of scavenging processes and chemical reactions during atmospheric transport. Therefore, vertical profiles of the stable isotope and chemical composition of surface snow on the Greenland ice sheet can provide valuable information relating to the temporal variations of transport processes of water vapor and aerosols over the Greenland ice sheet.

Snow samples for chemical analyses were collected on June 30, 2012, from the snow pit wall using a stainless steel tool to transfer snow into polyethylene bags. The snow samples were melted at ambient temperature in a camping tent and then transferred into 10 mL polypropylene bottles. The liquid samples were then transported to the Institute of the Low Temperature, Hokkaido University, Japan.

The δD values were measured by a cavity ring-down spectroscopy (model L2130-i, Picarro) with a vaporizer (model A0212, Picarro). The typical analytical precision was $\pm 0.5\%$. The concentrations of ion species were determined by ion chromatography (model DX-500x, Dionex). The typical analytical precision was 10%, and the detection limit of each species was 20 ppb.

Fig. 10 shows the vertical profiles of δD and concentrations of Cl^- , NO_3^- , SO_4^{2-} , and Ca^{2+} . The δD values above 66 cm height were high. Generally, the δD value in snow is mainly determined by the air temperature at which vapor condenses (Johnsen *et al.*, 1989). Therefore,

Digital photographs for analyses of SSA were taken every other day using a Nikon D3S, which was improved to detect near-infrared areas with a 25-mm Zeiss ZF-IR lens and gelatin filter (X-Nite 850 nm Filter) that selected wavelengths from 840 to 940 nm.

Fig. 9 shows the distribution of reflectance (Ref : %), calculated SSA (mm^{-1}) using measured Ref , and optical grain size (d_{opt} : mm) estimated from SSA ($d_{opt} = 6/SSA$). Although the data near the surface are plotted in the figure, these data may include errors because of the influence of light from the surface (Kuchiki *et al.*, 2013), thus these data should be treated with caution during analyses. The fluctuations in SSA for each day generally agreed closely with the stratigraphy, namely, large changes in SSA (Ref) appeared at the boundaries of each layer. The values of d_{opt} were usually smaller than grain size (E) from the snow pit observation (Fig. 4), and should correspond to different definitions of grain size, as has been reported by several researchers (*e.g.* Grenfell and




Fig. 10. Results of chemical analyses of snow samples obtained on June 30, 2012.

we assumed that the snow layer above 66 cm height corresponds to spring and early summer. According to the seasonally determined δD , Cl^- , which mainly originates from sea salt, showed a peak in late winter, and NO_3^- , which mainly originates from fossil fuel combustion and biomass burning, and Ca^{2+} , which mainly originates from terrestrial dust, showed peaks in the spring and early summer. These seasonal variations of ion species are roughly consistent with those observed in the inland ice sheet (Summit) (Whitlow *et al.*, 1992). The correspondence of seasonal variations of ion species between the coastal site (SIGMA-A) and inland suggest that aerosols deposited at SIGMA-A are supplied by long range transportation. In the figure, δD and Cl^- seem to increase again near the thick ice layer (IF_{ref}). Moreover, Ca^{2+} shows another peak near the thick ice layer (IF_{ref}). We interpret these trends as indication that the snow cover above the IF_{ref} consists of one year of accumulation since the previous summer (2011).

2.6 Firn cores drilled with a hand auger

To reconstruct the temporal variations of mass balance, 19-m firn cores were drilled on July 3 and 6, 2012, with a hand auger designed by the technical division of the Institute of Low Temperature Science, Hokkaido University. In addition, 5.5-m firn cores were also drilled with a hand auger on July 10, to test the application of the NIR photometry method to core analyses.

The 19-m firn cores had several thick ice layers in their shallower positions from the surface (Fig. 11a), but few ice layers in the deeper positions (Fig. 11b). In fact, the melt feature percentage (MFP) of the 19-m firn-ice core shows that high MFP values appear in the upper layers above a depth of 7 m W. E., while the MFP becomes almost zero in the deeper layers below a depth of 8 m W. E. (Aoki *et al.*, 2014a). This result indicated that more frequent snow melting events have occurred in recent years.

Fig. 11c shows the NIR image of the 5.5-m firn core at 2.5 m depth using the NIR photometry method. In the NIR Image, ice and firn are clearly distinguished, and




Fig. 11. Example photos of firn core obtained in 2012.

- Image of the 19 m firn core at 5 m depth (shallower layer).
- Image of the 19 m firn core at 15 m depth (deeper layer).
- Images of 5.5 m firn core at 2.5 m depth using the NIR photometry method. The diameter of the core is 10 cm.

detailed distributions of ice in the firn are also shown. This result implies that the NIR photometry method is useful for firn core analysis (*e.g.*, quantitative analyses of MFP). Based on this result, the NIR photometry method for firn core analysis has been improved (Aoki *et al.*, 2013).

Using the 19-m bore hole, temperatures at 10 m, 15 m, and 19 m depths from the surface were measured using temperature loggers (Ondotori Jr. TR-51i, T&D corporation), which were installed in the bore-hole and kept at

each depth for one day (Table 2). In general, the temperature at 10m depth was considered to be the annual mean temperature at the observation site, thus the mean annual temperature at SIGMA-A is considered to be approximately -20°C , while the measured annual mean air temperature from July 2012 to June 2013, which was calculated from the data of AWS in SIGMA-A, was -19.3°C .

3. Glaciological observations in 2013

3.1 Snow-stake measurements

A snow-stake network was established on July 24, 2013, in the same manner as the 2012 observations.

Table 2. Bore-hole temperatures measured in 2012.

Depth from the surface (m)	Temperature ($^{\circ}\text{C}$)
10	-19.9
15	-20.0
19	-20.3




Fig. 12. Results of snow-stake observations in 2013.

Blue circles indicate the averaged surface lowering over all measurements from 9:00 a.m. of the previous day to 9:00 a.m. of the present day. Red circle indicates the averaged cumulative surface lowering from 9:00 a.m. on July 24 to 9:00 a.m. on July 26. Bars indicate the dispersion of the data. Black rhombuses indicate the mean air temperature at each observation interval.

Snow-stakes were inserted into the snow cover until they reached a thick ice layer, which was at approximately 120cm below the snow surface (the average value was 117.3cm, with a standard deviation of $\pm 2.5\text{cm}$). Each height from the snow surface to the top of the snow stake (h) was measured each day at 9:00 a.m. from July 24 to July 29, 2013.

Fig. 12 shows the value of the surface change (Δh_{ave}) (cm), which is the surface change (Δh) averaged over all the snow-stake data at each measurement. The dispersion of the data is also shown by bars. Here, the positive Δh values indicate accumulation. Data from July 26, which is shown in a different color, has a different measurement interval due to lack of data from July 25, namely, its value was derived from the cumulative data of 2 days, from July 24 to July 26. The fluctuation of the mean air temperature during the observation intervals, which was measured at the AWS at SIGMA-A (Aoki *et al.*, 2014a), is also shown in the figure. Because the mean air temperature values during the observation period were negative, the Δh_{ave} values were positive, that is, accumulation occurred. The fluctuation of Δh was much larger in 2013 than in 2012, which was probably caused by the redistribution of snow by a blowing snow event (Aoki *et al.*, 2014a).

3.2 Snow pit observation and snow sampling for chemical analyses

The observation period in 2013 was only one week long and the weather conditions were bad (Aoki *et al.*, 2014a), thus snow pit observations and snow sampling for chemical analyses, which were performed in the same way as in 2012, were conducted only once, on July 27, 2013.

Fig. 13 shows the results of the snow pit observation and δD analyses. A pit was dug to a depth of 132cm from the surface, in which a melt-freeze crust layer existed. We considered this layer to be the melted snow surface from the summer of 2012. The dry snow crystals were dominantly grain-shaped, although some thin ice layers were present. This result indicated that




Fig. 13. Measurement results of snow pit observation and chemical analyses of snow obtained on July 23, 2013.

The colors and symbols representing snow classifications are in accordance with the snow classification system of Fierz *et al.* (2009). F is grain shape, T is snow temperature, and ρ is wet density.

significant snow melting had not occurred since the summer of 2012. Therefore, δD values showed a more obvious seasonal cycle in 2013 than in 2012, when the snow layer was affected by melt water (Fig. 10).

4. Summary

Glaciological observations were conducted in 2012 and 2013 at the SIGMA-A site on the northwest Greenland ice sheet (78°03'06"N, 67°37'42"W, 1490 m a.s.l.) as part of the SIGMA project. The meteorological conditions during the two observations were quite different: the meteorological condition in 2012 was warm, and heavy rainfall occurred during the observation period, while the meteorological condition in 2013 was cold, with a blowing snow event. These different meteorological conditions were reflected in the snow conditions: the snow condition in 2012 was very wet because of melting and rainfall, while the snow condition in 2013 was quite dry.

The glaciological observations in 2012 consisted of 1) snow-stake measurements, 2) snow pit observations, including observations of detailed grain size and specific surface area of snow measurement, 3) drilling of firn cores, and 4) snow sampling for chemical analyses. The glaciological observations in 2013 consisted of 1) snow-stake measurements, 2) snow pit observations, and 3) snow sampling for chemical analyses.

The data reported in this paper should be helpful for understanding the glaciological conditions in the accumulation area of the northwest Greenland ice sheet and can be used to improve the snow physical model SMAP (Niwano *et al.*, 2012) and to validate satellite-derived snow products for the SIGMA Project.

Acknowledgements

We thank the field campaign members of the SIGMA Project and GRENE-Arctic Project in Greenland in 2012 and 2013. This work was supported by (1) Japan Society for the Promotion of Science (JSPS), Grant-in-Aid for Scientific Research (S), number 23221004, (2) Green Network of Excellence (GRENE) Arctic Climate Change Research Project funded by MEXT Japan.

References

- Akitaya, E. (1974): Studies on depth hoar. *Contrib. Inst. Low Temp. Sci., Ser. A*, **26**, 1–67. <http://hdl.handle.net/2115/20238>.
- Aoki, T., Hori, M., Motoyoshi, H., Tanikawa, T., Hachikubo, A., Sugiura, K., Yasunari, T. J., Strovol, R., Eide, H. A., Stamnes, K., Li, W., Nieke, J., Nakajima, Y., Takahashi, F. (2007): ADEOS-II/GLI snow/ice products: Part II - Validation results using GLI and MODIS data. *Remote Sens. Environ.*, **111**, 274–290. doi: 10.1016/j.rse.2007.02.035.
- Aoki, T., Niwano, M., Yamaguchi, S., Motoyoshi, H., Kuchiki, K., Matoba, S. and Hachikubo, A. (2013): Optical measurements of snow and ice sample using near-infrared image sensors. *Summaries of JSSI & JSSE Joint Conference on Snow and Ice Research 2013 in Kitami*, 13 (in Japanese).
- Aoki, T., Matoba, S., Uetake, J., Takeuchi, N. and Motoyama, H. (2014a): Field activities of the “Snow Impurity and Glacial Microbe effects on abrupt warming in the Arctic” (SIGMA) Project in Greenland in 2011–2013. *Bull. Glaciol. Res.*, **32**, <http://dx.doi.org/10.5331/bgr.32.3>.
- Aoki, T., Matoba, S., Yamaguchi, S., Tanikawa, T., Niwano, M., Kuchiki, K., Adachi, K., Uetake, J., Motoyama, H. and Hori, M. (2014b): Light-absorbing snow impurity concentrations measured on Northwest Greenland ice sheet in 2011 and 2012. *Bull. Glaciol. Res.*, **32**, <http://dx.doi.org/10.5331/bgr.32.21>.
- Arnaud, L., Picard, G., Champollion, N., Domine, F., Gallet, C. J., Lefebvre, E., Fily, M., Barnola, M. J. (2011): Measurement of vertical profiles of snow specific surface area with a 1cm resolution using infrared reflectance: instrument description and validation. *J. Glaciol.*, **57**(201), 17–29. <http://www.igsoc.org/8080/journal/57/201/j10j080.pdf>.
- Box, J. E., Fettweis, X., Stroeve, J. C., Tedesco, M., Hall, D. K. and Steffen, K. (2012): Greenland ice sheet albedo feedback: thermodynamics and atmospheric drivers. *The Cryosphere*, **6**, 821–839, doi: 10.5194/tc-6-821-2012.
- Denoth, A., Foglar, A., Weiland, P., Mätzler, C., Aebischer, H., Tiuri, M., Shivola, A. (1984): A comparative study of instruments for measuring the liquid water content of snow. *J. Appl. Phys.* **56**(7), 2154–2160. <http://dx.doi.org/10.1063/1.334215>.
- Fierz, C., Armstrong, L. R., Durand, Y., Etchevers, P., Greene, E., McClung, M. D., Nishimura, K., Satyawali, K. P., Sokratov, A. S. (2009): The International classification for seasonal snow on the ground. *Technical documents in Hydrology (UNESCO document)*, **83**, 80 pp. http://www.cryosphericciences.org/products/snowClassification/snowclass_2009-11-23-tagged-highres.pdf.
- Fukuzawa, T. and Akitaya, A. (1993): Depth-hoar crystal growth in the surface layer under high temperature gradient. *Ann. Glaciol.*, **18**, 39–45. http://www.igsoc.org/annals.old/18/igs_annals_vol18_year1993_pg39-45.pdf.
- Gallet, J. C., Domine, F., Zender, C. S., Picard, G. (2009): Measurement of the specific surface area of snow using infrared reflectance in an integrating sphere at 1310 and 1550 nm. *The Cryosphere*, **3**, 167–182. doi: 10.5194/tc-3-167-2009.
- Grenfell, T. and Warren, G. S. (1999): Representation of a nonspherical ice particle by a collection of independent spheres for scattering and absorption of radiation. *J. Geophys. Res.*, **104**(D24), 697–709. doi: 10.1029/1999JD900496.
- Hori, M., Aoki, T., Stamnes, K. and Li, W. (2007): ADEOS-II/GLI snow/ice products - part III: Retrieved results. *Remote Sens. Environ.*, **111**, 274–319. doi: 10.1016/j.rse.2007.01.025.
- Hori, M., Aoki, T., Tanikawa, T., Kuchiki, K., Niwano, M., Yamaguchi, S. and Matoba, S. (2014): Dependence of thermal infrared emissive behaviors of snow cover on the surface snow type. *Bull. Glaciol. Res.*, **32**, <http://dx.doi.org/10.5331/bgr.32.33>.
- Jamieson, B. and Fierz, F. (2004): Heat flow from wet to dry snowpack layers and associated Faceting. *Ann. Glaciol.*, **38**, 187–194. <http://dx.doi.org/10.3189/172756404781814762>.
- Johnsen, S. J., Dansgaard, W. and White, J. W. C. (1989): The origin of Arctic precipitation under present and glacial conditions. *Tellus*, **41B**, 452–468, doi: 10.1111/j.1600-0889.1989.tb00321.x.
- Kuchiki, K., Aoki, T., Niwano, M., Yamaguchi, S., Hachikubo, A., Takeuchi, Y., Hori, M., Tanikawa, T. (2013): Retrieval of snow grain size using optical measurements in near-infrared wavelengths. *Summaries of JSSI & JSSE Joint Conference on Snow and Ice Research 2013 in Kitami*, 14 (in Japanese).
- Langlois, A., Royer, A., Montpetit, B., Picard, G., Brucker, L., Arnaud, L., Harvey-Collard, P., Fily, M., and Goïta, K. (2010): On the relationship between snow grain morphology and in-situ near infrared calibrated reflectance photographs. *Cold Reg. Sci. Tech.*, **61**, 34–42. doi: 10.1016/j.coldregions.2010.01.004.

- Matzl, M. and Schneebeli, M. (2006): Measuring specific surface area of snow by near-infrared photography. *J. Glaciol.*, **52**(179), 558–564. doi: 10.3189/172756506781828412.
- Mitchell, D. L. (2002): Effective Diameter in radiation Transfer: General Definition, Applications, and Limitation. *Journal of the Atmospheric Science*, **59**, 2330–2346. doi: 10.1175/1520-0469(2002)059<2330:EDIRTG>2.0.CO;2.
- Nghiem, S. V., Hall, D. K., Mote, T. L., Tedesco, M., Albert, M. R., Keegan, K., Shuman, C. A., DiGirolamo, N. E. and Neumann, G. (2012): The extreme melt across the Greenland ice sheet in 2012. *Geophys. Res. Lett.*, **39**, L20502, doi: 10.1029/2012GL053611.
- Nihon Seppyo Gakkai (2010): Glaciological observation guidebook. *Asakura Shoten*, 136 pp.
- Niwano, M., Aoki, T., Kuchiki, K., Hosaka, M. and Kodama, Y. (2012): Snow Metamorphism and Albedo Process (SMAP) model for climate studies: Model validation using meteorological and snow impurity data measured at Sapporo, Japan. *J. Geophys. Res.*, **117**, F03008, doi: 10.1029/2011JF002239.
- Ohmura, A. (2001): Physical basis for temperature-based melt-index method. *J. Appl., Meteorol.*, **40**(4), 753–761, [http://dx.doi.org/10.1175/1520-0450\(2001\)040<0753:PBFTTB>2.0.CO;2](http://dx.doi.org/10.1175/1520-0450(2001)040<0753:PBFTTB>2.0.CO;2).
- Rignot, E., Velicogna, I., van den Broeke, M. R., Monaghan, A. and Lenaerts, J. (2011): Acceleration of the contribution of the Greenland and Antarctic ice sheets to sea level rise. *Geophys. Res. Lett.*, **38**, L05503, doi: 10.1029/2011GL046583.
- Stamnes, K., Li, W., Eide, H., Aoki, T., Hori, M. and Storvold, R. (2007): ADEOS-II/GLI snow/ice products - part I: Scientific Basis. *Remote Sens. Environ.*, **111**, 258–273, doi: 10.1016/j.rse.2007.03.023.
- Steffensen J. P., Andersen K. K., Bigler M., Clausen H. B., Dahl-Jensen D., Fischer H., Goto-Azuma K., Hansson M., Johnsen S. J., Jouzel J., Masson-Delmotte V., Popp T., Rasmussen S. O., Röthlisberger R., Ruth U., Stauffer B., Siggaard-Andersen M. L., Sveinbjörnsdóttir A. E., Svensson A. and White J. W. (2008): Abrupt climate change happens in few years high-resolution Greenland ice core data show abrupt climate change happens in few years. *Science*, **321**, 680–684, doi: 10.1126/science.1157707.
- Tedesco, M., Fettweis, X., Mote, T., Wahr, J., Alexander, P., Box, J. E. and Wouters, B. (2013): Evidence and analysis of 2012 Greenland records from spaceborne observations, a regional climate model and reanalysis data. *The Cryosphere*, **7**, 615–630, doi: 10.5194/tc-7-615-2013.
- van den Broeke, M., Bamber, J., Ettema, J., Rignot, E., Schrama, E., van de Berg, W. J., van Meijgaard, E., Velicogna, I. and Wouters, B. (2009): Partitioning recent Greenland mass loss. *Science*, **326**, 984–986, doi: 10.1126/science.1178176.
- Whitlow, S., Mayewski, P. A. and Dibb, J. E. (1992): A comparison of major chemical species seasonal concentration and accumulation at the South Pole and summit, Greenland. *Atmos. Environ.*, **26A**(11), 2045–2054, doi: 10.1016/0960-1686(92)90089-4.
- Yamaguchi, S., Motoyoshi, H., Tanikawa, T., Aoki, T., Niwano, M., Takeuchi, Y. and Endo, Y. (2014): Application of snow specific surface area measurement using an optical method based on near-infrared reflectance around 900-nm wavelength to wet snow zones in Japan. *Bull. Glaciol. Res.*, **32**, <http://dx.doi.org/10.5331/bgr.32.55>.
- Zuanon, N. (2013): IceCube, a portable and reliable instruments for snow specific surface area measurement in the field. *International Snow Science Workshop Grenoble - Chamonix Mont-Blanc- 2013 proceedings*, 1020–1023.