



HOKKAIDO UNIVERSITY

Title	100 years Zoraptera : a phantom in insect evolution and the history of its investigation
Author(s)	Mashimo, Y.; Matsumura, Y.; Machida, R. et al.
Issue Date	2014
Doc URL	https://hdl.handle.net/2115/57321
Type	journal article
File Information	2014zorarev.pdf



100 years Zoraptera – a phantom in insect evolution and the history of its investigation

Mashimo, Y., Y. Matsumura, R. Machida, R. Dallai, M. Gottardo, K. Yoshizawa, F. Friedrich, B. Wipfler & R.G. Beutel

Y. Mashimo, R. Machida, Sugadaira Montane Research Center, University of Tsukuba, Sugadaira Kogen, Ueda, Nagano 386-2204, Japan

R. Dallai, M. Gottardo, Department of Life Sciences, University of Siena, Via A. Moro 2, I-53100 Siena, Italy

Y. Yoshizawa, Systematic Entomology, Hokkaido University, Sapporo, Hokkaido 060-8589, Japan

F. Friedrich, Biozentrum Grindel und Zoologisches Museum Hamburg, , Martin-Luther-King-Platz 3, Universität Hamburg, 20146 Hamburg, Germany

R.G. Beutel, B. Wipfler, Y. Matsumura, Institut für Spezielle Zoologie und Evolutionsbiologie, Erbertstr. 1, FSU Jena, 07743 Jena, Germany (rolf.beutel@uni-jena.de)

Zoraptera are a cryptic and enigmatic group of insects. The species diversity is lower than in almost all other groups of Hexapoda, but may be distinctly higher than presently known. Several new species were described from different regions recently. The systematic placement was discussed controversially since the group was discovered 100 years ago. Affinities with Isoptera and Psocoptera were discussed in earlier studies. A sistergroup relationship with Acercaria (Psocodea, Thysanoptera, Hemiptera) was proposed by W. Hennig, for the first time based on a strictly phylogenetic argumentation. More recent studies consistently suggest a

placement among the “lower neopteran orders” (Polyneoptera). Close affinities to Dictyoptera were proposed and alternatively a sistergroup relationship with Embioptera or with Embioptera + Phasmatodea (Eukinolabia), respectively. The precise placement is still controversial and the intraordinal relationships are largely unclear. Recent transcriptome analyses tentatively suggest a clade Zoraptera + Dermaptera as sistergroup of all other polyneopteran orders. The oldest fossils are from Cretaceous amber. An extinct genus from this era may be the sistergroup of all the remaining zorapterans. The knowledge of the morphology, development and features related to the reproductive system greatly increased in recent years. The general body morphology is very uniform, whereas the genitalia differ strongly between species. This is likely due to different kinds of selection, i.e. sexual selection in the case of the genital organs. The mating pattern also differs profoundly within the order. A unique external sperm transfer occurs in *Zorotypus impolitus*. A species-level phylogeny and more investigations of the reproductive system should have high priority.

Introduction

Zoraptera, also known as groundlice or angel insects (e.g., Grimaldi & Engel 2005), are a cryptic, inconspicuous and still enigmatic group of insects. Weidner (1969) pessimistically pointed out that they are in an evolutionary dead end and can only lead a wretched life (“kümmerliches Dasein”) in a very limited habitat. The systematic placement has been controversial since the group was introduced as an order by Silvestri (1913) (e.g., Trautwein et al. 2012) (see Table 1). Consequently the term “the Zoraptera problem” was coined by Beutel & Weide (2005).

1 Zorapterans were characterized by Silvestri (1913) as „Insecta terrestria, parva, aptera,
2 agila, praedantia“, which means „living in earth, small, apterous, agile and predacious
3 insects” (Weidner 1969). The scientific name given to the order („purely apterous ones”,
4 Greek: zoros = pure, strong; aptera = apterous) is a misnomer as zorapterans are primarily
5 winged (Caudell 1920) and small and poorly sclerotized besides. The wing dimorphism is one
6 of few autapomorphies of the order, correlated with the presence or absence of compound
7 eyes and ocelli, and the presence or absence of a distinct pigmentation, with distinctly darker
8 alate specimens.

9 Zoraptera currently comprise 39 extant species and nine species are known as fossils
10 (Engel 2008; Mashimo et al. 2013). They were considered as the least known insect order by
11 Kristensen (1995). The fact that the situation has changed profoundly in recent years (see Fig.
12 1) stimulated us to present this review. The aims are to summarize the current knowledge, to
13 point out remaining problems, and to illuminate the history of the investigation of this small
14 group of cryptic insects.

16 1. Distribution, diversity and taxonomy

17 Zoraptera are largely restricted to tropical and subtropical regions. They live under bark or
18 inside galleries made in rotting wood by other insects. Only *Zorotypus hubbardi* has expanded
19 its range as far north as Indiana, Iowa and Illinois. This species might survive in colder
20 regions by hiding in sawdust (Riegel 1963). Zoraptera have not been recorded from Australia,
21 but *Zorotypus novobritannicus* was recently described from New Britain (Terry & Whiting
22 2012) and the group was also recorded from New Zealand and Easter Island (Weidner 1969;
23 Choe 1989, 1992; Grimaldi & Engel 2005).

24 Silvestri (1913) introduced the order and described the genus and three species (in
25 Latin). He collected the type species *Zorotypus guineensis* himself in Ghana (“Costa d’Oro”),
26 and specimens of *Zorotypus ceylonicus* (Sri Lanka) and *Zorotypus javanicus* (Java) were

provided by other entomologists. Eight new species from different parts of the world were described in the next 15 years (e.g., Karny 1922, 1927), and four species including the North American *Z. hubbardi* were introduced in a study also containing a key and a discussion of possible relationships of the order (Caudell 1918). A catalog of the Order published by Hubbard (1990) contained 29 extant species and one from the Eocene. Two new species were described by Chao & Chen (2000) and Engel (2000). A distributional checklist of zorapteran species was published in Engel & Grimaldi (2002) and an updated checklist of “World Zoraptera” by Rafael & Engel (2006). In the latter, 34 extant and six fossil species were listed, and the authors provided information on sexes, winged forms, and nymphs of each species. The most recent account is given in Mashimo et al. (2013). This study contains the descriptions of three new species from Peninsular Malaysia, bringing the number of extant zorapterans to 39. Zoraptera have been regarded as rare and one of the least diverse group of hexapods. However, apparently their diversity remains underexplored (Rafael & Engel 2006; Mashimo et al. 2013).

In taxonomic studies the shape of the basal antennomeres, the chaetotaxy of the ventral metafemoral surface, the shape of the cerci, and the male genitalia are traditionally recognized as useful to define species. The male genitalia are highly variable and potentially suitable for classifying the group. It has been noted that closely related species with very similar external features can be clearly discriminated based on male genital structures (Paulian 1949, 1951; Hwang 1974, 1976; New, 1978; Rafael & Engel 2006; Rafael et al. 2008; Mashimo et al. 2013). Some species were described based solely on immature or female specimens (e.g., Silvestri 1913; Caudell 1923, 1927; New 1995), or the information on the male genitalia was insufficient (e.g., Choe 1989; Zompro 2005; Rafael & Engel 2006; Terry & Whiting 2012). In future taxonomic studies, a detailed investigation of both sexes or at least the male genitalia should be obligatory for an unambiguous circumscription of species (Mashimo et al. 2013).

2. Morphology

The illustrations provided by Silvestri (1913) were of high standard, displaying external and internal features, including the setation, the mouthparts, endoskeletal and pretarsal structures, and also internal organs such as the central nervous system, digestive tract, tracheal system and excretory organs (six free Malpighian tubules). A series of comparative studies were carried out by G.C. Crampton. They covered head structures, the neck region, the thorax, the wings, and wing base sclerites of Zoraptera and other groups of insects (Crampton 1918, 1920, 1921, 1926, 1927). Gurney (1938) provided more morphological information, mainly based on the North American species *Z. hubbardi*. Like Silvestri (1913) and Crampton (e.g., 1918, 1927) he used simple dissection techniques and light microscopy. He described and illustrated external body parts but also male and female internal genital organs, the digestive tract, Malpighian tubules and even eggs. Delamare-Deboutteville (1947) compared alate and wingless exemplars. The thoracic skeletomuscular system was described for the first time by Rasnitsyn (1998). The availability of only one damaged wingless exemplar and the application of simple preparation techniques led to incomplete and not fully satisfying results. Weidner (1970) summarized the available information in the "Handbuch der Zoologie" series.

In the 21st century the study of Zoraptera accelerated. Beutel & Weide (2005) provided the first complete study of head structures (*Z. hubbardi*) using semithin microtome sections and SEM. Friedrich & Beutel (2008) described the thoracic anatomy of alate and wingless specimens of the same species and emphasized the highly conserved condition of the skeletomuscular apparatus, presumably close to the neopteran groundplan. Based on the results of this study the authors proposed a new consistent and extendable nomenclature for the thoracic muscle system of neopteran insects (Friedrich & Beutel 2008). Wing base structures of *Z. caudelli* were investigated by Yoshizawa (2007, 2011a). A computer-based 3D reconstruction of the male postabdomen of *Z. hubbardi* was presented by Hünefeld

(2007), based on microtome sections. Structures of the male postabdomen were described by Bolívar y Pieltain (1940), Hwang (1974), New (1978), Rafael & Engel (2006) and others. They are extremely variable, probably related to different mating patterns (e.g., Dallai et al. 2013).

Recently different aspects of Zoraptera were investigated by a collaborative group including entomologists from Japan (University of Tsukuba), Italy (University of Siena), and Germany (Friedrich-Schiller-Universität Jena), mainly specialized on development (R. Machida), ultrastructure (R. Dallai) and the skeletomuscular system (R.G. Beutel), respectively. Combined with collecting efforts in Malaysia (R. Machida and coworkers) and Ecuador (Y. Matsumura) (Fig. 2), several studies focused on the genital region were presented, using a broad array of techniques including transmission electron microscopy (TEM), confocal laser scanning microscopy (CLSM), and micro computed tomography (μ CT) (Fig. 3). One result was the marked discrepancy between a far-reaching uniformity of the group in the general body morphology and conspicuous differences in the male genital structures (Dallai et al. 2011, 2012a, b; Y. Matsumura, pers. comm.).

Zorapteran eggs were described in earlier studies for several species (Caudell 1920; Gurney 1938; Silvestri 1946). SEM micrographs of eggs of *Zorotypus gurneyi* and *Zorotypus barberi* (Panama) were shown in Choe (1989). Mashimo et al. (2011) described eggs of *Z. caudelli* (Malaysia) with different approaches including TEM. The whitish egg is 0.6 mm long and 0.3 mm wide, the two-layered chorion shows a honeycomb pattern, and an operculum or a hatching line are missing. A pair of micropyles is present at the equator on the dorsal side, similar to the condition found in eggs of *Timema* (Phasmatodea) (Mashimo et al. 2011).

Spermatozoa were already observed by Silvestri (1913). The spermatogenesis and sperm ultrastructure was investigated by Dallai et al. (2011) using transmission electron microscopy. A common feature of Zoraptera is the great length of the spermatozoa. An

unusual feature of *Z. caudelli* is a disconnection of sub-tubules A and B at the level of microtubule doublets 1 and 6 of the mature sperm cells (Dallai et al. 2011), whereas disproportionately large mitochondrial derivatives are characteristic for *Z. hubbardi* (Dallai et al. 2012b). Character combinations found in different species suggest that spermatozoa do not evolve as a unit, but that like in other body regions components can be modified independently from each other. A derived feature, dense laminae radiating in a cartwheel array between neighboring centriolar triplets, is shared with Phasmatodea and Embioptera. An apomorphy shared with Phasmatodea is the presence of 17 protofilaments in the tubular wall of the outer accessory microtubules (Dallai et al. 2011, 2012b; Gottardo et al. 2012).

3. The phylogenetic position of Zoraptera (Figs 4, 5)

3.1. Pre-Hennigian approaches

When Silvestri (1913) described the first zorapteran species and introduced the family Zorotypidae he assumed that they must be close relatives (“collocare vicino”) of roaches (“Blattoidei”) and Isoptera. He listed several differences separating Zoraptera from these polyneopteran groups, such as for instance the presence of a bundle of setae on the left mandible. He also mentioned similarities with Dermaptera, but explicitly referred to them as superficial. After Silvestri, affinities (not necessarily in a phylogenetic sense) with Isoptera were emphasized by Caudell (1918) and the “distinguished albeit eccentric” G.C. Crampton (1920) (Grimaldi & Engel 2005). Potential arguments for a closer relationship included a general resemblance in the habitus, certain structural details of the thorax and abdomen, colonial habits, and dehiscent wings. While recognizing these similarities, Crampton (1920) clearly pointed out affinities with the acercarian orders, for which he hypothesized an origin from “Prothorthopteroid ancestors in the common Prothorthopteran-Protoblattid stem”. He explicitly suggested a very close relationship with Psocoptera (“Corrodentia”). A possible

relationship with Psocoptera was also discussed in Gurney (1938), concluding that “affinities with Corrodentia are more noticeable than those with orthopteroid insects”.

Weidner (1969) suggested “strong thoracic synapomorphies” of Zoraptera + Isoptera, but did not uphold this view in his *Handbuch der Zoologie* volume (Weidner 1970) where he explicitly rejected superficial arguments for such a hypothesis and emphasized the difficulty of placing Zoraptera. He rather vaguely referred to the order as a specialized, “today obsolete (“verkümmert”) branch of Blattodea” (including roaches and termites).

3.2. Hennig’s interpretation and follow up studies

W. Hennig revolutionized insect systematics with his classical work “Die Stammesgeschichte der Insekten” (Hennig 1969). In an earlier study he rather vaguely suggested monophyletic Paraneoptera¹ (Zoraptera + Acercaria [= Psocodea, Thysanoptera and Hemiptera]) with Zoraptera basal and definitely outside of Psocoptera (Hennig 1953). A similar hypothesis was proposed in a little-known study by Wille (1960: Fig. 1), who addressed Zoraptera as “the most primitive group of the hemipteroids [= Acercaria], at the base of their evolutionary branch and close to the orthopteroids”. Hennig (1969) indicated a single “certain derived feature” supporting Paraneoptera, the greatly condensed condition of the abdominal ganglionic chain, with two separate ganglionic masses in zorapterans, and only one in the remaining groups (Hennig 1953, 1969). He considered the reduced number of three tarsomeres (groundplan) as an additional potential synapomorphy, but it is apparent that losses of tarsal segments occurred in many groups. The hypotheses suggested by Hennig (1953, 1969) and Wille (1960), were tentatively followed by Kristensen (e.g., 1975), Willmann (2005), and Beutel & Weide (2005). The presence of only six Malpighian tubules (four in Acercaria), a cercus composed of only one segment or entirely missing (Acercaria), a

¹ The term Paraneoptera is presently often used for a group that does not include Zoraptera (e.g., Grimaldi & Engel 2005)

fork of the cubitus posterior (areola postica) (Willmann 2005), a strengthened cibarial sucking pump, and a blade-like lacinia lacking mesally directed spines (Beutel & Weide 2005) were considered as additional arguments. The last feature is likely an autapomorphy of *Zorotypus hubbardi*, the species examined by Beutel & Weide (2005). A largely unmodified lacinia is present in other species (e.g., Silvestri 1913). The strongly developed cibarial dilator is a rather unspecific feature which has also developed in other groups of insects (e.g., Antliophora; Beutel & Baum 2008). In his later review studies, Kristensen (e.g., 1995) questioned Hennig's Paraneoptera concept and treated acercarian and polyneopteran affinities of Zoraptera as competing working hypotheses.

Similarities of the antennae of Zoraptera and Isoptera (chemoreceptors, tactile setae, Johnston's organ) were described by Slifer & Sekhon (1978). However, it is evident that these features have evolved independently given the strong support for a subordinate placement of Isoptera within Blattodea (e.g., Deitz et al. 2003; Lo 2003; Klass 2009). A sistergroup relationship with Embioptera was first proposed by Minet & Bourgoin (1986), who suggested an entire series of potential synapomorphies including a reduced wing venation (groundplan), a hypertrophied metafemur and metathoracic tibial depressor, moniliform antennae, fusion of tarsomeres, loss of the arolium, reduction of the ovipositor, and gregarious habits. Even though some of these features are rather unspecific modifications or reductions, a close relationship with Embioptera gained further support by Engel & Grimaldi (2000) and Grimaldi (2001), like in Minet & Bourgoin (1986) based on informal character evaluations.

In a study on the zorapteran wing venation Kukalová-Peck & Peck (1993) addressed the intraordinal relationships and also the placement of the order within neopteran insects. Based on an informal analysis of wing characters they suggested that Zoraptera "probably diverged from the Blattoneoptera (= Grylloblattodea, †Protelytroptera, Dermaptera and Dictyoptera), almost certainly before the (†Protelytroptera + Dermaptera) line, and much before the (Isoptera + (Blattodea + Mantodea)) line". The hypothesis was illustrated with a phylogenetic

1 tree (Kukalová-Peck & Peck 1993: Fig. 23) showing Zoraptera as the second branch of
2 Blattoneoptera after Grylloblattodea.

3 A sistergroup relationship with Holometabola (“Scarabaeiformes”) was suggested by
4 Rasnitsyn (1998) based on the alleged presence of a precursor of the medial mesocoxal
5 articulation in *Z. hubbardi*. As pointed out above (2. Morphology), the morphological data
6 were insufficient. Moreover, the phylogenetic argumentation was based on an ad hoc
7 interpretation of a single vague character. Rasnitsyn (1998) provisionally placed Zoraptera in
8 an otherwise extinct superorder Caloneurida, supposedly subordinate to a “Cohors
9 Cimicoformes” (= Acercaria). He hypothesized Caloneurida “to be ancestral to other
10 cimicoform superorders as well as to the Cohors Scarabaeiformia” (= Holometabola). This
11 taxonomic treatment of Zoraptera is not compatible with the suggested phylogenetic position
12 as sistergroup of Holometabola.

13 Kusnetsova et al. (2002) examined the karyotype ($2n = 38, 36 + \text{neo-XY}$) and genital
14 structures of *Z. hubbardi*. Based on the presence of panoistic ovaries and holokinetic
15 chromosomes they rather vaguely suggested that Zoraptera may “represent a group of
16 Polyneoptera nearest to the origin of Paraneoptera”.

17 Zompro (2005) suggested a position of Zoraptera in “Orthopteromorpha”, supposedly
18 comprising the polyneopteran orders excluding Plecoptera, Embioptera and Phasmatodea.
19 This placement was based on elongate coxae and eggs without a strongly sclerotized egg
20 capsule and also lacking an operculum. However, a stringent character discussion and
21 analytical approach were lacking. Furthermore, the coxae of Zoraptera are not elongated (e.g.,
22 Friedrich & Beutel 2008) and the condition of the egg is obviously plesiomorphic (Mashimo
23 et al. 2011).

24 A character system previously not explored with respect to the position of Zoraptera is
25 the antennal heart and associated structures. The hitherto unknown antennal pulsatile organ of
26 Zoraptera is characterized by a t-shaped configuration of muscles (pers. obs. B. Wipfler), with

1 a fairly broad transverse muscle attached to the head capsule on both sides (assigned as
2 "Mxy" in Beutel & Weide 2005) and a narrower vertical muscle. Such a configuration occurs
3 in several groups of Polyneoptera but is completely unknown in Acercaria. This feature
4 clearly supports a placement of Zoraptera among the polyneopteran lineages.

5 Recent detailed morphological studies related to the reproduction suggested a
6 sistergroup relationship of Zoraptera with Embioptera or Eukinolabia (Embioptera +
7 Phasmatodea), by Mashimo et al. (2011, in press) based on the egg structure and development
8 of egg tooth and by Dallai et al. (2011, 2012b) based on the spermatozoan structures, i.e. 17
9 protofilaments comprising accessory tubules of axonemes and L-shaped electron-dense
10 lamellae accompanying microtubular triplets in the centriole adjunct.

12 **3.3. Cladistic approaches**

13 The first morphology-based formal cladistics analysis including all hexapod orders were
14 presented by Wheeler et al. (2001) and Beutel & Gorb (2001). Wheeler et al. (2001) was
15 primarily based on molecular data but also presented an independent morphological analysis
16 based on a matrix with 275 characters. The analysis of the morphological data yielded a
17 sistergroup relationship between Acercaria and Zoraptera in agreement with Hennig (1969).
18 Zoraptera were placed as sistergroup of Eumetabola (Acercaria + Holometabola) in the study
19 of Beutel & Gorb (2001), which had a main focus on the evolution of attachment structures.
20 The same result was obtained in a series of studies on the head morphology (Wipfler et al.
21 2011; Blanke et al. 2012; 2013; Wipfler 2012). However, it was pointed out by the authors
22 that the unit Zoraptera + Eumetabola was not supported by convincing synapomorphies. In a
23 follow-up study Beutel & Gorb (2006; expanded morphological data set, Mantophasmatodea
24 included) retrieved Zoraptera as sister to Acercaria, essentially supported by the potential
25 synapomorphies listed above (e.g., blade-like lacinia, condensed abdominal ganglionic chain).

Cladistic analyses of characters of the wing articulation (Yoshizawa 2007, 2011a) yielded a sistergroup relationship Zoraptera + Embioptera, like in Minet & Bourgoïn (1986). This was also supported when additional characters from Beutel & Gorb (2001, 2006) were added to the matrix. The fusion between the first axillary sclerite and the posterior notal wing process was emphasized as a potential synapomorphy, as this modification also involves a modification of the wing hinge, which is extremely conservative throughout the winged insects (Yoshizawa & Ninomiya 2007; Ninomiya & Yoshizawa 2009).

3.4. Molecular phylogenetic and phylogenomic approach

Wheeler et al. (2001; 18S, 28S, morphology) and Terry & Whiting (2005; 18S, 28S, Histone 3, morphology) analyzed the relationships of the entire Hexapoda or of the polyneopteran orders, respectively. Very different placements of Zoraptera order resulted from separate analyses of 18S and 28S rRNA in Wheeler et al. (2001: Figs 13, 14) (see Table 1), and analyses of both genes combined suggested an unlikely unit comprising Zoraptera, Trichoptera and Lepidoptera. Terry & Whiting (2005) placed Zoraptera as sistergroup of Dermaptera and referred to this possible clade as Haplocecata. The analytical methods used in both studies (POY) have been criticized (Kjer et al. 2007; Ogden & Rosenberg 2007; Yoshizawa 2010). A contaminant of a dermapteran sequence in the zorapteran dataset (Yoshizawa 2010, 2011b) suggests that the clade Zoraptera + Dermaptera may be artificial.

Yoshizawa & Johnson (2005; 18S rRNA) aligned their data using a secondary structure model. Maximum likelihood and Bayesian analyses retrieved Zoraptera as sister to Dictyoptera. However, very unusual molecular evolutionary trends observed in Zoraptera (e.g., extremely accelerated substitution rates and modifications of secondary structures) made the placement of the order highly unstable. Wang et al. (2013; 28S rRNA) also employed secondary structure-based alignments and obtained the same result for Zoraptera. However, for both 18S and 28S rRNA an extreme acceleration of the substitution rate and modification

of secondary structures is evident in Zoraptera, making the results unstable. Wang et al. (2013) identified unique features of the secondary structure of 28S rRNA shared by Zoraptera and Dictyoptera.

Misof et al. (2007; 18S rRNA) specifically addressed problems caused by character interdependence by employing secondary structure information and a Bayesian approach. A partly robust tree was obtained based on a hexapod consensus secondary structure model and mixed DNA/RNA substitution models. Zoraptera were placed as sistergroup of Dermaptera + Plecoptera, but with very low Bayesian support (0.45). This underlines that the frequently used 18S rRNA is apparently insufficient to fully resolve supraordinal insect interrelationships.

Ishiwata et al. (2011; DPD1, RPB1, RPB2) retrieved Polyneoptera incl. Zoraptera with good support, and a sistergroup relationship Zoraptera + Dictyoptera in some of the trees with low support. Consequently, Zoraptera were shown in an unresolved polyneopteran polytomy in a summary tree (Ishiwata et al. 2011).

Simon et al. (2012) placed Zoraptera as sister to all other polyneopteran terminals in a preliminary phylogenomic study. The support values were extremely low and several orders (Embioptera, Phasmatodea, Mantophasmatodea, Grylloblattodea) were not included. In a follow-up study (Letsch & Simon 2013) Embioptera and Phasmatodea were added. The analyses also supported a sistergroup relationship between Zoraptera and the remaining polyneopteran lineages. Recent unpublished results from the 1KITE project (ca. 100 hexapod terminals representing all orders) suggest a sistergroup relationship between a unit Zoraptera + Dermaptera (confirming the Haplocercata s. Terry & Whiting 2005) and a clade comprising all other polyneopteran orders. However, this pattern is not strongly supported statistically.

4. Intraordinal phylogeny

Kukalová-Peck & Peck (1993) suggested a classification based on an informal evaluation of wing characters. They recognized two families of Zoraptera and several genera. A new genus *Formosozoros* (from Taiwan) was described by Chao & Chen (2000). However, like the genera introduced by Kukalová-Peck & Peck (1993) it was not considered as valid in recent studies, and all extant species are currently classified into a single genus *Zorotypus* (e.g., Engel & Grimaldi 2002; Grimaldi & Engel 2005).

No molecular approach explicitly addressing the intraordinal phylogeny has been made to date. Yoshizawa & Johnson (2005; 18S rRNA) included four representatives of Zoraptera in their study (Old World and New World samples) and Yoshizawa (2010, 2011b; 18S rRNA) six. A deep divergence between the Old World and New World species was demonstrated. This is seemingly in conflict with evidence from male genitalia. The New World *Zorotypus snyderi* shares an elongated and coiled intromittent organ with the Old World species (Gurney 1938; Dallai et al. 2011; Mashimo et al. 2013), but molecular data suggest that it is closely related with *Z. hubbardi* (also New World) which lacks this unusual structure. Therefore it is conceivable that the coil was present in the groundplan of Zoraptera (or a large subgroup) and was secondarily lost in the majority of species. It cannot be fully excluded that it was acquired independently in several lineages, but independent gain of such a complex structure appears less likely.

5. Fossils

Grimaldi & Engel (2005) suggested a possible origin of Zoraptera in the lower Mesozoic. The first described extinct species was an apterous female of †*Zorotypus palaeus* from Dominican amber (Middle Miocene; Poinar 1988; Iturralde-Vinent & MacPhee 1996). The first winged specimen was described by Engel & Grimaldi (2000) from the same formation. Four species were identified in Burmese Cretaceous amber (Engel & Grimaldi, 2002). Three of them were very similar to extant zorapterans and consequently assigned to the genus *Zorotypus*. A basal

position of two of them (*†Zorotypus anathothorax*, *†Zorotypus nascimbenei*) was tentatively suggested by Engel (2003), and both were placed in a subgenus *†Octozoros* based on a reduced number of eight antennomeres and the presence of jugate setae along a part of the posterior forewing border (Engel, 2003). One of the four species was placed in a separate genus *†Xenozorotypus* (Engel & Grimaldi, 2002) and considered as sister to all other zorapterans. *†Palaeospinosus hudaie* was described from Jordanian amber by Kaddumi (2005), but the genus has been synonymized with *Zorotypus* (Engel 2008).

6. General biology

Zorapterans are omnivorous scavengers and feed on fungal hyphae and spores and on very small arthropods like mites and collembolans (Choe 1992; Grimaldi & Engel 2005). They are usually found in colonies under bark of decaying logs. As social behavior has not been reported yet and is probably missing in all species, zorapterans should be referred to as gregarious. The individuals spend much time grooming various body parts (Valentin 1986). In *Zorotypus barberi* some parts that cannot be reached are groomed by conspecifics (Choe 1992). Some species emit a slight but characteristic odor, reminiscent of a fragrance produced by the osmeterium of some swallowtail butterfly (Shetlar 1978; pers. obs. Y. Mashimo). Little is known about the life cycle of Zoraptera. The length of the nymphal period is about 1-2 months and adult lifespan is a few months, as reported by Gurney (1938) and Shetlar (1978). While Shetlar (1974) estimated five nymphal instars by measuring morphological features, Riegel & Eytalis (1974) suggested four instars based on different head widths. However, these descriptions are insufficient and the conclusions largely speculative. The specific wing dimorphism, a potential autapomorphy of the order, is also insufficiently investigated. According to observations mentioned in a review by Choe (1992) crowding and heredity seem to affect the production of the winged morphs. However, relevant details are unknown. Further intensive observations and investigations of the biology of Zoraptera are required.

7. Embryonic development

The embryonic development of Zoraptera was described recently by Mashimo et al, (in press). The embryo is formed by a fusion of paired blastoderm regions with higher cellular density extends along the egg surface. After reaching its full length, it migrates into the yolk and finally moves to take its position on the ventral surface of the egg accompanied by a reversion of its anteroposterior axis. These embryological features are widely known in Polyneoptera, and strongly suggest a placement of Zoraptera in this lineage (Mashimo et al. in press).

8. Mating behavior

Several studies on the mating behaviour were carried out by Choe (1994a, b, 1995, 1997), focused on the precopulatory courtship and copulation of *Z. gurneyi* and *Z. barberi* from Panama. The two sympatric species display a distinctly different mating behaviour. In *Z. gurneyi* a linear dominance hierarchy is established by mutual antennation and females were fertilized only once with very few exceptions. Dominant males performed 72% of the copulations, while the others were able to mate only occasionally. *Z. barberi* males lack this hierarchy. They are characterized by a highly unusual courtship ritual including nuptial feeding on a secretion of a cephalic gland (Choe 1997).

The mating behaviour of several Asian species was documented by Dallai et al. (2013) (Fig. 3). Mating sequences were recorded for many hours and observations were compared to earlier findings of Choe (1994a, b, 1995, 1997). An unusual case of external sperm transfer was described for *Z. impositus*, a feature unknown in other pterygote insects. The high variability not only of the genital structures but also of the mating behaviour and reproductive strategies was underlined by Dallai et al. (2013).

9. Discussion

1 Despite of their small size, inconspicuous appearance, cryptic habits and very low diversity it
2 would be misleading to consider Zoraptera as “unsuccessful” in their evolution. An origin in
3 the early Mesozoic was suggested by Grimaldi & Engel (2005). Considering a possible
4 placement as sistergroup of a large polyneopteran subunit (possibly together with
5 Dermaptera), an origin in the Carboniferous appears likely. In any case the group has survived
6 for a period of at least 200 Mya. Very small size and cryptic habits, which have prevented
7 discovery by insect collectors until the early 20th century, may have contributed to the long
8 survival. Recently collected and described species from different regions (e.g., Mashimo et al.
9 2013) indicate that the true diversity of the order is likely distinctly higher than known at
10 present.

11 The morphological investigation of the group mainly profited from three factors. The
12 availability of innovative techniques greatly facilitated detailed anatomical and ultrastructural
13 studies (Fig. 1; e.g., Friedrich & Beutel 2008; Dallai et al. 2012a, b). The coordinated effort of
14 researchers with different specialized skills also led to a markedly improved knowledge of the
15 group. Moreover, improved collecting and fixation yielded more and better conserved
16 material for anatomical investigations.

17 Unlike in Strepsiptera (“the Strepsiptera problem”; e.g., Kristensen 1991, 1995; Beutel
18 & Pohl 2006; Beutel et al. 2011) the systematic position of Zoraptera is not obscured by
19 numerous autapomorphies, but rather by many preserved plesiomorphic features, combined
20 with far-reaching reductions in some body regions (e.g., attachment structures, ovipositor).
21 Friedrich & Beutel (2008) showed that the thoracic skeleto-muscular system of winged
22 morphs is probably close to the groundplan of Neoptera. Whereas Strepsiptera have been
23 recently placed as the sistergroup of Coleoptera with strong support (Niehuis et al. 2012; Pohl
24 & Beutel 2013), the “Zoraptera problem” is still not completely resolved.

25 Widely divergent placements were suggested for Zoraptera (Figs 4, 5) and Strepsiptera
26 and in both cases early attempts were impeded by the lack of a sound phylogenetic

methodology (Pohl & Beutel 2013). That the spectrum of proposed positions is somewhat narrower in the case of Zoraptera may be partly due to its later discovery, about 120 years after the description of the first strepsipteran species (Rossius 1793; see Pohl & Beutel 2013). The placement of Zoraptera as close relatives of termites (e.g., Weidner 1969) was based on superficial similarities (see Weidner 1970) and an insufficient character evaluation. The structural affinities with Psocoptera emphasized by Crampton (e.g., 1921) and others are either plesiomorphic, superficial, or due to homoplasy. The placement of barklice (paraphyletic with respect to Liposcelididae) within a clade Psocodea is undisputed (e.g., Yoshizawa & Saigusa 2001; Yoshizawa & Johnson 2006, 2010; Friedemann et al. in press). The sistergroup relationship with Holometabola suggested by Rasnitsyn (1998) was based on an insufficient evaluation of very incomplete morphological data. A clade Zoraptera + Eumetabola (Beutel & Gorb 2001) is an artifact mainly caused by parallel reductions (e.g., ocelli in immature stages, number of Malpighian tubules).

The hypothesized sistergroup relationship between Zoraptera and Acercaria (Hennig 1953, 1969; Wille 1960; Kristensen 1975; Willmann 2003, 2005; Beutel & Weide 2005; Beutel & Gorb 2006) is presently largely refuted (Figs 4, 5). Most characters suggesting monophyletic Paraneoptera (incl. Zoraptera) are reductions (e.g., number of tarsomeres, cercomeres, and abdominal ganglia) and polyneopteran affinities (e.g., Yoshizawa 2007, 2011a; Ishiwata et al. 2011) imply that these structural modifications evolved independently.

A placement of Zoraptera within Polyneoptera is gaining more and more support (Fig. 5). However, the monophyly of this unit is not sufficiently supported yet (e.g., Whitfield & Kjer 2008; Trautwein et al. 2012) and the precise placement of Zoraptera is still controversial. A close relationship with Dictyoptera (Boudreaux 1979; Wheeler et al. 2001: p. 148 [“based on the discussion”]; Yoshizawa & Johnson 2005; Wang et al. 2013) is a possible option, but convincing synapomorphies are missing. Boudreaux’ (1979) arguments were not based on a formal character analyses and the features are unspecific or widespread in Insecta (e.g.,

“backwardly directed hind coxae”). The conclusion presented by Wheeler et al. (2001) in a
 summary tree is weakened by conflicting results obtained with the different data sets
 (morphology, 18S rRNA, 28S rRNA, 18S + 28S rRNA) (see Table 1). It is conceivable that
 the results based on molecular data were negatively affected by the use of POY (e.g.,
 Yoshizawa 2010, 2011b). Kukalová-Peck & Peck (1993) provided useful data, but the
 hypothesized placement of Zoraptera in a clade Blattoneoptera is weakened by the lack of a
 formal analysis and the exclusive use of wing characters. Wing base characters and
 morphological features linked to reproduction suggest a placement as sistergroup Embioptera
 (Yoshizawa 2007, 2011a) or Eukinolabia (Embioptera + Phasmatodea) (Mashimo et al. 2011,
 in press; egg and embryonic development; Dallai et al. 2011, 2012b; spermatozoa),
 respectively. Results of single gene analyses of single genes did not yield congruent results
 yet (e.g., Yoshizawa & Johnson 2005; Misof et al. 2007). Likewise, presently available
 transcriptome-based studies are not fully convincing. The basal placement within
 Polyneoptera suggested by Simon et al. (2012) and Letsch & Simon (2013) is weakened by
 the incomplete ordinal taxon sampling. As the pattern in the recent 1KITE-tree ((Zoraptera +
 Dermaptera) + (remaining Polyneoptera)) is not strongly supported statistically, the
 precise placement of Zoraptera remains a challenge.

Despite a considerable progress in zorapteran studies it is apparent that important
 problems remain to be solved, not only concerning the systematic position. To understand the
 phenomenon of different forms of selective pressure shaping different body parts (natural
 selection [ecological]/sexual selection) detailed information on genital structures of more
 species is required. Moreover, a species level phylogeny based on morphological *and*
 molecular data should have high priority in future studies. A solid intraordinal phylogeny is
 an essential basis for reconstructing the evolution of the group and to understand evolutionary
 processes.

It is probably safe to assume that zorapterans will remain one of the smallest hexapod orders and only few entomologists would address them as attractive insects. Nevertheless the group is intriguing in different aspects, phylogenetically, as a model case in evolutionary biology, and as cryptic survivors with a hidden diversity still to explore.

Acknowledgements

We thank G. Brehm for allowing YoM to join his field survey in Ecuador.

References

- Beutel, R.G. & Baum, E. (2008) A longstanding entomological problem finally solved? Head morphology of *Nannochorista* (Mecoptera, Insecta) and possible phylogenetic implications. *Journal of Zoological Systematics and Evolutionary Research* **46**: 346–367.
- Beutel, R.G., Friedrich, F., Hörnschemeyer, T., Pohl, H., Hünefeld, F., Beckmann, F., Meier, R., Misof, B., Whiting, M.F. & Vilhelmsen, L. (2011). Morphological and molecular evidence converge upon a robust phylogeny of the megadiverse Holometabola. *Cladistics* **27**: 341–355.
- Beutel, R.G. & Gorb, S.N. (2001) Ultrastructure of attachment specializations of hexapods (Arthropoda): Evolutionary patterns inferred from a revised ordinal phylogeny. *Journal of Zoological Systematics and Evolutionary Research* **39**: 177–207.
- Beutel, R.G. & Gorb, S.N. (2006) A revised interpretation of the evolution of attachment structures in Hexapoda with special emphasis on Mantophasmatodea. *Arthropod Systematics and Phylogeny* **64**: 3–25.
- Beutel, R.G. & Pohl, H. (2006) Endopterygote systematics – where do we stand and what is the goal (Hexapoda, Arthropoda)? *Systematic Entomology* **31**: 202–219.

- 1 Beutel, R.G. & Weide, D. (2005) Cephalic anatomy of *Zorotypus hubbardi* (Hexapoda:
2 Zoraptera): New evidence for a relationship with Acercaria. *Zoomorphology* **124**: 121–
3 136.
- 4 Blanke, A., Wipfler, B., Letsch, H., Koch, M., Beckmann, F., Beutel, R. & Misof, B. (2012)
5 Revival of Palaeoptera – head characters support a monophyletic origin of Odonata and
6 Ephemeroptera (Insecta). *Cladistics* **28**: 560–581.
- 7 Blanke, A., Greve, C., Wipfler, B., Beutel, R., Holland, B. & Misof, B. (2013) The
8 identification of concerted convergence in insect heads corroborates Palaeoptera.
9 *Systematic Biology* **62**: 250–263.
- 10 Bolívar y Pieltain, C. (1940) Estudio de un Nuevo Zoráptero de México. *Anales de la Escuela*
11 *Nacional de Ciencias Biologicas* **1**: 515–523.
- 12 Boudreaux, H.B. (1979) Arthropod Phylogeny, with Special Reference to Insects. John Wiley
13 & Sons: New York: viii+320 pp.
- 14 Caudell, A.N. (1918) *Zorotypus hubbardi*, a new species of the order Zoraptera from the
15 United States. *The Canadian Entomologist* **50**: 375–381.
- 16 Caudell, A.N. (1920) Zoraptera not an apterous order. *Proceedings of the Entomological*
17 *Society of Washington* **22**: 84–97.
- 18 Caudell, A.N. (1923) *Zorotypus swezeyi*, a new species of order Zoraptera from Hawaii.
19 *Transactions of the American Entomological Society* **48**: 133–135.
- 20 Caudell, A.N. (1927) *Zorotypus longicercatus*, a new species of Zoraptera from Jamaica.
21 *Proceedings of the Entomological Society of Washington* **29**: 144–145.
- 22 Chao, R.-F. & Chen, C.-S. (2000) *Formosozoros newi*, a new genus and species of Zoraptera
23 (Insecta) from Taiwan. *Pan-Pacific Entomologist* **76**: 24–27.
- 24 Choe, J.C. (1989) *Zorotypus gurneyi*, new species, from Panama and redescription of *Z.*
25 *barberi* Gurney (Zoraptera: Zorotypidae). *Annals of the Entomological Society of*
26 *America* **82**: 149–155.

- 1 Choe, J.C. (1992) Zoraptera of Panama with a review of the morphology, systematics, and
2 biology of the order. In: Quintero, D. & Aiello, A. (eds) *Insects of Panama and*
3 *Mesoamerica: Selected Studies*. Oxford University Press, Oxford: pp. 249–256.
- 4 Choe, J.C. (1994a) Sexual selection and mating system in *Zorotypus gurneyi* Choe (Insecta:
5 Zoraptera): I. Dominance hierarchy and mating success. *Behavioral Ecology and*
6 *Sociobiology* **34**: 87–93.
- 7 Choe, J.C. (1994b) Sexual selection and mating system in *Zorotypus gurneyi* Choe (Insecta:
8 Zoraptera): II. Determinants and dynamics of dominance. *Behavioral Ecology and*
9 *Sociobiology* **34**: 233–237.
- 10 Choe, J.C. (1995) Courtship feeding and repeated mating in *Zorotypus barberi* (Insecta:
11 Zoraptera). *Animal Behaviour* **49**: 1511–1520.
- 12 Choe, J.C. (1997) The evolution of mating systems in the Zoraptera: Mating variations and
13 sexual conflicts. In: Choe, J.C. & Crespi B.J. (eds) *The Evolution of Mating Systems in*
14 *Insects and Arachnids*. Cambridge University Press, Cambridge: pp. 130–145.
- 15 Crampton, G.C. (1918) The genitalia and terminal abdominal structures of male Neuroptera
16 and Mecoptera with notes on the Psocidae, Diptera and Trichoptera. *Psyche* **25**: 47–58.
- 17 Crampton, G.C. (1920) Some anatomical details of the remarkable winged zorapteron,
18 *Zorotypus hubbardis* [sic] Caudell, with notes on its relationships. *Proceedings of the*
19 *Entomological Society of Washington* **22**: 98–106.
- 20 Crampton, G.C. (1921) A further comparison of the wings of Zoraptera, psocids, and aphids,
21 from the standpoint of phylogeny. *The Canadian Entomologist* **53**: 110–117.
- 22 Crampton, G.C. (1926) A comparison of the neck and prothoracic sclerites through the orders
23 of insects from the standpoint of phylogeny. *Transactions of the American*
24 *Entomological Society* **52**: 199–248.
- 25 Crampton, G.C. (1927) The thoracic sclerites and wing bases of the roach *Periplaneta*
26 *americana* and the basal structures of the wings of insects. *Psyche* **34**: 59–72.

- 1 Dallai R., Gottardo, M., Mercati, D., Machida, R., Mashimo, Y. & Beutel, R.G. (2013)
2 Divergent mating patterns and a unique mode of external sperm transfer in Zoraptera:
3 An enigmatic group of pterygote insects. *Naturwissenschaften* **100**: 581–594.
- 4 Dallai, R., Mercati, D., Gottardo, M., Machida, R., Mashimo, Y. & Beutel, R.G. (2011) The
5 male reproductive system of *Zorotypus caudelli* Karny (Zoraptera): Sperm structure
6 and spermiogenesis. *Arthropod Structure and Development* **40**: 531–547.
- 7 Dallai, R., Mercati, D., Gottardo, M., Machida, R., Mashimo, Y. & Beutel, R.G. (2012a) The
8 fine structure of the female reproductive system of *Zorotypus caudelli* Karny
9 (Zoraptera). *Arthropod Structure and Development* **41**: 51–63.
- 10 Dallai, R., Mercati, D., Gottardo, M., Dossey, A.T., Machida, R., Mashimo, Y. & Beutel,
11 R.G. (2012b). The male and female reproductive systems of *Zorotypus hubbardi*
12 Caudell, 1918 (Zoraptera). *Arthropod Structure and Development* **41**: 337–359.
- 13 Deitz, L.L., Nalepa, C. & Klass, K.-D. (2003) Phylogeny of the Dictyoptera re-examined
14 (Insecta). *Entomologische Abhandlungen* **61**: 69–91.
- 15 Delamare-Deboutteville, C. (1947) Sur la morphologie des adultes aptères et ailés de
16 Zoraptères. *Annales des Sciences Naturelles - Zoologie et Biologie Animale* **11**: 145–
17 154.
- 18 Engel, M.S. (2000) A new *Zorotypus* from Peru, with notes on related neotropical species
19 (Zoraptera: Zorotypidae). *Journal of the Kansas Entomological Society* **73**: 11–20.
- 20 Engel, M.S. (2003) Phylogeny of the Zoraptera. *Entomologische Abhandlungen* **61**: 147–148.
- 21 Engel, M.S. (2008) A new apterous *Zorotypus* in Miocene amber from the Dominican
22 Republic (Zoraptera: Zorotypidae). *Acta Entomologica Slovenica* **16**: 127–136.
- 23 Engel, M.S. & Grimaldi, D.A. (2000) A winged *Zorotypus* in Miocene amber from the
24 Dominican Republic (Zoraptera: Zorotypidae), with discussion on relationships of
25 and within the order. *Acta Geológica Hispánica* **35**: 149–164.

- 1 Engel, M.S. & Grimaldi, D.A. (2002) The first Mesozoic Zoraptera (Insecta). *American*
2 *Museum Novitates* **3362**: 1–20.
- 3 Friedemann, K., Spangenberg, R., Yoshizawa, K. and Beutel, R.G. (in press) Evolution of
4 attachment structures in the highly diverse Acercaria. *Cladistics*. DOI:
5 10.1111/cla.12030
- 6 Friedrich, F. & Beutel, R.G. (2008) The thorax of *Zorotypus* (Hexapoda, Zoraptera) and a new
7 nomenclature for the musculature of Neoptera. *Arthropod Structure and Development*
8 **37**: 29–54.
- 9 Gottardo, M., Mercati, D. & Dallai, R. (2012). The spermatogenesis and sperm structure of
10 *Timema poppensis* (Insecta: Phasmatodea). *Zoomorphology* **131**: 209–223.
- 11 Grimaldi, D. (2001) Insect evolutionary history from Handlirsch to Hennig, and beyond.
12 *Journal of Paleontology* **75**: 1152–1160.
- 13 Grimaldi, D.A. & Engel, M.S. (2005) Evolution of the Insects. Cambridge University Press,
14 Cambridge: xv+755 pp.
- 15 Gurney, A.B. (1938) A synopsis of the order Zoraptera, with notes on the biology of
16 *Zorotypus hubbardi* Caudell. *Proceedings of the Entomological Society of Washington*
17 **40**: 57–87.
- 18 Hennig, W. (1953) Kritische Bemerkungen zum phylogenetischen System der Insekten.
19 *Beiträge zur Entomologie* **3**: 1–85.
- 20 Hennig, W. (1969) Die Stammesgeschichte der Insekten. Waldemar Kramer: Frankfurt am
21 Main: 436 pp.
- 22 Hubbard, M.D. (1990) A catalog of the order Zoraptera (Insecta). *Insecta Mundi* **4**: 49–66.
- 23 Hünefeld, F. (2007) The genital morphology of *Zorotypus hubbardi* Caudell, 1918 (Insecta:
24 Zoraptera: Zorotypidae). *Zoomorphology* **126**: 135–151.
- 25 Hwang, F.-S. (1974). *Zorotypus chinensis* [sic], a new species from China. *Acta*
26 *Entomologica Sinica* **17**: 423–427. [in Chinese, English summary].

- 1 Hwang, F.-S. (1976). A new species of Zoraptera. *Acta Entomologica Sinica* **19**: 225–227. [in
2 Chinese, English summary].
- 3 Ishiwata, K., Sasaki, G., Ogawa, J., Miyata, T. & Su, Z.-H. (2011) Phylogenetic relationships
4 among insect orders based on three nuclear protein-coding gene sequences. *Molecular*
5 *Phylogenetics and Evolution* **58**: 169–180.
- 6 Iturralde-Vinent, M.A. & MacPhee, R.D.E. (1996) Age and paleogeographic origin of
7 Dominican Amber. *Science* **273**: 1850–1852.
- 8 Kaddumi, H.F. (2005) Amber of Jordan, the Oldest Prehistoric Insects in Fossilized Resin.
9 Eternal River Museum of Natural History, Amman: 298 pp.
- 10 Karny, H.H. (1922) Zorapteren aus Süd-Sumatra. *Treubia* **3**: 14–37.
- 11 Karny, H.H. (1927) Spolia Mentawiensia. Zoraptera. *Treubia* **9**: 1–5.
- 12 Kjer, K.M., Gillespie, J.J. & Ober, K.A. (2007) Opinions on multiple sequence alignment, and
13 an empirical comparison of repeatability and accuracy between POY and structural
14 alignment. *Systematic Biology* **56**: 133–146.
- 15 Klass, K.-D. (2009) A critical review of current data and hypotheses on hexapod phylogeny.
16 *Proceedings of the Arthropodan Embryological Society of Japan* **43**: 3–22.
- 17 Kristensen, N.P. (1975) The phylogeny of hexapod "orders". A critical review of recent
18 accounts. *Journal of Zoological Systematics and Evolutionary Research* **13**: 1–44.
- 19 Kristensen, N.P. (1991) Phylogeny of extant hexapods. In: CSIRO (ed.) The Insects of
20 Australia. 2nd edn. Volume 1. Melbourne University Press, Carlton: pp. 125–140.
- 21 Kristensen, N.P. (1995) Forty years' insect phylogenetic systematics. *Zoologische Beiträge*,
22 *NF* **36**: 83–124.
- 23 Kukalová-Peck, J. & Peck, S.B. (1993) Zoraptera wing structures: Evidence for new genera
24 and relationship with the blattoid orders (Insecta: Blattoneoptera). *Systematic*
25 *Entomology* **18**: 333–350.

- 1 Kuznetsova, V.G., Nokkala, S. & Shcherbakov, D.E. (2002) Karyotype, reproductive organs,
2 and pattern of gametogenesis in *Zorotypus hubbardi* Caudell (Insecta: Zoraptera,
3 Zorotypidae), with discussion on relationships of the order. *Canadian Journal of*
4 *Zoology* **80**: 1047–1054.
- 5 Letsch, H. & Simon, S. (2013) Insect phylogenomics: new insights on the relationships of
6 lower neopteran orders (Polyneoptera). *Systematic Entomology* **38**, 783–793.
- 7 Lo, N. (2003) Molecular phylogenetics of Dictyoptera: Insights into the evolution of termite
8 eusociality and bacterial endosymbiosis in cockroaches. *Entomologische*
9 *Abhandlungen* **61**: 137–138.
- 10 Mashimo, Y., Beutel, R.G., Dallai, R., Lee, C.-Y. & Machida, R. (in press) Embryonic
11 development of Zoraptera with special reference to external morphology, and its
12 phylogenetic implications (Insecta). *Journal of Morphology*.
- 13 Mashimo, Y., Machida, R., Dallai, R., Gottardo, M., Mercati, D. & Beutel, R.G. (2011) Egg
14 structure of *Zorotypus caudelli* Karny (Insecta, Zoraptera, Zorotypidae). *Tissue and*
15 *Cell* **43**: 230–237.
- 16 Mashimo, Y., Yoshizawa, K., Engel M.S., Ghani, A.B., Dallai, R., Beutel, R.G. & Machida,
17 R. (2013). *Zorotypus* in Peninsular Malaysia (Zoraptera: Zorotypidae), with the
18 description of three new species. *Zootaxa* **3717**: 498–514
- 19 Minet, J. & Bourgoïn, T. (1986) Phylogenie et classification des Hexapodes (Arthropoda).
20 *Cahiers Liaison, OPIE* **20**: 23–28.
- 21 Misof, B., Niehuis, O., Bischoff, I., Rickert, A., Erpenbeck, D. & Staniczek, A. (2007)
22 Towards an 18S phylogeny of hexapods: Accounting for group-specific character
23 covariance in optimized mixed nucleotide/doublet models. *Zoology* **110**: 409–429.
- 24 New, T.R. (1978) Notes on neotropical Zoraptera, with descriptions of two new species.
25 *Systematic Entomology* **3**: 361–370.

- 1 New, T.R. (1995) The order Zoraptera (Insecta) from Christmas Island, Indian Ocean.
2 *Invertebrate Taxonomy* **9**: 243–246.
- 3 Niehuis, O., Hartig, G., Grath, S., Pohl, H., Lehmann, J., Tafer, H., Donath, A., Krauss, V.,
4 Eisenhardt, C., Hertel, J., Peterson, M., Mayer, C., Meusemann, K., Peters, R.S.,
5 Stadler, P.F., Beutel, R.G., Bornberg-Bauer, E., McKenna, D.D. & Misof, B. (2012)
6 Genomic and morphological evidence converge to resolve the enigma of Strepsiptera.
7 *Current Biology* **22**: 1309–1313.
- 8 Ninomiya, T. & Yoshizawa, K. (2009) A revised interpretation of the wing base structure in
9 Odonata. *Systematic Entomology* **34**: 334–345.
- 10 Ogden, T.H. & Rosenberg, M.S. (2007) Alignment and topological accuracy of the direct
11 optimization approach via POY and traditional phylogenetics via ClustalW + PAUP*.
12 *Systematic Biology* **56**: 182–193.
- 13 Paulian, R. (1949). Découverte de l'ordre des Zoraptères à Madagascar et description d'une
14 nouvelle espèce. *Mémoires de l'Institut Scientifique de Madagascar (A)* **3**: 77–80.
- 15 Paulian, R. (1951). Découverte de l'ordre des Zoraptères à l'île Maurice. *Zorotypus vinsoni*,
16 n. sp. *Le Naturaliste Malgache* **3**: 33–35.
- 17 Pohl, H. & Beutel, R.G. (2013) The Strepsiptera-Odyssey - the history of the systematic
18 placement of an enigmatic parasitic insect order. *Entomologia*. DOI: 10.4081
19 /entomologia.2013.e4
- 20 Poinar Jr., G.O. (1988) *Zorotypus palaeus*, new species, a fossil Zoraptera (Insecta) in
21 Dominican amber. *Journal of the New York Entomological Society* **96**: 253–259.
- 22 Rafael, J.A., Engel, M.S. (2006) A new species of *Zorotypus* from central Amazonia, Brazil
23 (Zoraptera: Zorotypidae). *American Museum Novitates* **3528**: 1–11.
- 24 Rafael, J.A., Godoi, F.S.P. & Engel, M.S. (2008) A new species of *Zorotypus* from eastern
25 Amazonia, Brazil (Zoraptera: Zorotypidae). *Transactions of the Kansas Academy of*
26 *Science* **111**: 193–202.

- 1 Rasnitsyn, A.P. (1998) On the taxonomic position of the insect order Zorotypida = Zoraptera.
2 *Zoologischer Anzeiger* **237**: 185–194.
- 3 Riegel, G.T. (1963) The distribution of *Zorotypus hubbardi* (Zoraptera). *Annals of the*
4 *Entomological Society of America* **56**: 744–747.
- 5 Riegel, G.T. & Eytalis, S.J. (1974) Life history studies on Zoraptera. Proceedings North
6 Central Branch, Entomological Society of America **29**: 106–107.
- 7 Rossius, P. (1793) Observation de M. Rossi sur un nouveau genre d’Insecte, voisin
8 d’Ichneumons. *Bulletin de la Societé Philomatique de Paris* **1**: 49.
- 9 Shetlar, D.J. (1974) The biology, morphology, and taxonomy of *Zorotypus hubbardi*
10 Caudell (Insecta: Zoraptera). Master’s thesis, The University of Oklahoma.
- 11 Shetlar, D.J. (1978) Biological observations on *Zorotypus hubbardi* Caudell (Zoraptera).
12 *Entomological News* **89**: 217–223.
- 13 Silvestri, F. (1913) Descrizione di un nuovo ordine di insetti. *Bollettino del Laboratorio di*
14 *Zoologia Generale e Agraria, Portici* **7**: 193–209.
- 15 Silvestri, F. (1946) Descrizione di due specie neotropicali di *Zorotypus* (Insecta, Zoraptera).
16 *Bollettino del Laboratoria di Entomologia Agraria, Portici* **7**: 1–12.
- 17 Simon, S., Narechania, A., DeSalle, R. & Hadrys, H. (2012) Insect phylogenomics: Exploring
18 the source of incongruence using new transcriptomic data. *Genome Biology and*
19 *Evolution* **4**: 1295–1309.
- 20 Slifer, E.H. & Sekhon, S.S. (1978) Structures on the antennal flagellum of *Zorotypus*
21 *hubbardi* (Insecta, Zoraptera). *Notulae Natural. Academy of Natural Sciences of*
22 *Philadelphia* **453**: 1–8
- 23 Snodgrass, R.E. (1935) Principles of insect morphology. McGraw-Hill, New York: 667 pp.
- 24 Terry, M.D. & Whiting, M.F. (2005) Mantophasmatodea and phylogeny of the lower
25 neopterous insects. *Cladistics* **21**: 240–257.

- 1 Terry, M.D. & Whiting, M.F. (2012) *Zorotypus novobritannicus* n. sp., the first species of the
2 order Zoraptera (Zorotypidae) from the Australasian Ecozone. *Zootaxa* **3260**: 53–61.
- 3 Trautwein, M.D., Wiegmann, B.M., Beutel, R., Kjer, K.M., Yeates, D.K. (2012) Advances in
4 insect phylogeny at the dawn of the postgenomic era. *Annual Review of Entomology* **57**:
5 449–468.
- 6 Valentine, B.D. (1986) Grooming behavior in Embioptera and Zoraptera (Insecta). *The Ohio*
7 *Journal of Science* **86**: 150–152.
- 8 Wang, Y., Engel, M.S., Rafael, J.A., Dang, K., Wu, H., Wang, Y., Xie, Q. & Bu, W. (2013) A
9 unique box in 28S rRNA is shared by the enigmatic insect order Zoraptera and
10 Dictyoptera. *PLOS ONE* **8**: e53679.
- 11 Weber, H. (1933) Lehrbuch der Entomologie. Gustav Fischer Verlag, Jena: 726 pp.
- 12 Weidner, H. (1969) Die Ordnung Zoraptera oder Bodenläuse. *Entomologische Zeitschrift* **79**:
13 29–51.
- 14 Weidner, H. (1970) 15. Zoraptera (Bodenläuse). In: Helmcke, J.-G., Starck, D. & Wermuth, H.
15 (eds) Handbuch der Zoologie. Bd. IV: Arthropoda. 2. Hälfte: Insecta, 2. Teil:
16 Spezielles. Walter de Gruyter, Berlin: pp. 1–12.
- 17 Wheeler, W.C., Whiting, M., Wheeler, Q.D. & Carpenter, J.M. (2001) The phylogeny of the
18 extant hexapod orders. *Cladistics* **17**, 113–169.
- 19 Whitfield, J.B. & Kjer, K.M. (2008) Ancient rapid radiations of insects: Challenges for
20 phylogenetic analysis. *Annual Review of Entomology* **53**: 449–472.
- 21 Wille, A. (1960) The phylogeny and relationships between the insect orders. *Revista de*
22 *Biología Tropical*, **50** 93–123.
- 23 Willmann, R. (2003) Die phylogenetischen Beziehungen der Insecta: Offene Fragen und
24 Probleme. *Verhandlungen Westdeutscher Entomologentag* **2001**: 1–64.

- 1 Willmann, R. (2005) Phylogenese und Systemk der Insecta. In: Dathe, H. H. (ed.) Lehrbuch
2 der Speziellen Zoologie, begründet von A. Kaestner, 2. Aufl. Bd. I: Wirbellose Tiere. 5.
3 Teil: Insecta. Spektrum, Gustav Fischer, Heidelberg, Berlin: pp. 1–65.
- 4 Wipfler, B. (2012) Polyneopteran head morphology and its phylogenetic implications.
5 Unpubl. PhD thesis, Biologisch Pharmazeutische Fakultät, FSU Jena.
- 6 Wipfler, B., Machida, R., Müller, B. & Beutel, R.G. (2011) On the head morphology of
7 Grylloblattodea (Insecta) and the systematic position of the order, with a new
8 nomenclature for the head muscles of Dicondylia. *Systematic Entomology* **36**: 241–266.
- 9 Wipfler, B. (2012). Polyneopteran head morphology and its phylogenetic implications.
10 Unpubl. PhD thesis, Biologisch Pharmazeutische Fakultät, FSU Jena.
- 11 Yoshizawa, K. (2007) The Zoraptera problem: Evidence for Zoraptera + Embiodea from the
12 wing base. *Systematic Entomology* **32**: 197–204.
- 13 Yoshizawa, K. (2010) Direct optimization overly optimizes data. *Systematic Entomology* **35**:
14 199–206.
- 15 Yoshizawa, K. (2011a) Monophyletic Polyneoptera recovered by wing base structure.
16 *Systematic Entomology* **36**: 377–394.
- 17 Yoshizawa, K. (2011b) Problems in direct optimization. *Taxa, Proceedings of the Japanese*
18 *Society of Systematic Zoology* **30**: 17–29 (in Japanese with English summary).
- 19 Yoshizawa, K. & Johnson, K.P. (2005) Aligned 18S for Zoraptera (Insecta): Phylogenetic
20 position and molecular evolution. *Molecular Phylogenetics and Evolution* **37**: 572–
21 580.
- 22 Yoshizawa, K & Johnson, K.P (2006) Morphology of male genitalia in lice and their relatives
23 and phylogenetic implications. *Systematic Entomology* **31**: 350–361.
- 24 Yoshizawa, K., Johnson, K .P. (2010) How stable is the "polyphyly of lice" hypothesis
25 (Insecta: Psocodea)? A comparison of phylogenetic signal in multiple genes.
26 *Molecular Phylogenetics and Evolution* **55**: 939–951.

- 1 Yoshizawa, K. & Ninomiya, T. (2007) Homology of the wing base sclerites in Ephemeroptera
2 (Insecta: Pterygota)-A reply to Willkommen and Hörnschemeyer. *Arthropod Structure*
3 *and Development* **36**: 277–279.
- 4 Yoshizawa, K. & Saigusa, T. (2001) Phylogenetic analysis of paraneopteran orders (Insecta:
5 Neoptera) based on forewing base structure, with comments on monophyly of
6 Auchenorrhyncha (Hemiptera). *Systematic Entomology* **26**: 1–13.
- 7 Zompro, O. (2005) Inter- and intra-ordinal relationships of the Mantophasmatodea, with
8 comments on the phylogeny of polyneopteran orders (Insecta: Polyneoptera).
9 *Mitteilungen aus dem Geologisch-Paläontologischen Institut der Universität*
10 *Hamburg* **89**: 85–116.

1 Table 1. Systematic placements of Zoraptera with different approaches and character sets.

Systematic approach/ Study	Characters	Systematic placement
Pre-Hennigian concepts		
Silvestri (1913)	morphology	closely related with roaches and termites (Blattodea)
Caudell (1918)	morphology	affinities with Isoptera
Crampton (1920, 1921, 1926)	different morphological character systems	affinities with Psocoptera ("Corrodentia")
Weber (1933)	morphology	tentatively in a superorder Blattoidea (Mantodea, Zoraptera, Blattaria, Isoptera)
Gurney (1938)	morphology	affinities with Psocoptera ("Corrodentia")
Hennigian concepts		
Hennig (1953)	morphological characters	tentatively as basal lineage of Paraneoptera , outside of Psocoptera
Hennig (1969)	morphological characters, abdominal ganglionic chain, number of tarsomeres	sistergroup of Acercaria (Paraneoptera concept)
Weidner (1969)	mainly thoracic characters	tentatively as sistergroup of Isoptera
Weidner (1970)	mainly thoracic characters	obsolete branch of Blattodea
Kristensen (1975)	different morphological	Paraneoptera concept as

	characters	working hypothesis, polyneopteran affinities tentatively refuted
Boudreaux (1979)	wing venation, metacoxae	sistergroup of Dictyoptera („Cursorida = Zorapterida + Blattarida“
Minet et Bourgoin (1986)	wing venation, thoracic and abdominal morphology	sistergroup of Embioptera
Kristensen (1995)	different morphological characters	either acercarian or polyneopteran affinities
Kukalová-Peck & Peck (1997)	wing base and venation	sistergroup of †Protelytroptera, Dermaptera and Dictyoptera (Blattoneoptera excl. Grylloblattodea)
Kusnetsova et al. (2002)	chromosome structure and genitalia	“in Polyneoptera close to the origin of Paraneoptera”
Zompro (2005)	morphology and egg structure	“Orthopteromorpha” (=Polyneoptera excl. Plecoptera, Embioptera and Phasmatodea)
Dallai et al. (2011)	sperm ultrastructure	In a clade with Embioptera and Phasmatodea
Mashimo et al. (2011, in press)	egg structure and development	In a clade with Embioptera and Phasmatodea

Cladistic approach based on morphology		
Kukalová-Peck & Peck (1993)	wing base and venation	sistergroup of †Protelytroptera, Dermaptera and Dictyoptera (Blattoneoptera)
Wheeler et al. (2001)	morphological matrix with 275 characters, extracted from literature	sistergroup of Acercaria
Beutel & Gorb (2001)	characters of adults and larvae including attachment structures	sistergroup of Eumetabola (Acercaria + Holometabola)
Beutel & Gorb (2006)	characters of adults and larvae including attachment structures, Mantophasmatodea and some developmental features added	sistergroup of Acercaria
Yoshizawa (2011)	wing base	Embioptera
Single gene analyses		
Wheeler et al. (2001)	18S rRNA (analyzed with POY)	Sistergroup of Psocodea
	28S rRNA (POY)	sistergroup of a clade comprising all pterygote

		groups except for Strepsiptera and two ephemeropteran terminals
	18S + 28S rRNA	sistergroup of Amphiesmenoptera (trichopteran and lepidopteran terminals)
	18S, 28S rRNA + morphological data extracted from literature (“total evidence”)	sistergroup of Dictyoptera
Yoshizawa & Johnson (2005)	18S rRNA	Dictyoptera
Misof et al. (2007)	18S rRNA, special focus on character interdependence	sistergroup of Plecoptera + Dermaptera
Ishiwata et al. (2011)	3 nuclear protein-coding genes	In Polyneoptera , possibly sistergroup of Dictyoptera
Wang et al. (2013)	28S rRNA	Dictyoptera
Phylogenomic approach		
Simon et al. (2012)	Transcriptomes, Embioptera, Phasmstodea, Grylloblattodea, and Mantophasmatodea not included	sistergroup of remaining polyneopteran terminals with low support
Letsch & Simon (2013)	Transcriptomes,	sistergroup of remaining

	Grylloblattodea and Mantophasmatodea not included	polyneopteran terminals
--	---	-------------------------

1

1 Fig. 1. Histogram showing number of publications on Zoraptera under different aspects.

2
3
4 2 Fig. 2. Collecting site of different *Zorotypus* species in Ecuador, Zamora-Chinchipe,
5
6 3 Copalinga Lodge Reserve Rainforest, with nymph (lower right).

7
8
9
10 4 Fig. 3. Copula of *Zorotypus caudelli*, volume render of μ -Ct image stacks.


11
12
13 5 Fig. 4. Cladograms showing different placements of Zoraptera.

14
15
16
17 6 Fig. 5. Cladograms from recent studies showing different placements of Zoraptera.

18
19
20
21 7

22
23
24 8

Figure



Figure

[Click here to download high resolution image](#)



Figure

[Click here to download high resolution image](#)







Figure **A Hennig 1969**
Morphology: discussion




B Kristensen 1981, 1991
Morphology: discussion




C Beutel & Gorb 2001
Morphology: cladistic analysis




D Wheeler et al. 2001
18S rDNA, 28S rDNA & morphology




E Grimaldi & Engel 2005
Morphology: discussion




F Willmann 2005
Morphology: discussion




G Yoshizawa & Johnson 2005
18S rDNA




H Terry & Whiting 2005
18S rDNA, 28SrDNA, Histone3
& morphology (PAUP)




I Terry & Whiting 2005
18S rDNA, 28SrDNA, Histone3
& morphology (ClusterX)




J Beutel & Gorb 2006
Morphology: cladistic analysis




K Misof et al. 2007
18S rDNA




M Yoshizawa 2011
Morphology: cladistic analysis




M Yoshizawa 2011
Morphology (wing joint):
cladistic analysis




N Ishiwata et al. 2011
DPD1, RPB1, RPB2




O Wipfler 2012
Cephalic morphology:
cladistic analysis



P Simon et al. 2012
Transcriptomic data



Q Blanke et al. 2012
Cephalic morphology:
cladistic analysis



R Letsch & Simon in press
Transcriptomic data

