HOKKAIDO UNIVERSITY

Title	Human health risks from metals and metalloid via consumption of food animals near gold mines in Tarkwa, Ghana: Estimation of the daily intakes and target hazard quotients (THQs)
Author(s)	Bortey-Sam, Nesta; Nakayama, Shouta M. M.; Ikenaka, Yoshinori et al.
Citation	Ecotoxicology and environmental safety, 111, 160-167 https://doi.org/10.1016/j.ecoenv.2014.09.008
Issue Date	2015-01
Doc URL	https://hdl.handle.net/2115/57933
Type	journal article
File Information	Ecotox.Environ.Safe_v.111.pdf

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

Human health risks from metals and metalloid via consumption of food animals near Gold Mines in Tarkwa, Ghana: Estimation of the daily intakes and target hazard quotients (THQs)

Nesta Bortey-Sam[1‡], Shouta M.M. Nakayama[1‡], Yoshinori Ikenaka[1], Osei Akoto[2], Elvis Baidoo[2], Yared Beyene Yohannes[1], Hazuki Mizukawa[1], Mayumi Ishizuka[1*]

[1]Graduate School of Veterinary Medicine, Hokkaido University, Kita 18, Nishi 9, Kita ku, Sapporo 060-0818, Japan

[2]Department of Chemistry, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana

‡ The authors contributed equally to this study.

* Corresponding author: Mayumi ISHIZUKA, e-mail: ishizum@vetmed.hokudai.ac.jp

Laboratory of Toxicology, Department of Environmental Veterinary Sciences, Graduate School of Veterinary Medicine, Hokkaido University, N18, W9, Kita-ku, Sapporo 060-0818, Japan Tel: +81-11-706-6949; Fax: +81-11-706-5105.

Abstract

Heavy metal and metalloid contamination in food resulting from mining is of major concern due to the potential risk involved. Food consumption is the most likely route for human exposure to metals. This study was therefore to estimate the daily intake and health risk (based on target hazard quotients, THQ) from metals via consumption of free-range chicken, goat and sheep near gold mines in Tarkwa, Ghana. The concentrations of Cr, Mn, Fe, Co, Ni, Cu, Zn, As, Cd, and Pb were measured with an inductively coupled plasma-mass spectrometer and Hg analysis was done using the mercury analyzer. The mean concentrations of metals ranged from nd–542 mg/kg wet weight. Principal component analysis of the results showed a clear separation between chicken, grouped on one side, and the ruminants clustered on another side in both offal and muscle. Interestingly, As, Cd, Hg, Mn and Pb made one cluster in the offal of chicken. Chicken muscle also showed similar distribution with As, Hg and Pb clustered together. The daily intake of As (µg/kg body weight/day) were in the following ranges; [0.002 (kidneys of goat and sheep)–0.19 (chicken gizzard)], Cd [0.003 (chicken muscle)–0.55 (chicken liver)], Hg [0.002 (goat muscle)–0.29 (chicken liver)], Pb [0.01 (muscles and kidneys of goat and sheep)–0.96 (chicken gizzard)] and Mn [0.13 (goat kidney)–8.92 (sheep liver)]. From the results, daily intake of As, Cd, Hg, Pb and Mn in these food animals were low compared to the provisional tolerable daily intake guidelines. The THQs although less than one, indicated that contributions of chicken gizzard and liver to toxic metal exposure in adults and especially children could be significant.

Keywords: Offal, Muscle, Estimated daily intake, Target hazard quotient, Ghana

1. Introduction

Heavy metals and metalloid (arsenic) are ubiquitous in the environment either naturally or anthropogenically, and their concentrations were elevated promptly through waste disposal, smelter stacks, atmospheric deposition, mining, fertilizer and pesticide use and the application of sewage sludge in arable land (Cui et al., 2005). Mining and processing metal ore can be a significant source of metal and metalloid contamination of the environment (Navarro et al., 2008; Singh et al., 2005). The increased occurrence of metal and metalloid pollution in the environment has been associated with anthropogenic activities as effluents and emissions from mines and smelters often contain elevated concentrations of toxic metals including arsenic, cadmium, mercury and lead (Tubaro and Hungerford, 2007).

Food consumption has been identified as the major pathway of human exposure, accounting for >90% compared to other ways of exposure such as inhalation and dermal contact (Loutfy et al., 2006). In a review by Yabe et al. (2010), contamination of food animals, fish, soil, water, and vegetables with heavy metals has reached unprecedented levels over the past decade in some parts of Africa. As a result, human exposure to toxic metals has become a major health risk.

Tarkwa (05°18′00″N; 01°59′00″W) is a town in the southwest of Ghana, located about 120 miles west of the capital city, Accra. As of 2010, Tarkwa was estimated to have a population of 90,477 (Ghana Statistical Service, 2010). It is a noted centre of gold and manganese mining. Tarkwa mine, which is a large open-cast gold mine, is situated to the northwest of the town, and Nsuta manganese mine is situated to the east. Tarkwa has nearly a century of gold mining history and has the largest concentration of mining companies in a single district in Ghana and the West African sub-region (Akabzaa and Darimani, 2001). A study conducted by Hayford et al. (2008) on the impact of the gold mining on soil and staple foods collected around mining communities in Tarkwa showed high levels of some toxic metals including arsenic and mercury in the soil and staple foods. In Tarkwa, the extraction and processing of gold has given rise to various environmental related diseases and accidents. According to the District Medical Officer of Health, the common mining-related diseases observed in the area over the years include, but are not limited to: vector-borne diseases such as malaria, schistomiasis and onchocerciasis; respiratory tract diseases,

81 especially pulmonary tuberculosis and silicosis; skin diseases; eye diseases, especially
82 acute conjunctivitis; accidents resulting from galamsey activities, and mental cases
83 (Akabzaa and Darimani, 2001). The aim of this study was therefore to determine the
84 concentrations of heavy metals and metalloid in offal and muscle of free-range chicken,
85 goat and sheep; to reveal the species differences in accumulation features of metals among
86 free-range chicken, goat and sheep; and to estimate the daily intake (EDI) and target hazard
87 quotient (THQ) (toxic metals) through consumption of these food animals in Tarkwa.
88

2. Materials and methods

2.1. Sampling and pre-treatment

In June 2012, liver, kidney and muscle samples were collected at slaughterhouses from both male and female local goat and sheep (n = 10 species each) aged three years and above, and stored in labeled plastic bags in cooler boxes. 10 free-range chickens were also bought from townships surrounding the two major gold mines. All the townships were close to the mine, and some households were within 1 km of the mine. The live adult chickens were transported to the laboratory for dissection after exsanguination. Liver, kidney, muscle and gizzard samples were collected. Samples were transported to the laboratory and kept frozen at the Chemistry Department of the Kwame Nkrumah University of Science and Technology, KNUST, Ghana before being transported and analyzed for metal concentrations at the Laboratory of Toxicology, Graduate School of Veterinary Medicine, Hokkaido University, Japan.

2.2. Analysis of total Mercury (Hg)

The concentration of total mercury (Hg) was measured by thermal decomposition, gold amalgamation and atomic absorption spectrophotometry (Mercury analyzer, MA-3000; Nippon Instruments Corporation, Tokyo, Japan), after preparation of the calibration standard. Recovery rates of Hg for the certified reference material, DOLT-4 (Dogfish liver, the National Research Council, Canada) ranged from 92% to 103% (94.3 ± 4.2%). The detection limit (Hg) was 2.0 pg of total Hg.

2.3. Sample preparation and metal analysis

Approximately 0.5 g of individual samples were dried in an oven at 40 ^{0}C and digested in a closed microwave extraction system, Speed Wave MWS-2 microwave digestion (Berghof, Germany). Briefly, the dried samples were placed in prewashed digestion vessels followed by acid digestion using 5 mL of (65%) nitric acid, HNO_3 (Kanto Chemical Corp., Tokyo, Japan) and 1 mL of (30%) hydrogen peroxide, H_2O_2 (Kanto Chemical Corp., Tokyo, Japan). The digestion vessels were capped and placed into a 10-position turntable

conditions followed by a ramped temperature programme: ramp to 160 ^{0}C (5 min hold); and increase to 190 ^{0}C (15 min hold). After cooling, samples were transferred into plastic tubes and diluted to a final volume of 10 mL with milli Q water. A reagent blank was prepared using the same procedure. An Inductively Coupled Plasma-Mass Spectrometer (ICP-MS; 7700 series, Agilent technologies, Tokyo, Japan) was used for quantification. The instrument was calibrated using standard solutions of the respective metals (to establish standard curves before metal analysis). All chemicals and standard stock solutions were of analytical-reagent grade (Wako Pure Chemicals). The detection limits (μg/L) of chromium (Cr), manganese (Mn), iron (Fe), cobalt (Co), nickel (Ni), copper (Cu), zinc (Zn), arsenic (As), cadmium (Cd), and lead (Pb) were 0.003, 0.025, 0.154, 0.0004, 0.024, 0.007, 0.226, 0.002, 0.001 and 0.001 respectively. Concentrations of metals were expressed in mg/kg wet weight (mg/kg ww)

2.4. Quality assurance and quality control

For heavy metals, replicate blanks and the reference materials DORM-3 (Fish protein, the National Research Council, Canada) and DOLT-4 were used for method validation and quality control. Replicate analysis of these reference materials showed good accuracy (relative standard deviation, RSD, ≤ 3%) and recovery rates ranged from 80% to 115%.

2.5. Data analysis

2.5.1 Estimated daily intake (EDI) of toxic metals

The estimated daily intake (EDI) of toxic metals (As, Cd, Hg, Pb and Mn) depended on both the metal concentration in offal or muscle and the amount of consumption of the respective food. The EDI of metals was determined by the following equation:

$$EDI = \frac{MC \times FDC}{BW}$$

where MC is average concentration of metal in food (μg/g, on fresh weight basis); FDC represents the average food daily consumption of offal and muscle in this region (g/person/d); BW is the average body weight. It was assumed that the local inhabitants consumed an average liver, muscle and gizzard of 150 and 100 g/day for adult (60 kg in BW) and children (30 kg in BW) respectively. For kidney, it was assumed that inhabitants

consumed 10 and 8 g/day for adults and children respectively (Ihedioha and Okoye, 2013), because the size of kidneys are generally smaller and is not favorite compared with liver, muscle, or gizzard. The metal intakes were compared with the tolerable daily intakes for metals recommended by the FAO/WHO (2010), WHO (2000) and FSA (2006).

2.5.2 Target Hazard Quotient (THQ)

The health risks from consumption of offal and muscle of chicken, goat and sheep by the local inhabitants were assessed based on the THQ. The THQ is a ratio of determined dose of a pollutant to a reference dose level. If the ratio is less than 1, the exposed population is unlikely to experience obvious adverse effects. The method of estimating risk using THQ was provided in the U.S. EPA Region III risk-based concentration table (USEPA IRIS, 2007) and in Chien et al., (2002), and is based on the equation below:

$$THQ = \frac{EFr \times ED \times FI\ R \times MC}{Rf\,D \times BW \times AT} \times 10^{-3}$$

where THQ is target hazard quotient; EFr is exposure frequency (365 days/year); ED is exposure duration (70 years); FIR is food ingestion rate (g/person/d); MC is average concentration of metal in food (µg/g, on fresh weight basis); RfD is the oral reference dose (mg/kg/d); BW is the average body weight, adult (60 kg); children (30 kg); AT is the average exposure time (365 days/year × number of exposure years, assuming 70 years in this study). Oral reference doses were based on 3E-04, 1E-03, 5E-04, 4E-03, 0.14 mg/kg/d for As, Cd, Hg, Pb and Mn respectively (USEPA, 1997; USEPA IRIS, 2007).

2.6. Statistical analysis

IBM SPSS 20.0 Statistical software (SPSS Inc., Illinois, USA) was used to perform ANOVA analysis followed by Tukey's test. ANOVA and Tukey analyses were used to compare metal concentrations in the liver, kidney, muscle and gizzard and differences were considered statistically significant with p value < 0.05. Principal component analysis (PCA) was done to determine the distribution pattern of metals in organs using JMP statistical software v. 10 (SAS Institute).

3. Results and discussion

3.1. Levels of heavy metals and metalloid

Concentrations of Cr, Mn, Fe, Co, Ni, Cu, Zn, As, Cd, Hg, and Pb in the liver, kidney muscle and gizzard of free-range chickens, as well as metal and metalloid levels in liver, kidney and muscle of goat and sheep from Tarkwa were determined (Table 1). Table 1 shows that the mean Cd concentrations were in the range of nd–0.73 mg/kg ww in all samples while the mean concentrations of Pb were in the range of nd–0.39 mg/kg ww, Cr (0.01–0.24), Mn (0.15–3.57), Fe (6.62–542.31), Co (nd–0.11), Ni (0.01–0.55), Cu (0.35–105.8), Zn (5.12–65.89), As (nd–0.14) and Hg (nd–0.12) mg/kg ww. Cu and Zn were detected in 100% of all offal and muscle samples for all the food animal studied.

Ni can cause respiratory problems and is a carcinogen. The permissible limit of Ni in food according to WHO (2000) standard is 0.5 mg/kg. In this study concentrations of Ni ranged between 0.01–0.55 mg/kg ww, in chicken liver and muscle respectively. As shown in Table 1, the highest mean concentration of Ni was found in the local chickens while Zn levels in goat muscle exceeded the permissible limit of 50 mg/kg in meat (USDA, 2006; OJEC, 2001; European Commission Regulation, 2006).

3.1.1. Arsenic

As was detectable in 100%, 83%, 80% and 52% of all gizzard, liver, kidney and muscle samples respectively. Concentrations of As in muscle and offal samples of free-range chicken, goat and sheep ranged from nd to 0.14 ± 0.11 mg/kg ww in goat muscle and chicken kidney respectively (Table 1). The As distribution in liver, kidney and muscle of chicken was found to be higher and differ statistically with that of goat and sheep ($p < 0.05$). However, there was no significant difference when As levels in goat offal and muscle was compared with sheep (Table 1, $p > 0.05$). Mean concentration of As in the kidney (0.14 ± 0.11 mg/kg ww) and gizzard (0.08 ± 0.05 mg/kg ww) of free-range chicken from Tarkwa were found to be higher than the USDA (2006); OJEC (2001); European Commission Regulation (2006) and SAC/MOHC, 2005 standard of 0.05 mg/kg ww in food. The high levels of As in the free-range chicken may be due to the nature of the gold bearing ore

206 which is mineralized pyrites and arsenopyrates. Processing of the ore involves roasting and
207 this results in the production of arsenic trioxide gas which is distributed throughout the
208 study area by air current. As is a toxic substance and due to its non-biodegradable nature, it
209 could accumulate in surface soil and water. It could also enter the food chain through plant
210 assimilation (Amonoo-Neizer et al., 1995; Griffis et al., 2002). This could be the possible
211 reason for the high As concentrations in free-range chickens from Tarkwa as they
212 continuously pick food particles and water from the ground.
213
214 *3.1.2. Cadmium*
215 Mean level of Cd in chicken kidney (0.73 ± 0.81 mg/kg ww) was within the
216 recommended residual concentrations of 0.5 to 1.0 mg/kg ww in offal for human
217 consumption (European Commission, 2001). Cd was detectable in 14% of all muscle,
218 100% of all liver and kidney, 90% of chicken gizzard. The concentrations of Cd in livers
219 and kidneys of chicken were significantly higher compared to those of goat and sheep
220 (Tukey test; $p < 0.05$). The higher frequencies of Cd in liver and kidney (100%) are likely
221 due to their special functions; liver as storage organ and kidney as excretory organ (Stoyke
222 et al., 1995). There have also been suggestions that animals exposed to Cd accumulate it in
223 their kidneys because of the presence of free protein-thiol groups which leads to a strong
224 fixation of heavy metals (Pompe-Gotal and Crnic, 2002). The likely unidentified sources of
225 Cd which the animals could come in contact with include municipal waste, electroplating,
226 plastic and paint waste. Other sources include leachates from nickel–cadmium based
227 batteries and cadmium plated items which are so carelessly discarded by battery chargers
228 and users in Ghana. Recently, electronic wastes are disposed and often burnt at refuse
229 dumps.
230
231 *3.1.3. Mercury*
232 Hg is a neurotoxic poison that causes neurobehavioral effects (especially on
233 psychomotor coordination), neuroendocrine, and renal damage and gastrointestinal toxicity
234 (SAC/MOHC, 2005). Hg was detectable in 38% of all muscle, 69% and 97% of all liver
235 and kidney respectively, and 100% of chicken gizzard. The levels of Hg in free-range

236 chicken from Tarkwa were higher compared to that of goat and sheep. The result of this
237 study indicated that chicken kidney (0.12 ± 0.08 mg/kg ww) and liver (0.11 ± 0.07 mg/kg
238 ww) contained the highest concentrations of Hg followed by sheep kidney (0.07 ± 0.07
239 mg/kg ww). The concentrations of Hg in liver and muscle of free-range chicken from
240 Tarkwa was found to be higher and differed statistically with that of goat and sheep (Tukey
241 test; $p < 0.05$). However, there was no significant difference when Hg levels in goat offal
242 and muscle was compared with sheep (Tukey test; $p > 0.05$). The lowest and highest mean
243 concentrations of Hg in chicken were 0.01 ± 0.01 mg/kg ww in muscle and 0.12 ± 0.08
244 mg/kg ww in kidney, respectively; goat (nd–0.03 ± 0.02 in muscle and kidney
245 respectively); sheep (nd–0.07 ± 0.07 in the muscle and kidney respectively) (Table 1). The
246 USDA (2006); OJEC (2001); and European Commission Regulation (2006) permissible
247 limit for Hg is 0.05 mg/kg and the Standardization Administration of China/Ministry of
248 Health of China established maximum limits of Hg in food to be 0.01 to 0.05 mg/kg
249 (SAC/MOHC, 2005). The high levels of Hg especially in free-range chicken could be
250 problematic, as concentration levels exceeded the maximum values permitted in food. In
251 Ghana, amalgamation is the preferred gold recovery method employed by almost all
252 artisanal gold miners because it is a very simple, inexpensive and an easier to use technique.
253 These activities by local and small scale miners are widespread, and Hg is introduced into
254 the environment (via amalgamation process popularly known as "galamsey") (Amonoo-
255 Neizer et al., 1995). The high levels of Hg in offal of free-range chicken could therefore be
256 due to contamination from soil, feed or water.
257
258 *3.1.4. Lead*
259 Excess Pb is known to reduce the cognitive development and intellectual performance in
260 children and to increase blood pressure and cardiovascular disease incidence in adults
261 (Commission of the European Communities, 2001). Pb was detectable in all samples of
262 liver, kidney, 80% of chicken gizzard and 38% of muscle. The result of this study indicated
263 that chicken gizzard (0.39 ± 1.17 mg/kg ww) contained the highest concentrations of Pb
264 followed by chicken kidney (0.26 ± 0.31 mg/kg ww) and liver (0.13 ± 0.14 mg/kg ww).
265 This is because, to be toxic, Pb must be in a form which will be retained in the gizzard, thus

resulting in continuous absorption over a prolonged period (Salisbury et al., 1958). In another study by Kendall et al. (1996), analyses of field and laboratory data indicated that two scenarios can be envisioned when Pb is ingested: (1) the ingested grit/pellets containing Pb might be eliminated (via regurgitation or passage through the gastrointestinal tract) before any significant dissolution or absorption of Pb occurs; (2) the grit/pellets may be partially or totally dissolved in the gizzard, in which case causes Pb accumulation resulting in the occurrence of a range of toxicologic effects.

The levels of Pb in organs of free-range chickens may emanate mainly from contamination of soil, feeds and/ or water sources. Tukey's test was used to compare the levels of Pb in the three food animals, and results showed no statistical significance in liver and muscle when chicken was compared with goat and sheep ($p > 0.05$). In this study, concentration of Pb in some organs of chicken exceeded the USDA (2006); OJEC (2001); and European Commission Regulation (2006) standard of 0.5 and 0.1 mg/kg ww for Pb in offal and muscle respectively (Table 1). Generally, the low levels of Pb in the offals and muscles of the mentioned food animals could be due to the fact that, Pb accumulates in bone and the metal concentration increases with age (Caggiano et al., 2005; Rubio et al., 1998).

3.1.5. Manganese

The role of Mn in neurosychiatric disorders is also documented (Schild, 1980; Jackson and Morris, 1989). Mn was detectable in 100% of all offal and muscle samples. This study showed that mean concentrations of Mn ranged from 0.15–3.51 (chicken); 0.62–2.43 (goat) and 0.71–3.57 mg/kg ww (sheep) (Table 1). The concentration of Mn in kidney of free-range chicken was found to be higher and differed statistically with that of goat and sheep ($p < 0.05$). However, there was no significant difference when Mn levels in the muscle of the three food animals were compared (Table 1, $p > 0.05$). In the liver, the levels of Mn between free-range chickens and sheep was statistically insignificant with mean values of 3.50 ± 0.92 and 3.57 ± 1.02 mg/kg ww respectively (Table 1, $p > 0.05$). The mean concentrations of Mn in offal and muscles of all three food animals were above the WHO (1996) reference standard of 0.5 mg/kg except in chicken muscle (Table 1). The high Mn

levels in these food animals could be due to the fact that the sampling site at Tarkwa was close to the Nsuta Mn mine (which is a famous site for Mn mining).

This study was compared with similar studies in different countries (Table 2). Results from this study was found to be comparable with study by Husain et al. (1996), in goat and sheep livers and kidneys, in Kuwait, but higher than similar study by Villar et al. (2005) in the liver of poultry from the Philippines. Study done by Yabe et al. (2013) near a Pb and Zn mine in Kabwe, Zambia indicated that the levels of Cd and Pb in the livers and kidneys of free-range chickens were higher compared to this study but levels in muscle were comparable. On the other hand, the levels of Pb and Cd in chicken gizzard from this study was higher than the study by Yabe et al. (2013). From Table 2, the mean concentrations of Zn in this study was comparable in all organs except for gizzard when compared to the study by Yabe et al. (2013).

3.2. Distribution of metals

ANOVA was used to compare the accumulation and distribution of metals in the organs of free-range chicken, goat and sheep. The metal distribution in offal and muscle of chicken was statistically significant ($p < 0.05$) except for Cr, As and Pb. There was also statistical significance ($p < 0.01$) in the distribution of Mn, Co, Cu, Zn and Cd when the organs of goat were compared. On the other hand metal distribution in organs of sheep were significant except for As ($p > 0.05$). The liver and kidneys are target tissues for monitoring metal contamination in animals because both organs function in removing toxic metals from the body (Husain et al., 1996; Abou-Arab, 2001). Similarly, organs of all species in this study showed that the liver and kidney contained the highest levels of metals (Table 1).

Distribution of metals in offal and muscle samples in the food animals was analyzed using PCA. PCA results (Fig. 1) showed a clear separation between chicken (C), grouped on one side, and the ruminants, goat (G), and sheep (S), clustered on another side in both offal and muscle. Interestingly, As, Cd, Hg, Mn and Pb made one cluster or group in the liver and kidney of free-range chicken (Fig. 1). Chicken muscle also showed similar distribution pattern, with As, Hg and Pb clustered together (Fig. 1). This could be attributed to the difference in their feeding habits and/ or the different levels of metallothionein (MT)

326 from the sampled food animals (species differences in hepatic MT content). MT is a low
327 molecular weight protein in various mammalian and non-mammalian tissues and is
328 involved in the detoxification of metals (Margoshes and Valle, 1957; Kagi and Vallee,
329 1960). According to study by Bryan-Henry et al. (1994), the hepatic MT levels in these
330 three food animals studied decreased in order of livestock > non-mammals; i.e. goat >
331 sheep > chicken.
332
333 *3.3. Dietary intake of metals and target hazard quotient*
334 The estimated daily intakes (EDI) of toxic metals for adults in the vicinity of Tarkwa
335 mine through consumption of offal and meat of free-range chicken, goat and sheep are
336 presented in Table 3. The highest EDIs of As (0.19 µg/kg bw/day), Pb (0.96 µg/kg bw/day),
337 Cd (0.55 µg/kg bw/day) and Hg (0.29 µg/kg bw/day), through consumption were from the
338 gizzard and liver of chicken (Table 3). Meanwhile, the highest EDI of Mn (8.92 µg/kg
339 bw/day) was through consumption of sheep liver (Table 3). The calculated EDIs of As, Cd,
340 Hg, Pb and Mn in offal and muscle samples of the 3 food animals from Tarkwa were low
341 compared to the FAO/WHO (2010), WHO (2000) and FSA (2006) tolerable daily intakes
342 (Table 3). Children are especially vulnerable to acute, sub-acute and chronic effects of
343 ingestion of chemical pollutants, since they (children) consume more (twice of the amount)
344 of food per unit of body weight as adults (ENHIS, 2007). As a result intakes of these toxic
345 metals through food could be higher for children from Tarkwa.
346 The health risk from consumption of offal and muscle from the three food animals were
347 assessed based on the THQ. THQ value less than 1 means the exposed population is
348 unlikely to experience obvious adverse effects (USEPA, 2000). Although the calculated
349 individual THQ were below 1.0 (Fig. 2), attention should be paid particularly to some high
350 individual THQ values approaching 1.0 (As Cd, and Hg). The THQ for As in chicken
351 gizzard and liver which is largely consumed was higher in children (0.85 and 0.84
352 respectively) than in adults (0.63 in both cases). Similarly, the THQ for Cd in the liver of
353 chicken was 0.74 for children compared with 0.55 for adults (Fig. 2). As shown in Fig. 2,
354 THQ for Hg in the liver of chicken was 0.76 and 0.57 in children and adults respectively. In
355 spite of this, caution must be taken since perennial intake of these contaminated food

356 animals is likely to induce adverse health effects arising largely from As, Cd, Hg and Pb
357 exposure. Meanwhile, the THQ for Mn was very low among all the three food animals
358 (data not shown) ranging from 0.001–0.06 and 0.001–0.008 in adults and children
359 respectively. The THQ for As, Cd, Hg and Pb showed that contributions of chicken gizzard
360 and liver to toxic metal exposure in adults and especially children could be significant.
361

4. Conclusion

The dietary intakes of As, Cd, Hg, Pb and Mn from the offal and muscle of free-range chicken, goat and sheep in Tarkwa, Ghana were low when compared to the FAO/WHO (2010), WHO (2000) and FSA (2006) tolerable daily intakes. Children are especially vulnerable to acute, sub-acute and chronic effects of ingestion of chemical pollutants, since they (children) consume more (twice of the amount) of food per unit of body weight as adults (ENHIS, 2007). As a result intakes of these four toxic metals through food could be higher for children from Tarkwa. THQ of As, Cd and Hg showed that there could be potential health risk for inhabitants through consumption of larger quantities of contaminated offal of free-range chicken. Prolonged consumption of these toxic metals in the offal and muscle may lead to accumulation and cause toxicity and therefore there is a clear need to avoid consumption of these contaminated food animals, as well as restrict them from roaming and scavenging for food and/or water near the mining areas. Furthermore, continuous monitoring of As, Cd, Hg and Pb residues (in these food animals) in the vicinities of gold mines in Tarkwa is recommended in the interest of consumers.

Acknowledgement

This study was supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology of Japan awarded to M. Ishizuka (24248056, 24405004) and Y. Ikenaka (26304043) and the foundation of JSPS Core to Core Program (AA Science Platforms). We would like to acknowledge the financial support by the Mitsui & Co., Ltd. Environment Fund, The Akiyama Life Science Foundation and The Nihon Seimei Foundation. We also express our sincere gratitude and thanks to Mr Joseph Prah, Mr Joseph Addae and Mr. Takahiro Ichise who in various ways assisted to carry out this research.

388 **References**

389 Abou-Arab, A.A.K., 2001. Heavy metal contents in Egyptian meat and the role of detergent
390 washing on their levels. Food Chem. Toxicol. 39 (6), 593-599.

391 Akabzaa, T., Darimani, A., 2001. Impact of mining sector investment in Ghana: a study of
392 the Tarkwa mining region. Health Impacts 6.4 pp 60.

393 Amonoo-Neizer, E.H., Nyamah, D., Bakiamoh, S.B., 1995. Mercury and Arsenic Pollution
394 in soil and biological samples around mining towns if Obuasi, Ghana. Water, Air and
395 Soil Poll. 91, 363-373.

396 Bryan Henry, R., Liu, J., Choudhuri, S., Klaassen, D.C., 1994. Species variation in hepatic
397 metallothionein. Toxicol lett. 74, 23-33.

398 Caggiano, R., Sabia, S., D'Emilio, M., Macchiato, M., Anastasio, A., Ragosta, M., Paino,
399 S., 2005. Metal levels in fodder, milk, dairy products, and tissues sampled in ovine
400 farms of Southern Italy. Environ. Res. 99, 48-57.

401 Chien, L.C., Hung, T.C., Choang, K.Y., Choang, K.Y., Yeh, C.Y., Meng, P.J., et al. 2002.
402 Daily intake of TBT, Cu, Zn, Cd and As for fishermen in Taiwan. Sci. Total Environ.
403 285, 177–185.

404 Commission of the European Communities, 2001. Commission Regulation (EC) No.
405 221/2002 of 6 February 2002 amending regulation (EC) No. 466/2002 setting
406 maximum levels for certain contaminants in foodstuffs. Official Journal of the
407 European Communities, Brussels, 6 February 2002.

408 Cui, Y., Zhu, Y., Zhai, R., Huang, Y., Qiu, Y., Liang, J., 2005. Exposure to metal mixtures
409 and human health impacts in a contaminated area in Nanning, China. Environ. Int. 31,
410 784–790.

411 Demirbas, A., 1999. Proximate and heavy metal composition in chicken meat and tissues.
412 Food Chem. 6 (1), 27-31.

413 European Commission, 2001. Setting maximum levels for certain contaminants in
414 foodstuffs. Commission Regulation 466.Off J. Eur. Commun. L77, 1–13.

415 European Commission Regulation, 2006. No 1881/2006, Setting maximum levels for
416 certain contaminants in foodstuff. Official J. Eur. Union, L 364: 5-24.

417 European Environment and Health Information System (ENHIS), 2007. Exposure of
418 children to chemical hazards in food. Fact Sheet No. 4.4, CODE: RPG4_Food_EX1.
419 World Health Organization.
420 FAO/WHO, 2010. Summary report of the seventy-third meeting of JECFA. Joint
421 FAO/WHO Expert Committee on Food Additives. Geneva.
422 FSA (Food Standard Agency), 2006. Metals and other elements in processed fish and fish
423 products: Food Survey Information Sheet 08/06,
424 http://www.food.gov.uk/science/surveillance.
425 Ghana Statistical Service, 2010. Population and Housing census.
426 https://www.google.co.uk/#q=Ghana+Statistical+Service%2C+2010.+Population+an
427 d+Housing+census+ pp 93.
428 Griffis, R.J., Barning, K., Agezo, F.L., Akosah, F.K., 2002. Gold deposits of Ghana.
429 Minerals Commission of Ghana, Accra.
430 Hayford, E.K., Amin, A., Osae, E.K., Kutu, J., 2008. Impact of gold mining on soil and
431 some staple foods collected from selected mining communities in and around Tarkwa
432 Prestea Area. West African J. Appl. Ecol. 14, 1–12.
433 Husain, A., Al-Rashdan, A., Al-Awadhi, A., Mahgoub, B., Al-Amiri, H., 1996. Toxic
434 metals in food products originating from locally reared animals in Kuwait. Bull
435 Environ. Contam. Toxicol. 57, 549–555.
436 Ihedioha, J.N., Okoye, C.O.B., 2013. Dietary intake and health risk assessment of lead and
437 cadmium via consumption of cow meat for an urban population in Enugu State,
438 Nigeria, Ecotox. Environ. Safe. 93, 101–106.
439 Jackson, H.M., Morris, G.P., 1989. Environmental health reference book. Hartonolls Ltd.
440 Bodmin.
441 Kagi J.H.R., Vallee, B.L., 1960. Metallothionein: a cadmium- and zinc-containing protein
442 from equine renal cortex. J. Biol. Chem. 235, 3460–3465.
443 Kendall, R.J., Lacher, T.E., Bunck, C., Daniel, B., Driver, C., Grue, C.E., Leighton, F.,
444 Stansley, W., Watanabe, P.G., Whitworth, M., 1996. An ecological risk assessment of
445 lead shot exposure in non-waterfowl avian species: Upland game birds and raptors.
446 Environ. Toxicol. Chem. 15, (1), 4–20.

447 Loutfy, N., Fuerhacker, M., Tundo, P., Raccanelli, S., El Dien, A.G., Ahmed, M.T., 2006.
448 Dietary intake of dioxins and dioxin-like PCBs, due to the consumption of dairy
449 products, fish/seafood and meat from Ismailia city, Egypt. Sci. Tot. Environ. 370, 1–8.
450 Margoshes, M., Valle, B.L., 1957. A cadmium protein from equine kidney cortex. J. Am.
451 Chem. Sot. 799, 4813-4814.
452 Navarro, M.C., Perez-Sirvent, C., Martinez-Sanchez, M.J., Vidal, J., Tovar, P.J., Bech, J.,
453 2008. Abandoned mine sites as a source of contamination by heavy metals: a case
454 study in a semi-arid zone. J. Geochem. Explor. 96, 183–193.
455 Official J. of the Eur. Communities (OJEC), 2001. L 77/1 COMMISSION REGULATION
456 (EC) No 6/200. Setting maximum levels for certain contaminants in foodstuffs.
457 Okoye, C.O.B., Ugwu, J.N., 2010. Impact of environmental cadmium, lead, copper and
458 zinc on quality of goat meat in Nigeria. Bull. Chem. Soc. Ethiop. 24 (1), 133–138.
459 Pompe-Gotal, J., Crnic, A.P., 2002. Cadmium in tissue of roe deer (Capreolus capreolus) in
460 Croatia. Vet. Arh. 72, 303–310
461 Rubio, M.R., Sigrist, M.E., Encinas, E.T., Baroni, E., Coronel, J.E., Boggio, J.C., et al.
462 1998. Cadmium and lead levels in cow's milk from a milking region in region in
463 Santa Fe, Argentine. Bull. Environ. Contam. Toxicol. 60, 164–167.
464 SAC/MOHC (The Standardization Administration China/Ministry of Health, China). 2005.
465 Maximum levels of contaminants in foods (GB2762-2005). Beijing.
466 Salisbury, R.M., Staples, E.L.J., Sutton, M., 1958. Lead poisoning of chickens, New. Zeal.
467 Vet. J. 6, 1, 2-7.
468 Schild, H.O., 1980. Applied Pharmacology. 12th. The Eng. Lang.bk.soc and Churchill
469 Livington. Lon. 465
470 Singh, A.N., Zeng, D.H., Chen, F.S., 2005. Heavy metal concentrations in redeveloping
471 soil of mine spoil under plantations of certain native woody species in dry tropical
472 environment, Indian J. Environ. Sci. 1, 168–174.
473 Stoyke, M., Doberschutz, K.D., Lusky, K., 1995. Heavy metal contents (cadmium, lead and
474 mercury) in selected feedstuff, organs and tissue in cattle found different sites of
475 Brandenburg. Mengen. Spureneler. Arb. 15, 269–276.

476 Tubaro, A., Hungerford, J., 2007. Environmental toxicology. In Gupta RC, ed, Veterinary
477 Toxicology: Basic and clinical principles. Academic, New York, NY, USA, pp. 663–
478 725.
479 USEPA, Integrated Risk Information System-database (IRIS), 2007. Philadelphia PA;
480 Washington, DC.
481 USEPA, 2000. Risk-based concentration table. Philadelphia PA: United States
482 Environmental Protection Agency, Washington DC;
483 USEPA, Integrated Risk Information System-database; 1997. Philadelphia PA; Washington,
484 DC.
485 USDA, 2006. Foreign Agricultural Service GAIN Report Global Agriculture Information
486 Network Voluntary Report-public distribution GAIN Report Number CH 6064, China
487 Peoples Republic of FAIRS Product. Specific maximum levels of contaminants in
488 foods, Jim Butterworth and Wu Bugang.
489 Uluozlua, O.D., Tuzen, M., Mendil, D., Soylak, M., 2009. Assessment of trace element
490 contents of chicken products from turkey. J. Hazard. Mater. 163, 982–987.
491 Villar, T.C., Elaine, J., Kaligayahan, P., Flavier, M.E., 2005. Lead and cadmium levels in
492 edible internal organs and blood of poultry chicken. J. Appl. Sci. 5 (7), 1250–1253.
493 WHO, 2000. Evaluation of certain food additives and contaminants. Report of the Fifty-
494 Third of the Joint FAO/WHO Expert Committee on Food Additives. Technical
495 Report Series No. 896. Geneva.
496 WHO, 1996. Trace Elements in Human Nutrition and Health. Geneva.
497 Yabe, J., Ishizuka, M., Umemura, T., 2010. Current levels of heavy metal pollution in
498 Africa. J. Vet. Med. Sci. 72, 1257–1263.
499 Yabe, J., Nakayama, M.M.S., Ikenaka, Y., Muzandu, K., Choongo, K., Mainda, G., Mabeta,
500 M., Ishizuka, M., Umemura, T., 2013. Metal distribution in tissues of free-range
501 chickens near a lead–zinc mine in Kabwe, Zambia. Environ. Toxicol. Chem. 32 (1),
502 189–192.
503

Figure captions:

Fig. 1. Distribution patterns of metals in food animals characterized by PCA (C: chicken; G: goat and S: sheep).

Fig. 2. Target Hazard Quotient (THQ) of As, Cd, Hg and Pb in children and adults (C: chicken; G: goat; S: sheep; G: gizzard; K: kidney; L: liver and M: muscle).

Table 1

Mean concentrations and range of metals (mg/kg ww) in various organs of free range chicken (C), goat (G) and sheep (S).

Sample		Cr	Mn	Fe	Co	Ni	Cu	Zn	As	Cd	Hg	Pb
CG(n=10)	Mean±SD	0.06±0.08	0.64±0.46	81.4±87.2	nd	0.10±0.04	1.92±0.81	30.3±3.5	0.07±0.05	0.08±0.09	0.01±0.01	0.38±1.17
	Range	0.01-0.27	0.20-1.75	37.1-321.4	nd-0.01	0.06-0.19	0.74-3.19	24.3-34.9	0.02-0.16	nd-0.30	0.01-0.03	nd-3.70
CK(n=10)	Mean±SD	0.24±0.40	2.44±0.55 [a]	94.4±33.2	0.03±0.01	0.07±0.14	2.82±0.37	28.4±3.1	0.14±0.11 [a]	0.72±0.81 [a]	0.12±0.08 [a]	0.25±0.31 [a]
	Range	0.01-1.35	1.65-3.59	59.7-150.5	0.01-0.07	nd-0.46	2.01-3.38	25.2-35.6	0.02-0.37	0.16-2.26	0.02-0.28	0.02-1.05
GK(n=10)	Mean±SD	0.01±0.00	0.77±0.25 [b]	88.8±72.8	0.07±0.03	0.14±0.07	2.89±0.46	20.0±3.9	0.01±0.01 [b]	0.06±0.04 [b]	0.03±0.02 [b]	0.03±0.04 [b]
	Range	nd-0.02	0.50-1.35	39.2-278.5	0.01-0.12	0.05-0.27	2.42-3.76	15.0-27.5	nd-0.04	0.01-0.13	0.01-0.07	0.01-0.14
SK(n=10)	Mean±SD	0.01±0.00	1.14±0.27 [b]	49.1±15.4	0.03±0.02	0.09±0.04	3.170±0.45	22.2±4.1	0.01±0.01 [b]	0.15±0.15 [b]	0.07±0.08 [ab]	0.03±0.01 [b]
	Range	nd-0.01	0.87-1.70	24.8-80.6	0.01-0.06	0.04-0.15	2.58-3.90	18.1-28.1	nd-0.03	0.01-0.43	nd-0.23	0.01-0.06
CL(n=10)	Mean±SD	0.02±0.01	3.50±0.92 [a]	542.3±604.0	0.03±0.01	0.01±0.02	3.67±0.67	65.8±50.6	0.07±0.07 [a]	0.22±0.18 [a]	0.11±0.07 [a]	0.13±0.14 [a]
	Range	nd -0.04	1.57-4.84	72.0-1629.3	0.02-0.05	nd-0.06	2.72-4.74	32.5-182.2	0.01-0.21	0.05-0.56	0.05-0.25	0.01-0.41
GL(n=10)	Mean±SD	0.02±0.01	2.42±0.71 [b]	89.9±47.7	0.11±0.05	0.20±0.05	60.3±44.7	47.1±26.6	nd [b]	0.02±0.01 [b]	0.01±0.01 [b]	0.02±0.04 [b]
	Range	0.01-0.04	1.06-3.41	44.3-212.6	0.04-0.21	0.15-0.29	3.3-140.7	21.9-103.3	nd-0.01	0.01-0.04	nd-0.05	0.01-0.13
SL(n=10)	Mean±SD	0.02±0.01	3.57±1.02 [a]	126.3±71.3	0.07±0.03	0.33±0.13	105.8±74.3	43.6±9.6	0.01±0.01 [b]	0.05±0.02 [b]	0.03±0.03 [b]	0.06±0.03 [ab]
	Range	0.01-0.04	1.50-4.92	55.4-294.1	0.04-0.12	0.22-0.68	7.9-224.3	31.7-63.3	0.01-0.05	0.02-0.09	nd-0.09	0.02-0.11
CM(n=10)	Mean±SD	0.05±0.01	0.14±0.08 [a]	6.6±1.6	nd	0.55±0.30	0.34±0.03	5.1±1.7	0.04±0.03 [a]	nd [a]	0.01±0.01 [a]	0.01±0.03 [a]
	Range	0.02-0.09	0.09-0.38	4.7-9.5	nd	0.24-1.29	0.28-0.41	3.8-10.0	nd-0.13	nd	0.01-0.02	nd-0.10
GM(n=10)	Mean±SD	0.05±0.08	0.6±0.81 [a]	31.7±17.0	0.02±0.05	0.14±0.05	1.50±0.76	51.3±17.6	nd [b]	nd [a]	nd [b]	nd [a]
	Range	0.01-0.28	0.21-2.83	19.2-76.8	nd-0.15	0.08-0.23	0.91-3.38	17.0-78.2	nd-0.01	nd-0.02	nd	nd-0.01
SM(n=10)	Mean±SD	0.03±0.03	0.71±0.55 [a]	37.8±10.9	0.01±0.00	0.46±0.16	2.09±1.19	45.9±20.8	0.01±0.01 [b]	0.01±0.01 [a]	nd [b]	0.01±0.00 [a]
	Range	0.01-0.09	0.16-2.02	22.7-58.9	nd-0.01	0.15-0.66	0.84-4.85	27.0-83.4	nd-0.02	nd-0.03	nd-0.02	nd-0.01

n: number of samples; C: chicken; G: goat; S: sheep; K: kidney; L: liver; M: muscle.

different letters (a and b) between animals in the same organ indicates significant difference (Tukey's test; $p < 0.05$).

nd: not detected.

Table 2

Published data of mean metal levels (mg/kg dw) in organs of different animal groups from different regions.

Study/Country	Pollution Factor	Animal	Organ	Cr	Co	Ni	Cu	Zn	As	Cd	Hg	Pb
Yabe et al., 2013*, Zambia	mining	C (broiler)	L	0.06	0.05	0.03	56.80	28.60		0.00		0.06
		C(free-range)	L	0.07	0.09	0.07	3.40	30.90		1.60		4.15
			K	0.08	0.16	0.08	2.12	24.70		3.50		7.62
			M	0.08	0.01	0.02	nd	4.16		0.01		0.23
			G	0.10	0.04	0.03	0.13	30.50		0.02		0.23
Uluozlua et al., 2009*, Turkey	markets		L	0.04	0.02	0.01	12.10	22.50	0.06	2.24		0.12
Demirbas, 1999*, Turkey	control		L				2.95	28.10		0.04	0.08	0.09
Villar et al., 2005, Phillipines			L							0.03		0.08
Husain et al., 1996, Kuwait	wholesalers	G	L							0.05		0.13
			K							0.44		0.43
Okoye and Ugwu, 2010, Nigeria	Industrial & urban		L				134.02	120.44		0.35		0.65
Abou-Arab, 2001*, Egypt	Industrial & urban		L				46.6	32.6		0.26		0.46
	industrial & urban		K				2.4	34.1		0.91		0.68
	industrial & urban		M				1.2	41.4		0.04		0.08
Husain et al., 1996, Kuwait	wholesalers	S	L							0.04		0.13
			K							0.30		0.15
Caggiano et al., 2005, Italy	urban		L							0.33		1.50
			K							6.71		2.00
			M							0.16		1.60
Abou-Arab, 2001*, Egypt	Industrial & urban		L				48.6	36.8		0.26		0.42
			K				3	34.2		0.82		0.54

C: chicken; G: goat; S: sheep; L: liver; K: kidney; M: muscle; G: gizzard.

* means mg/kg ww.

Table 3

Estimated Daily Intake (µg/kg bw/day) of As, Cd, Hg, Pb and Mn.

Sample name	As	Cd	Hg	Pb	Mn
Chicken Gizzard	0.19	0.21	0.03	0.96	1.61
Chicken Kidney	0.02	0.12	0.02	0.04	0.41
Chicken Liver	0.19	0.55	0.29	0.34	8.77
Chicken Muscle	0.12	0.003	0.03	0.03	0.37
Goat Kidney	0.002	0.01	0.005	0.01	0.13
Goat Liver	0.02	0.05	0.03	0.07	6.07
Goat Muscle	0.01	0.01	0.002	0.01	1.55
Sheep Kidney	0.002	0.03	0.01	0.01	0.19
Sheep Liver	0.04	0.13	0.07	0.14	8.92
Sheep Muscle	0.02	0.02	0.01	0.01	1.77
TDI by FAO/WHO/FSA	2.0	1.0	0.71	3.57	200

FAO: Food and Agricultural Organization.

WHO: World Health Organization.

FSA: Food Safety Agency.


Liver
Component 2 (16.5 %)
Component 1 (43.8 %)
Ni
Cu
Co
Cr
Mn
Zn
Pb
Fe
Cd
Hg
As
Kidney
Component 2 (15.9 %)
Component 1 (31.6 %)
Muscle
Component 2 (19 %)
Component 1 (41.7 %)

Fig. 1.


Child
(As, Cd, Hg & Pb) THQ
1.0
0.8
0.6
0.4
0.2
0.0
As
Cd
Hg
Pb
CG
CK
CL
CM
GK
GL
GM
SK
SL
SM
Adult
(As, Cd, Hg & Pb) THQ
0.8
0.6
0.4
0.2
0.0
As
Cd
Hg
Pb
CG
CK
CL
CM
GK
GL
GM
SK
SL
SM

Fig. 2.