



HOKKAIDO UNIVERSITY

Title	Identification of the phytosphingosine metabolic pathway leading to odd-numbered fatty acids
Author(s)	Kondo, Natsuki; Ohno, Yusuke; Yamagata, Maki et al.
Citation	Nature communications, 5, 5338 https://doi.org/10.1038/ncomms6338
Issue Date	2014-10
Doc URL	https://hdl.handle.net/2115/58535
Type	journal article
File Information	WoS_67865_Kihara.pdf



Identification of the phytosphingosine metabolic pathway leading to odd-numbered fatty acids

**Natsuki Kondo^{1,2}, Yusuke Ohno^{1,2}, Maki Yamagata¹, Takashi Obara¹, Naoya Seki¹,
Takuya Kitamura¹, Tatsuro Naganuma¹, and Akio Kihara¹**

¹Laboratory of Biochemistry, Faculty of Pharmaceutical Sciences, Hokkaido University, Kita 12-jo, Nishi 6-chome, Kita-ku, Sapporo 060-0812, Japan

²These authors contributed equally to this work.

Correspondence and requests for materials should be addressed to

Professor Akio Kihara
Laboratory of Biochemistry
Faculty of Pharmaceutical Sciences
Hokkaido University
Kita 12-jo, Nishi 6-chome
Kita-ku, Sapporo 060-0812, Japan
Tel: +81-11-706-3754
Fax: +81-11-706-4900
E-mail: kihara@pharm.hokudai.ac.jp

Running Head: Phytosphingosine metabolic pathway

ABSTRACT

The long-chain base phytosphingosine is a component of sphingolipids and exists in yeast, plants, and some mammalian tissues. Phytosphingosine is unique in that it possesses an additional hydroxyl group compared to other long-chain bases. However, its metabolism is unknown. Here we show that phytosphingosine is metabolized to odd-numbered fatty acids and is incorporated into glycerophospholipids both in yeast and mammalian cells. Disruption of the yeast gene encoding long-chain base 1-phosphate lyase, which catalyzes the committed step in the metabolism of phytosphingosine to glycerophospholipids, causes a ~40% reduction in the level of phosphatidylcholines that contain a C15 fatty acid. We also find that 2-hydroxypalmitic acid is an intermediate of the phytosphingosine metabolic pathway. Furthermore, we show that the yeast *MPO1* gene, whose product belongs to a large, conserved protein family of unknown function, is involved in phytosphingosine metabolism. Our findings provide insights into fatty acid diversity and identify a pathway by which hydroxyl group-containing lipids are metabolized.

INTRODUCTION

Cell membranes contain thousands of lipid species, which have compositions specific to each organism, tissue, cell, and organelle and become altered in response to environmental changes. Cell membranes with certain lipid compositions exhibit characteristic properties of fluidity, thickness, curvature, and local crowding (lipid microdomain), affecting protein activities and cellular signaling. Maintenance of cell- or tissue-specific lipid composition is important for cellular activity, and its deregulation leads to disorders. Fatty acids (FAs), the hydrophobic backbones of most lipid classes, are diverse molecules differing in chain-length, degree of saturation, and modification state, for instance, hydroxylation. Most cellular FAs contain even-numbered chain-lengths. However, odd-numbered FAs also exist, albeit at much lesser levels compared to even-numbered FAs¹⁻⁴.

Sphingolipids are one of the major lipid classes in eukaryotes. Their roles are not restricted to building blocks of plasma membrane but also include cell adhesion, recognition of viruses and bacterial toxins, skin barrier formation, neural functions, apoptosis, and glucose metabolism⁵⁻⁹. Furthermore, the sphingolipid metabolite sphingosine 1-phosphate (S1P) functions as a lipid mediator that regulates various cellular responses, such as egress of T cells from the thymus and secondary lymphoid tissues and vascular stabilization, by binding to one of the five S1P receptors (S1P₁₋₅)^{6,10}. Its function in the immune system has already been applied clinically as a therapeutic agent for multiple sclerosis¹¹.

The hydrophobic backbone of sphingolipids ceramide is composed of a long-chain base (LCB) and a FA connected by an amide bond⁶. The major LCB in mammals is a *trans*-double bond-containing sphingosine, which is a precursor of S1P. Sphingosine/LCB is phosphorylated by sphingosine/LCB kinases to produce S1P/LCB 1-phosphate (LCBP). The saturated LCB dihydrosphingosine (DHS) also exists throughout the body, although the levels of DHS are less abundant than those of sphingosine⁶. Phytosphingosine (PHS), which contains a hydroxyl

group at the C4 position, is localized to specific tissues such as the skin and small intestine^{7,12-14}. PHS is a major LCB in yeast, and it also exists in plants¹⁵.

In addition to the function of S1P/LCBP as a lipid mediator, S1P/LCBP also plays an important role as an intermediate in the sphingolipid-to-glycerophospholipid metabolic pathway¹⁶. This metabolic pathway is conserved in eukaryotes from yeast to humans, indicating its much older origin than the function of S1P/LCBP as a lipid mediator. We recently identified the complete metabolic pathway of S1P. S1P is metabolized to palmitoyl-CoA via four reactions (cleavage, oxidation, CoA addition, and saturation) and incorporated into glycerophospholipids¹⁷⁻¹⁹. DHS is also metabolized to glycerophospholipids after phosphorylation through a pathway that is essentially same as the S1P metabolic pathway, but without the saturation step¹⁷ (Fig. 1a). However, current knowledge about the metabolism of PHS is severely lacking. For example, it is even unknown whether PHS is metabolized to glycerophospholipids. If it is metabolized to glycerophospholipids, it still remains unclear whether PHS is incorporated into glycerophospholipids as FA or 2-hydroxy FA.

Several organisms, including bacteria, fungi, plants, and animals, possess 2-hydroxy FAs²⁰. In mammals, 2-hydroxy FAs exist in specific tissues such as the epidermis, small intestine, and in myelin^{7,21,22}. An additional hydroxyl group in 2-hydroxy FA as well as in PHS may confer the unique ability to strongly interact with other lipids and proteins via an extra hydrogen bond. In myelin, 2-hydroxy FAs are enriched in the glycosphingolipids galactosylceramide and sulfatide, and may play important functions in the firm stacking of multiple lipid layers. Mutations in the gene encoding FA 2-hydroxylase (*FA2H*) cause hereditary spastic paraplegia, a neuronal disorder²³. In the stratum corneum of the epidermis, multiple lipid lamellae act as a permeability barrier. Ceramides containing 2-hydroxy FAs and PHS are particularly abundant in the lipid lamellae⁷, and their hydroxyl groups may also strengthen the interaction between lipid layers. Although the hydroxyl group confers this

particular property to lipids, its presence renders their metabolism complex. Since 2-hydroxy FAs are only incorporated into sphingolipids and are resistant to β -oxidation, 2-hydroxy FAs must be converted to other metabolizable compounds for utilization in other lipid species or for removal. However, as with PHS metabolism, the metabolic pathway of 2-hydroxy FAs is largely unknown.

In the present study, we prepared radiolabeled PHS and first examined its metabolism using yeast as a model organism, since PHS is a major LCB in yeast²⁴. We then expanded the analyses to mammalian cells. We found that PHS was metabolized to sphingolipids and glycerophospholipids as other LCBs. Interestingly, PHS was metabolized to odd-numbered FAs in glycerophospholipids via the intermediate 2-hydroxypalmitic acid. Deletion of the yeast LCBP lyase gene (*DPL1*), which is responsible for the first step of LCBP metabolism, caused a reduction in the levels of the C15 phosphatidylcholine (PC) species to ~40% of the control, indicating that PHS is the major source of odd-numbered FAs in yeast. Furthermore, we identified that the yeast *MPO1* (metabolism of phytosphingosine to odd-numbered FA) gene is involved in PHS metabolism. Mpo1 belongs to the large protein family DUF962 (DUF: domain of unknown function), whose function presently is completely unknown. We speculate that Mpo1 acts as an α -oxidase of 2-hydroxyacyl-CoA. Through the pathway revealed here, PHS and 2-hydroxy FAs are converted to metabolizable, saturated FAs (odd-numbered FAs). Since deregulated accumulation of biomolecules often leads to cellular dysfunction or disorder, our newly identified PHS metabolic pathway may also be important for the maintenance of proper cellular functions.

Results

PHS is metabolized to odd-numbered FAs

The metabolism of PHS is currently unknown mainly due to the lack of commercially available radioactive PHS. Therefore, we prepared radiolabeled PHS ([11,12-³H]PHS) from [9,10-³H]palmitic acid by metabolic conversion in yeast, followed by purification. The prepared [³H]PHS presented as a single band that migrated to a position identical to a cold PHS standard on TLC (Fig. 1b), suggesting that our [³H]PHS was highly pure. Furthermore, all [³H]PHS was converted to PHS 1-phosphate (PHS1P) upon incubation with LCB kinase Lcb4 (Fig. 1c), confirming its identity and purity. We observed additional lipid products in addition to PHS1P following Lcb4 treatment (Fig. 1c, asterisk). Similar lipids were reported in a previous study²⁵, where the authors concluded that these lipids were generated during the TLC separation.

To examine the metabolism of PHS, wild-type yeast cells were labeled with [³H]PHS and [³H]DHS, for comparison, and their metabolites were separated by TLC (Fig. 1d). Consistent with a previous report¹⁷ (Fig. 1a), DHS was metabolized to sphingolipids (ceramide, inositol phosphorylceramide (IPC)), mannosylinositol phosphorylceramide (MIPC), and mannosyldiinositol phosphorylceramide (M(IP)₂C)) and glycerophospholipids (PC, phosphatidylethanolamine (PE), phosphatidylserine (PS), and phosphatidylinositol (PI)). Similar to DHS metabolism, PHS was metabolized to both sphingolipids and glycerophospholipids, but with the metabolic flow to sphingolipids being predominant (Fig. 1e).

To characterize the FA composition of these glycerophospholipids, [³H]DHS/[³H]PHS metabolites were subjected to mild saponification that hydrolyzes ester-linked FAs from glycerophospholipids without hydrolysis of amide-linked FAs from sphingolipids. The liberated FAs were esterified to FA methyl-esters (FAMES) and analyzed by reverse-phase

TLC (Fig. 2a) along with standard FAMES (Supplementary Fig. 1). A portion of the FAME mixture was also treated with $\text{KIO}_4\text{-KMnO}_4$ before TLC analysis to identify saturated and unsaturated FAs (Fig. 2b). The $\text{KMnO}_4\text{-KIO}_4$ treatment oxidatively cleaves a double bond, thus fragmenting unsaturated FAs; consequently, the spot corresponding to the unsaturated FA disappears on the TLC. DHS was metabolized to palmitoyl-CoA (C16:0) and palmitoleoyl-CoA (C16:1), the latter of which was formed by $\Delta 9$ desaturase-catalyzed reaction of the former, and subsequently to glycerophospholipids. On the other hand, PHS was incorporated into glycerophospholipids as odd-numbered FAs (C15:0 and, to a lesser extent, C15:1) (Fig. 2a,b).

The contribution of PHS-to-glycerophospholipid metabolism to cellular odd-numbered FA levels was examined by analyzing the most abundant glycerophospholipid, PC, in LCBP lyase-deficient yeast cells ($\Delta dpl1$) using LC-MS. In $\Delta dpl1$ cells lacking the PHS-to-glycerophospholipid metabolic pathway (Fig. 1a; see also Fig. 4a), most C15 PC species were significantly reduced (Fig. 2c). Deletion of the *DPL1* gene caused a reduction in total C15 PC levels to ~40% of wild-type levels (Fig. 2d). Some PC species with an elongated C17 FA were also decreased (Fig. 2c), while those with two even-numbered FAs were mostly unaffected (Fig. 2e). Similar results were obtained when the deletion mutant of the *SUR2* gene, which encodes the DHS C4-hydroxylase producing PHS from DHS, was analyzed (Supplementary Fig. 2).

When the pAN9 plasmid, which encoded the 3xFLAG-tagged *DPL1* gene under the control of the *GAPDH* promoter, was introduced into the $\Delta dpl1$ cells, C15 and C17 PC levels increased greatly, even when compared to the wild-type control (Supplementary Fig. 3). Furthermore, addition of PHS to the culture medium caused a large increase in odd-numbered PCs (Supplementary Fig. 4). Levels of some even-numbered PCs were also slightly altered by exogenous PHS. In summary, PHS is metabolized to odd-numbered FAs, and PHS appears to

be a major source of odd-numbered FAs in yeast.

LCBP lyase is a key determinant of LCB/LCBP metabolism

The metabolism of PHS to glycerophospholipids was less efficient than that of DHS (Fig. 1d). However, it is unclear what determines LCB/LCBP metabolism. After LCBs are converted to LCBPs by LCB kinases, LCBPs are degraded via one of two reactions: cleavage by LCBP lyase to fatty aldehydes and phosphoethanolamine and dephosphorylation by LCBP phosphatases to LCBs⁶ (Fig. 1a). Fatty aldehydes produced by LCBP lyase are metabolized to glycerophospholipids, whereas LCBs generated by LCBP phosphatases are recycled back to sphingolipids⁶.

The yeast LCB kinases are reported to phosphorylate DHS and PHS with similar efficiencies²⁵; however, little is known about the relative activity of LCBP lyase/phosphatase toward DHS 1-phosphate (DHS1P) and PHS1P. *In vitro* LCBP lyase/phosphatase assays were therefore performed on [³²P]DHS1P and [³²P]PHS1P using membrane fractions, and the product [³²P]phosphoethanolamine/[³²P]phosphate was monitored by normal-phase TLC (Fig. 3a). Incubation of wild-type membranes with DHS1P produced almost equal amounts of phosphoethanolamine and phosphate (a 0.92:1 ratio), whereas phosphate was preferentially generated from PHS1P (a 0.45:1 ratio) (lines 3 and 4; see Fig. 3b for quantified data). Deletion of the sole LCBP lyase *DPL1*, ($\Delta dpl1$ cell membranes) did not produce phosphoethanolamine (lanes 5 and 6); however, deletion of the LCBP phosphatase *Lcb3*, the more predominant one of the two (*Ysr3* being the other)²⁶, was not sufficient to prevent the production of phosphate but reduced it with a concomitant increase of phosphoethanolamine (lanes 7 and 8). The lipid phosphate phosphatases *Lpp1* and *Dpp1* in yeast are known to exhibit broad specificity toward lipid phosphates²⁷. Accordingly, DHS1P/PHS1P dephosphorylation was eliminated by using membranes of the $\Delta lcb3 \Delta ysr3 \Delta lpp1 \Delta dpp1$ quadruple deletion mutant (lanes 9 and 10). This

result is consistent with our previous data indicating that membranes of the $\Delta lcb3 \Delta lpp1 \Delta dpp1$ triple deletion mutant had no phosphatase activity toward DHS1P²⁸. Even under these phosphatase-null conditions, conversion of PHS1P to phosphoethanolamine was low (lane 10).

The above results suggest that the differences in the metabolism of DHS and PHS (Fig. 1d) may be caused by different substrate preferences of the LCBP lyase Dpl1, but not of the LCBP phosphatases, toward DHS1P and PHS1P. To test this possibility, we next determined the kinetic parameters of Dpl1 and Lcb3 using cell membranes prepared from yeast cells lacking LCBP lyase and LCBP phosphatases but expressing 3xFLAG-tagged Dpl1 or Lcb3. Immunoblotting revealed that expression levels of 3xFLAG-Dpl1 and 3xFLAG-Lcb3 were equivalent (Fig. 3c). We determined the $V_{\max}$ and K_m values for Dpl1 to be 28.3 nmol/ μ g/min and 19.6 μ M for DHS1P and 8.9 nmol/ μ g/min and 10.6 μ M for PHS1P, respectively (Fig. 3d). Thus, the $V_{\max}$ value of Dpl1 toward PHS1P is $\sim 1/3$ of that toward DHS1P, indicating that the activity of Dpl1 toward PHS1P was indeed low. On the other hand, the kinetic parameters of Lcb3 toward DHS1P and PHS1P were similar (DHS1P, 6.0 nmol/ μ g/min for $V_{\max}$ and 8.3 μ M for K_m ; PHS1P, 7.3 nmol/ μ g/min for $V_{\max}$ and 17.4 μ M for K_m) (Fig. 3e). These results indicate that the activity of Dpl1 determines the efficiency of LCBP metabolism to glycerophospholipids.

2-hydroxypalmitic acid is an intermediate in PHS metabolism

We previously reported that DHS was metabolized to glycerophospholipids via DHS1P, hexadecanal, palmitic acid, and palmitoyl-CoA¹⁷ (Fig. 1a). The LCB kinase Lcb4, LCBP lyase Dpl1, fatty aldehyde dehydrogenase Hfd1, and acyl-CoA synthetases Faa1 and Faa4 are involved in each conversion step¹⁷ (Fig. 1a). We next examined whether these enzymes were also involved in PHS metabolism. The PHS-to-glycerophospholipid conversion was completely abolished in $\Delta lcb4$, $\Delta dpl1$, $\Delta hfd1$, and $\Delta faa1 \Delta faa4$ cells (Fig. 4a). Thus, the

reactions catalyzed by the LCB kinase Lcb4, LCBP lyase Dpl1, fatty aldehyde dehydrogenase Hfd1, and acyl-CoA synthetases Faa1 and Faa4 are commonly required for DHS and PHS metabolism. Our results indicate that PHS is converted to PHS1P by Lcb4, to 2-hydroxyhexadecanal by Dpl1, and to 2-hydroxypalmitic acid by Hfd1. Consistent with the conclusion that 2-hydroxypalmitic acid is an intermediate of the PHS metabolic pathway, addition of 2-hydroxypalmitic acid to the culture medium caused an increase in odd-numbered PCs, as did addition of PHS (Supplementary Fig. 4).

To determine the reaction catalyzed by the acyl-CoA synthetases Faa1 and Faa4, we examined the FA species that accumulated in $\Delta faa1 \Delta faa4$ cells using normal-phase TLC and hexane/diethyl ether/acetic acid (30:70:1) as a resolving buffer. Although only small amounts of FAs were detected in the cell fraction of [^3H]PHS labeled $\Delta faa1 \Delta faa4$ cells, FA and 2-hydroxy FA were observed in the medium (Fig. 4b). Methyl esterification and separation by reverse-phase TLC revealed that they were methyl esters of C15:0 FA (pentadecanoic acid) and 2-hydroxypalmitic acid (Fig. 4c). These FA intermediates were not observed in wild-type cells (Fig. 4b). When $\Delta faa1 \Delta faa4$ cells were labeled with [^3H]DHS, palmitic acid was found to accumulate in the medium (Fig. 4b,c), as expected from the metabolic pathway of DHS (Fig. 1a).

***MPO1* is involved in the metabolism of PHS**

The results described above indicate that PHS is metabolized to 2-hydroxypalmitic acid and then converted to pentadecanoic acid via α -oxidation. To identify a gene involved in α -oxidation of PHS metabolites, we performed bioinformatic analysis using a yeast database. Since LCBP metabolism takes place in the ER¹⁶, we speculated that the corresponding gene encodes an ER protein. Therefore, we searched for genes encoding ER proteins in the yeast GFP fusion localization database (<http://yeastgfp.yeastgenome.org/>)²⁹ and found 296

genes/open reading frames (ORFs). Of these, the 13 genes with unknown functions were selected and their deletion mutants were subjected to the [^3H]PHS-labeling experiment. We found that one of these genes, termed *MPO1* (metabolism of phytosphingosine to odd-numbered FA, ORF *YGL010w*), exhibited a defect in the metabolism of PHS to glycerophospholipids when the gene was deleted (Supplementary Fig. 5). Deletion of the *MPO1* gene had no effect on DHS metabolism (Fig. 5a). These results indicate that the *MPO1* gene is involved in a PHS-specific reaction, α -oxidation of PHS metabolites. Moreover, $\Delta mpo1$ cells with [^3H]PHS accumulated 2-hydroxypalmitic acid in the medium (Fig. 5b). These results suggest that 2-hydroxypalmitic acid or its reversible metabolite is the substrate of Mpo1 and is released into the medium without accumulating in cells.

Mpo1 belongs to the DUF962 (DUF: domain of unknown function) superfamily (<http://www.ebi.ac.uk/interpro/entry/IPR009305>). To date, over 3,000 proteins from bacteria, fungi, plants, and protists have been registered as DUF962 family members. However, no functions have been reported for any of these members. Mpo1 is a relatively small protein, and its predicted molecular mass is 20.2 kDa. It is also a highly hydrophobic protein and may possess 2 to 4 transmembrane segments (Fig. 5c,d). Mpo1 contains no apparent domain or sequence motif found in other protein families. However, the N-terminal region of Mpo1 (Fig. 5d, underlined) shares weak sequence similarity with the NAD^+ -binding domain of the histone deacetylase Sir2, implying that Mpo1 may function as an oxidase with NAD^+ acting as the oxidant. We confirmed the ER localization of Mpo1 using N-terminally GFP-tagged Mpo1 protein. GFP-Mpo1 exhibited a typical ER pattern, i.e., two-ring structures of nuclear ER that surround the nucleus, and cortical ERs (Fig. 5e). GFP-Mpo1 indeed co-localized with the mCherry-tagged ER marker Sec61.

PHS is metabolized to odd-numbered FAs in mammalian cells

We next examined the metabolism of PHS in mammalian cells using CHO-K1 cells via a [^3H]PHS labeling experiment. Comparing the metabolism of PHS with that of DHS, sphingomyelin slightly decreased, while monohexosylceramide increased (Fig. 6a; see Supplementary Fig. 6 for the sphingolipid biosynthetic pathway in mammals). Glycerophospholipid levels appeared to be equivalent between [^3H]DHS and [^3H]PHS labeling experiments, in contrast to the case of yeast. To examine the FA species derived from PHS in mammals, PHS metabolites were analyzed by reverse-phase TLC after methyl-esterification. We observed production of C15:0 FAME and, to a lesser extent, its elongated product C17:0 FAME, both of which were resistant to $\text{KIO}_4/\text{KMnO}_4$ treatment (Fig. 6b). By [^3H]DHS-labeling, C16:0 FAME and its elongated product C18:0 FAME were generated (Fig. 6b). Similar results were obtained when HEK 293T cells were subjected to the analyses (Fig. 6c,d). Thus, although PHS was metabolized to odd-numbered FAs in mammalian cells as well as in yeast, the odd-numbered FAs generated in yeast were partly desaturated, while mammals produced elongated odd-numbered FAs.

In yeast, PHS was metabolized to glycerophospholipids after conversion to PHS1P and cleavage by LCBP/S1P lyase. To investigate whether PHS is also metabolized via a similar pathway in mammals, we performed the [^3H]DHS and [^3H]PHS labeling experiments with mouse embryonic carcinoma F9 cells and its mutant cell line (S1P lyase *Spl* gene (the *DPL1* homolog)-deficient) (Fig. 6e). Consistent with our previous results¹⁷, the metabolism of DHS to glycerophospholipids was almost completely abolished in F9 *Spl*^{-/-} cells. Similarly, the metabolism of PHS to glycerophospholipids was also impaired in F9 *Spl*^{-/-} cells. Thus, the metabolism of PHS to glycerophospholipids is dependent on S1P lyase in mammalian cells as well as in yeast.

Discussion

In the present study, we demonstrate for the first time that PHS is metabolized to glycerophospholipids in both yeast and mammalian cells (Fig. 1d and Fig. 6a,c,e), as are other LCBs including sphingosine and DHS¹⁷. Although sphingosine and DHS were metabolized to palmitic acid (C16:0) and its desaturated (C16:1) or elongated (C18:0) derivative in glycerophospholipids, PHS was converted to odd-numbered pentadecanoic acid (C15:0) and its derivatives (C15:1, C17:0, and 17:1) (Fig. 2 and Fig. 6). The existence of odd-numbered FAs in mammals has been known since the 1950's^{30,31}. However, their origin remains unclear, although dietary intake and *de novo* FA synthesis by FA synthase using propionyl-CoA have been considered to be sources of odd-numbered FAs^{3,32,33}. FA synthase normally utilizes acetyl-acyl carrier protein (ACP) as a starting material for condensation with malonyl-ACP, producing even-numbered FAs, mainly palmitic acid. However, a small fraction of propionyl-CoA produced during amino acid and cholesterol metabolism appears to be converted to propionyl-ACP, which is then used for FA synthesis. When mice were fed tripropionin, levels of odd-numbered FAs were increased in depot fat³². In addition, propionic acidemia patients, who cannot metabolize propionyl-CoA due to a defect in propionyl-CoA decarboxylase, were found to have ~4–5-fold higher levels of odd-numbered FAs in their erythrocytes, plasma, liver, and subcutaneous tissue compared to controls³. However, the contribution of the propionate-derived odd-numbered FAs relative to total odd-numbered FAs under normal conditions and other tissues remains unknown. Here, we report that PHS was the major source of odd-numbered FAs, at least in yeast (Fig. 2c,d and Supplementary Fig. 2). In mammals, PHS is localized to specific tissues such as the skin and small intestine^{7,12-14}. Consistent with the abundance of PHS in skin, odd-numbered FAs also exist in skin at extremely high levels⁴, suggesting that PHS may be the major source of odd-numbered FAs in PHS-rich mammalian tissues.

We observed that the metabolism of PHS to glycerophospholipids depended on the LCB kinase Lcb4, LCBP lyase Dpl1, fatty aldehyde dehydrogenase Hfd1, and acyl-CoA synthetases Faa1 and Faa4 (Fig. 4a). Therefore, PHS is first converted to PHS1P by Lcb4 and degraded to 2-hydroxyhexadecanal by Dpl1 (Fig. 7). This degradation step is the key determinant of the metabolic flow of LCB/LCBP (Fig. 3). 2-hydroxyhexadecanal is then oxidized to 2-hydroxypalmitic acid by Hfd1 (Fig. 7). We detected 2-hydroxypalmitic acid and pentadecanoic acid in the medium of $\Delta faa1 \Delta faa4$ cells (Fig. 4b), suggesting that both 2-hydroxypalmitic acid and pentadecanoic acid are substrates of the acyl-CoA synthetases Faa1 and Faa4 in the PHS metabolic pathway. Thus, the Hfd1 product 2-hydroxypalmitic acid may be converted to 2-hydroxypalmitoyl-CoA by Faa1 and Faa4 in the next step (Fig. 7). Although the exact role of Mpo1 is unclear at present, Mpo1 must be involved in a PHS-specific reaction, i.e., a reaction that is not involved in DHS metabolism (Fig. 5a). Given that Mpo1 is a putative NAD⁺-binding protein, we hypothesize that Mpo1 may be involved in the α -oxidation of 2-hydroxypalmitoyl-CoA, producing pentadecanoic acid or other metabolites such as pentadecanal that can be metabolized to pentadecanoic acid. Pentadecanoic acid is finally converted to pentadecanoyl-CoA by Faa1 and Faa4 for glycerophospholipid synthesis (Fig. 7). The observation that pentadecanoic acid was detected in the medium of $\Delta faa1 \Delta faa4$ cells (Fig. 4b) suggests that metabolism of at least some 2-hydroxypalmitic acid can still proceed to pentadecanoic acid even in the absence of Faa1 and Faa4. Yeast contain 7 acyl-CoA synthetases (Faa1, Faa2, Faa3, Faa4, Fat1, Fat2, and Acs1). Therefore, acyl-CoA synthetases other than Faa1 and Faa4 may have residual activity toward 2-hydroxypalmitic acid in $\Delta faa1 \Delta faa4$ cells. As 2-hydroxypalmitic acid was detected in the medium of $\Delta mpo1$ cells (Fig. 5b), we speculate that 2-hydroxypalmitoyl-CoA is converted to 2-hydroxypalmitic acid by an acyl-CoA hydrolase and is released into the medium in the $\Delta mpo1$ cells.

Although peroxisomes are known to be involved in α -oxidation³⁴, we found that Mpo1 was localized to the ER (Fig. 5e), similar to other enzymes involved in LCBP metabolism¹⁶. We examined the involvement of peroxisomes in PHS metabolism by using yeast deletion mutants for the peroxisomal translocation machinery *PEX2* and *PEX12* and the ABC transporter subunits *PXA1* and *PXA2* that are required for importing long-chain FA into peroxisomes. However, PHS was metabolized as normal in these mutants (Supplementary Fig. 7). Thus, our results suggest that peroxisomes are unlikely to be involved in PHS metabolism; instead, all reactions in the PHS metabolic pathway are likely to occur in the ER.

Mpo1 belongs to the large protein family DUF962 (DUF: domain of unknown function), whose functions are unknown at present. Our findings suggest that Mpo1 is involved in the α -oxidation of 2-hydroxyacyl-CoAs. As the function of the DUF962 family has been elucidated here, we will refer to it as the Mpo1 family hereafter. Mpo1 family members exist in bacteria, fungi, plants, and protists; 2-hydroxy FAs also exist in these organisms²⁰, although its presence in protists has not been examined. It is therefore highly likely that Mpo1 family members are involved in the α -oxidation of 2-hydroxyacyl-CoAs in general.

Our results indicate that PHS is metabolized to odd-numbered FAs in mammals (Fig. 6). However, Mpo1 family members are not expressed in mammals, in contrast to other LCB/LCBP metabolic enzymes conserved between yeast and mammals¹⁶. PHS seems to have once disappeared in the course of evolution, when Animalia diverged from Plantae. Therefore, members of lower Animalia do not possess PHS. However, mammals again evolutionarily acquired the ability to produce PHS, likely due to the requirement for PHS to fulfill the functions in specific tissues. We speculate that Mpo1 homologs were lost at the same time as when PHS disappeared in Animalia. Instead, mammals may have independently evolved an α -oxidase from other protein families, when they began to utilize PHS.

LCBs have two hydroxyl groups and one amino group in common, both of which can

form hydrogen bonds. Since PHS contains an additional hydroxyl group, PHS may possess different properties compared to other LCBs and specific functions by interacting with other lipids or proteins with higher affinity due to additional hydrogen bond formation. Therefore, PHS may be important for PHS-containing tissues such as the skin and small intestine, and its reduction may lead to disorders. Indeed, a decrease in PHS-containing ceramide levels has been observed in psoriatic scale³⁵. In addition, the degradation pathway must also be important for the maintenance of appropriate levels of PHS, as it is for other biomolecules in general. Indeed, several metabolic disorders, including sphingolipidoses³⁶, are known to be due to defects in certain lipid degradation pathway. The PHS metabolite, odd-numbered FAs, is much less common than even-numbered FAs, and both are similarly metabolized to glycerophospholipids. Thus, odd-numbered FA itself might not have physiological significance. However, the PHS metabolic pathway that we have elucidated here may be important for the maintenance of cellular PHS levels and clearance of PHS metabolites such as 2-hydroxy FAs, and disorders can arise from its deregulation. Both PHS and 2-hydroxypalmitic acid are incorporated into sphingolipids only, and are not used in other lipids. In addition, they cannot be subjected to β -oxidation in their current forms. The PHS metabolic pathway revealed here converts them to the metabolizable compound, non-hydroxylated pentadecanoic acid. Pentadecanoic acid can undergo desaturation, elongation, incorporation into glycerophospholipids, and mitochondrial β -oxidation, where the final round product propionyl-CoA is converted to succinyl-CoA and subsequently metabolized in the citric acid cycle³⁷.

In mammals, there remain a large number of orphan G protein-coupled receptors (GPCRs). Recent developments in lipidomics analysis have led to the discovery of many novel lipid metabolites, some of which have been determined to be ligands of orphan GPCRs. Our elucidation of the PHS metabolic pathway encourages the search for novel bioactive lipid

molecules in the derivatives of PHS metabolites.

Methods

Yeast strains and media

The *Saccharomyces cerevisiae* strains used in this study are listed in Supplementary Table 1. BY4741 cells³⁸ were obtained from Open Biosystems (Huntsville, AL). Chromosome fusion of *mCherry* to the 3'-terminus of the *SEC61* gene was performed using PCR-based gene disruption and modification³⁹. Cells were grown in YPD medium (1% yeast extract, 2% peptone, and 2% D-glucose), synthetic complete (SC) medium (0.67% yeast nitrogen base, 2% D-glucose, 20 mg/l adenine, 20 mg/l uracil, and an amino acid mixture), or SC medium lacking uracil (SC-URA; 0.67% yeast nitrogen base, 2% D-glucose, 20 mg/l adenine, 20 mg/l tryptophan, and 0.5% casamino acids).

Cell culture

Human embryonic kidney (HEK) 293T (American Type Culture Collection (ATCC), Manassas, VA; Cat. No. CRL-3216) and mouse embryonic carcinoma and F9 cells (ATCC; CRL-1720) were grown in Dulbecco's modified Eagle's medium (Sigma/Sigma-Aldrich, St. Louis, MO), while Chinese hamster ovary CHO-K1 cells (ATCC; CRL-9618) were cultured in Ham's F-12 medium (Sigma). Each medium contained 10% fetal bovine serum (FBS), 100 U/ml penicillin, and 100 µg/ml streptomycin. HEK 293T and F9 cells were grown in dishes coated with 0.3% collagen and 0.1% gelatin, respectively. *Spl*-deficient F9 cells (F9-2) have been described previously⁴⁰.

Plasmids

The pAKNF316 plasmid (*URA3* marker) is a yeast expression vector and was designed to produce an N-terminal 3xFLAG-tagged protein under the control of the *GAPDH* (*TDH3*) promoter. The pAN9 (3xFLAG-*DPL1*) and pAN13 (3xFLAG-*LCB3*) plasmids were

constructed by cloning of *DPL1* and *LCB3* genes into the pAKNF316 plasmid, respectively. The pTS229 plasmid (*TRP1* marker) encodes 3xFLAG-*LCB4* under the control of the *GAPDH* promoter.

The pUG36 plasmid (*URA3* marker), a yeast expression vector constructed to produce a fusion protein with an N-terminal enhanced GFP under the control of the *MET15* promoter was a gift from J. H. Hegemann. The pNK47 (*GFP-MPO1*) plasmid was constructed by cloning the *MPO1* gene into the pUG36 plasmid.

Preparation of [11, 12-³H]PHS

BY4741 cells (100 ml) grown in YPD medium to log phase were labeled at 30 °C for 4 hr with 1 mCi [9,10-³H]palmitic acid (40 Ci/mmol, 10 mCi/ml; American Radiolabeled Chemical, St. Louis, MO). Lipids were extracted and hydrolyzed by HCl/methanol as described elsewhere⁴¹. Lipids suspended in 300 µl chloroform /methanol (2:1, v/v) were resolved by normal-phase TLC on Silica Gel 60 high performance TLC plates (Merck, Whitehouse Station, NJ) with chloroform/methanol/4.2 N ammonia (9:7:2, v/v) so the leading edge was positioned 2.5 cm from the top. Plates were then dried and resolved with 1-butanol/acetic acid/water (3:1:1, v/v) to the top. PHS was scraped from the TLC plate and was extracted three times using 200 µl ethanol. Silica gel was removed by centrifugation, and the collected lipids were combined. The obtained PHS was estimated to be 0.025 µCi/µl by measuring its radioactivity using a liquid scintillation counter (LSC-3600; Aloka, Tokyo, Japan) and to be 0.55 mM by comparing its ninhydrin-stained density with cold PHS standard (Enzo Life Sciences, Farmingdale, NY).

To phosphorylate [11, 12-³H]PHS, 8 nCi [11, 12-³H]PHS was incubated with 250 ng 3xFLAG-Lcb4 for 1 hr at 37 °C in 50 µl buffer A (50 mM HEPES-NaOH (pH 7.5), 150 mM NaCl, 200 mM sorbitol, 1x Complete protease inhibitor mixture (EDTA-free; Roche Diagnostics, Indianapolis, IN), 1 mM phenylmethylsulfonyl fluoride, 1 mM dithiothreitol, 5

mM ATP, and 10 mM MgCl₂) containing 0.5 mg/ml FA-free BSA (A6003; Sigma). Soluble, non-palmitoylated 3xFLAG-Lcb4 was purified as described previously⁴², by subjecting the soluble fraction prepared from the palmitoyltransferase mutant KHY543 ($\Delta akr1$) cells harboring the pTS229 (3xFLAG-LCB4) plasmid to incubation with anti-FLAG M2 agarose (Sigma), followed by extensive washing and elution with buffer containing 100 μ g/ml 3xFLAG peptide. Lipids were extracted by successive addition and mixing of 187.5 μ l chloroform/methanol/HCl (100:200:1, v/v), 62.5 μ l chloroform, and 62.5 μ l 1% KCl. Phases were separated by centrifugation, and the organic phase was recovered, dried, and suspended in chloroform /methanol (2:1, v/v). The labeled lipids were resolved by normal-phase TLC on Silica Gel 60 high performance TLC plates with chloroform/methanol/4.2 N ammonia (9:7:2, v/v).

[³H]LCB-labeling assay

To prepare [4,5-³H]DHS with its radioactivity and concentration identical to [11,12-³H]PHS (0.025 μ Ci/ μ l, 0.55 mM), [4,5-³H]DHS (50 Ci/mmol; American Radiolabeled Chemicals) was mixed with cold DHS (Enzo Life Sciences). The [³H]LCB-labeling assay using mammalian cells was performed as described previously^{17,19}. The [³H]LCB-labeling assay using yeast was conducted as follows. Yeast cells were grown to $\sim 5 \times 10^6$ cells/ml in YPD medium and then 0.75 ml of cells were labeled with [11,12-³H]PHS or [4,5-³H]DHS (0.1 μ Ci/2.2 nmol or 0.2 μ Ci/0.44 nmol as indicated in figure legends) at 30 °C for 3 hr. After labeling, the cells and medium were separated by centrifugation (2,000 *g*, room temperature, 2 min). Cells were washed with 0.5 ml water and resuspended in 200 μ l ethanol/water/diethyl ether/pyridine/15 N ammonia (15:15:5:1:0.018, v/v). After a 15-min incubation at 60 °C with mixing every 5 min, extracted lipids were recovered in the supernatant after centrifugation (8,000 *g*, room temperature, 1 min). Radioactivity was measured using the liquid scintillation system

LSC-3600, and lipid extracts of equal radioactivity were dried. Lipids from the medium were extracted by mixing with successive additions of 3.75 volumes of chloroform/methanol/HCl (100:200:1, v/v), 1.25 volumes of chloroform, and 1.25 volumes of 1% KCl. Phases were then separated by centrifugation (8,000 *g*, room temperature, 1 min), and the organic phase was recovered and dried. Lipids were resuspended in 20 μ l chloroform/methanol/water (5:4:1, v/v) and separated by normal-phase TLC on Silica Gel 60 high performance TLC plates with chloroform/methanol/4.2 N ammonia (9:7:2, v/v) or hexane/diethyl ether/acetic acid (30:70:1, v/v) as the solvent. Note that other than FAs, no labeled lipids were moved from the origin by the latter, highly nonpolar, resolving buffer. Lipids were visualized by spraying the TLC plates with a fluorographic reagent (2.8 mg/ml 2,5-diphenyl-oxazole in 2-methylnaphthalene/1-butanol (1:3.3, v/v)) and exposing them to X-ray film at -80 °C. The identities of the lipid bands on TLCs have been described previously¹⁷.

Methyl-esterification of FAs, double-bond cleavage assays using KMnO₄ and KIO₄, and separation of FAMES by reverse-phase TLC were performed as follows. Lipids extracted from labeled cells with ethanol/water/diethyl ether/pyridine/15 N ammonia (15:15:5:1:0.018, v/v) as described above were treated with 0.25 volumes of 0.5 M NaOH in methanol and incubated at 37 °C for 30 min to liberate the FAs from glycerophospholipids. After neutralization with acetic acid, lipids were extracted by mixing with successive additions of 2.25 volumes of chloroform/methanol (7:11, v/v), 0.94 volumes of chloroform, and 0.94 volumes of 1% KCl. Phases were then separated by centrifugation (8,000 *g*, room temperature, 1 min), and the organic phase was recovered and dried. Lipids were resuspended in 50 μ l chloroform/methanol (2:1, v/v) and separated by normal-phase TLC on Silica Gel 60 high performance TLC plates with chloroform/methanol/4.2 N ammonia (9:7:2, v/v). FAs were eluted from the silica with ethanol and dried. The prepared FAs or lipids extracted from the medium as described above were methyl-esterified by resuspending the dried lipids in 200 μ l 1 M methanolic HCl and 5%

2,2-dimethoxypropane, followed by incubation at 100 °C for 1 hr. Generated FAMEs were extracted from the methanol phase with 300 µl hexane twice, dried, and separated by reverse-phase TLC or subjected to double-bond cleavage assays. For the double-bond cleavage assay, lipids were resuspended in 100 µl 90% acetic acid and divided into two portions of 50 µl each. Lipids were treated with 75 µl 0.5% KMnO₄ and 0.5% KIO₄ (or 75 µl water for the control) at 37 °C for 1 hr, reduced with 15 µl 20% sodium bisulfite (or 15 µl water for the control), and neutralized with 50 µl 15 M ammonia. Lipids were then extracted by adding 250 µl chloroform. After phase separation by centrifugation (8,000 *g*, room temperature, 1 min), the organic phase was recovered and dried. Lipids were resuspended in 20 µl chloroform and separated by reverse-phase TLC on a Silica gel 60 RP-18 F₂₅₄s TLC plates (Merck) with chloroform/methanol/water (5:15:1, v/v).

Preparation of ³²P-labeled LCBPs

Cold DHS and PHS were purchased from Enzo Life Sciences and dissolved in ethanol. The ³²P-labeled LCBPs were prepared by incubating 30 µM LCBs complexed with 0.4 mg/ml FA-free BSA with 13 µCi [γ-³²P]ATP (3,000 Ci/mmol; PerkinElmer Life Sciences, Waltham, MA) and 62.5 ng purified 3xFLAG-Lcb4⁴² at 37 °C for 1 hr in 100 µl buffer B (50 mM HEPES-NaOH (pH 7.5), 150 mM NaCl, 10% glycerol, 1x Complete protease inhibitor mixture, 1 mM phenylmethylsulfonyl fluoride, and 1 mM dithiothreitol) containing 5 mM MgCl₂. Produced LCBPs were extracted by successive additions of 600 µl chloroform/methanol (1:2, v/v), 7.5 µl 15 M ammonia, 400 µl chloroform, and 400 µl 1% KCl, with mixing. Phases were then separated by centrifugation, and the upper, water phase was recovered and mixed with 600 µl chloroform and 77 µl of 5 M HCl. Phases were separated by centrifugation again, and the lower, organic phase was recovered, dried, and dissolved in ethanol.

Cold DHS1P (Sigma) and PHS1P (Avanti Polar Lipids, Alabaster, AL) were first

dissolved in methanol at 0.5 mM by incubating at 75 °C, followed by sonication. LCBPs were then diluted to 50 µM in buffer B containing 2 mg/ml FA-free BSA and stored at -30°C. Each ³²P-labeled LCBP prepared as above was mixed with cold LCBP to the desired concentration prior to use.

Preparation of total membrane fractions

Approximately 4×10^8 yeast cells were suspended in 3.3 ml buffer C (100 mM Tris-HCl (pH 9.4) and 40 mM 2-mercaptoethanol), and incubated at room temperature for 10 min. After centrifugation, cells were resuspended in 6.6 ml buffer D (50 mM Tris-HCl (pH 7.5) and 1.2 M sorbitol), and incubated at 30 °C for 30 min with 0.33 mg Zymolyase 100T (Seikagaku Corporation, Tokyo, Japan). The resulting spheroplasts were washed with 3.3 ml buffer D, suspended in 0.5 ml buffer B, and lysed by sonication. Cell debris was removed by centrifugation (1,000 g, 4 °C, 3 min), and the supernatant was subjected to centrifugation at 100,000 g at 4 °C for 30 min. The pellet (total membrane fractions) was suspended in 750 µl buffer B by sonication.

***In vitro* LCBP lyase/phosphatase assay**

The *in vitro* LCBP lyase/phosphatase assay was performed by incubation of total membrane fractions with LCBP (25 nCi ³²P-labeled LCBP plus desired concentration of cold LCBP), complexed with FA-free BSA, at 37 °C for 10 min in a 50 µl reaction mixture (50 mM HEPES-NaOH (pH 7.5), 150 mM NaCl, 10% glycerol, 1x Complete protease inhibitor mixture, 1 mM phenylmethylsulfonyl fluoride, 1 mM dithiothreitol, and 250 µM pyridoxal 5'-phosphate). After the reaction, the 7 µl reaction mixture was mixed with 33 µl chloroform/methanol/water/HCl (100:200:30:1, v/v), and 5 µl of the resultant mixture was spotted on Silica Gel 60 high-performance TLC plates and separated with 1-butanol/acetic

acid/water (2:1:1, v/v) as the solvent system. Labeled products were detected and quantified by a bioimaging analyzer BAS-2500 (Fuji Photo Film, Tokyo, Japan).

Lipid analysis by mass spectrometry

The FA composition of PC was determined by ultra-performance liquid chromatography (UPLC) coupled with electrospray ionization tandem quadrupole time-of-flight mass spectroscopy (Xevo G2 QTof; Waters, Milford, MA) as follows. Yeast cells were grown in SC medium, and lipids were extracted from $\sim 1.2 \times 10^8$ cells by incubating cell pellets with ethanol/water/diethyl ether/pyridine/15 N ammonia (15:15:5:1:0.018, v/v) at 60 °C for 15 min in the presence of 1 nmol C12:0 PC (Avanti Polar Lipids) as an internal standard. Extracted lipids were then separated by normal-phase TLC with chloroform/methanol/4.2 N ammonia (9:7:2, v/v), and silica containing PC was scraped from the TLC plate. PC was eluted from silica with chloroform/methanol (1:2, v/v), dried, suspended in methanol, resolved by UPLC on a reverse-phase column (AQUITY UPLC BEH C18 column, length 150 mm; Waters) at 45 °C, and detected by MS. The flow rate was 0.1 ml/min in the binary gradient system using mobile phase A (acetonitrile/water/methanol (4:4:2, v/v) containing 0.1% formic acid and 0.025 % ammonia) and mobile phase B (2-propanol/methanol (4:1, v/v) containing 0.1% formic acid and 0.025 % ammonia). The elution gradient steps were as follows: 0–10 min, 5% B; 10–50 min, gradient to 80% B; 50–55 min, gradient to 100% B; 55–65 min, 100% B; 65–66 min, gradient to 5% B; 66–80 min, 5% B. The ESI capillary voltage was set at 2.6 kV, sampling cone was at 35 V, and extraction cone was at 4 V in negative ion mode. The mass range was set at m/z 500-900, and scan duration was set at 1 sec.

To discriminate PC species having the same molecular mass but different FA compositions, the same samples used in the above MS analysis were also subjected to MS analysis in MS/MS mode. The m/z ($[M+HCOO]^-$) of specific PC species, as listed in

Supplementary Table 2, was selected at Q1 (Supplementary Fig. 8), and their FA compositions were determined by analyzing their fragment ions (Supplementary Table 3 and Supplementary Fig. 9). Argon gas was used as the collision gas, and collision energy was set at 30 V. Data analysis and quantification were performed using MassLynx software (Waters). For quantification, levels of PC species with the same molecular mass were first determined by comparing their ion intensities with those of the internal standards in the first MS analyses, and levels of each PC species having specific FA composition were then calculated by comparing the ion intensities of their fragments in the MS/MS analysis.

Immunoblotting

Immunoblotting was performed as described previously⁴³. Proteins were electrotransferred to an Immobilon PVDF membrane (Millipore, Billerica, MA), which was incubated with primary anti-FLAG M2 antibody (1 µg/ml; Stratagene, Agilent Technologies) for 1 hr and then with secondary HRP-conjugated anti-mouse IgG F(ab')₂ fragment antibody (1:7500 dilution; GE Healthcare Life Sciences, Buckinghamshire, UK) for an additional 1 hr. Labeling was detected using Pierce ECL Western Blotting Substrate (Thermo Fisher Scientific, Waltham, MA).

References

- 1 Hajra, A. K. & Radin, N. S. Isotopic studies of the biosynthesis of the cerebroside fatty acids in rats. *J. Lipid Res.* **4**, 270-278 (1963).
- 2 Svennerholm, L. & Stallberg-Stenhagen, S. Changes in the fatty acid composition of cerebroside and sulfatides of human nervous tissue with age. *J. Lipid Res.* **9**, 215-225 (1968).
- 3 Wendel, U., Baumgartner, R., van der Meer, S. B. & Spaapen, L. J. Accumulation of odd-numbered long-chain fatty acids in fetuses and neonates with inherited disorders of propionate metabolism. *Pediatr. Res.* **29**, 403-405 (1991).
- 4 Hinder, A., Schmelzer, C. E., Rawlings, A. V. & Neubert, R. H. Investigation of the molecular structure of the human stratum corneum ceramides [NP] and [EOS] by mass spectrometry. *Skin Pharmacol. Physiol.* **24**, 127-135 (2010).
- 5 Holthuis, J. C., Pomorski, T., Raggars, R. J., Sprong, H. & Van Meer, G. The organizing potential of sphingolipids in intracellular membrane transport. *Physiol. Rev.* **81**, 1689-1723 (2001).
- 6 Kihara, A., Mitsutake, S., Mizutani, Y. & Igarashi, Y. Metabolism and biological functions of two phosphorylated sphingolipids, sphingosine 1-phosphate and ceramide 1-phosphate. *Prog. Lipid Res.* **46**, 126-144 (2007).
- 7 Mizutani, Y., Mitsutake, S., Tsuji, K., Kihara, A. & Igarashi, Y. Ceramide biosynthesis in keratinocyte and its role in skin function. *Biochimie* **91**, 784-790 (2009).
- 8 Pontier, S. M. & Schweisguth, F. Glycosphingolipids in signaling and development: from liposomes to model organisms. *Dev. Dyn.* **241**, 92-106 (2012).
- 9 Hla, T. & Dannenberg, A. J. Sphingolipid signaling in metabolic disorders. *Cell Metab.* **16**, 420-434 (2012).
- 10 Kihara, A. & Igarashi, Y. Production and release of sphingosine 1-phosphate and the

- phosphorylated form of the immunomodulator FTY720. *Biochim. Biophys. Acta* **1781**, 496-502 (2008).
- 11 Bigaud, M., Guerini, D., Billich, A., Bassilana, F. & Brinkmann, V. Second generation S1P pathway modulators: Research strategies and clinical developments. *Biochim. Biophys. Acta* **1841**, 745-758 (2014).
 - 12 Okabe, K., Keenan, R. W. & Schmidt, G. Phytosphingosine groups as quantitatively significant components of the sphingolipids of the mucosa of the small intestines of some mammalian species. *Biochem. Biophys. Res. Commun.* **31**, 137-143 (1968).
 - 13 Iwamori, M., Costello, C. & Moser, H. W. Analysis and quantitation of free ceramide containing nonhydroxy and 2-hydroxy fatty acids, and phytosphingosine by high-performance liquid chromatography. *J. Lipid Res.* **20**, 86-96 (1979).
 - 14 Nishimura, K. Phytosphingosine is a characteristic component of the glycolipids in the vertebrate intestine. *Comp. Biochem. Physiol. B* **86**, 149-154 (1987).
 - 15 Pruett, S. T. *et al.* Biodiversity of sphingoid bases ("sphingosines") and related amino alcohols. *J. Lipid Res.* **49**, 1621-1639 (2008).
 - 16 Kihara, A. Sphingosine 1-phosphate is a key metabolite linking sphingolipids to glycerophospholipids. *Biochim. Biophys. Acta* **1841**, 766-772 (2014).
 - 17 Nakahara, K. *et al.* The Sjögren-Larsson syndrome gene encodes a hexadecenal dehydrogenase of the sphingosine 1-phosphate degradation pathway. *Mol. Cell* **46**, 461-471 (2012).
 - 18 Ohkuni, A., Ohno, Y. & Kihara, A. Identification of acyl-CoA synthetases involved in the mammalian sphingosine 1-phosphate metabolic pathway. *Biochem. Biophys. Res. Commun.* **442**, 195-201 (2013).
 - 19 Wakashima, T., Abe, K. & Kihara, A. Dual functions of the *trans*-2-enoyl-CoA reductase TER in the sphingosine-1-phosphate metabolic pathway and in fatty acid

- elongation. *J. Biol. Chem.* in press.
- 20 Ring, M. W., Schwär, G. & Bode, H. B. Biosynthesis of 2-hydroxy and iso-even fatty acids is connected to sphingolipid formation in myxobacteria. *ChemBioChem* **10**, 2003-2010 (2009).
- 21 Hoshi, M. & Kishimoto, Y. Synthesis of cerebronic acid from lignoceric acid by rat brain preparation. Some properties and distribution of the alpha-hydroxylation system. *J. Biol. Chem.* **248**, 4123-4130 (1973).
- 22 Umesaki, Y. *et al.* Presence of asialo GM₁ and glucosylceramide in the intestinal mucosa of mice and induction of fucosyl asialo GM₁ by conventionalization of germ-free mice. *J. Biochem.* **90**, 1731-1738 (1981).
- 23 Dick, K. J. *et al.* Mutation of FA2H underlies a complicated form of hereditary spastic paraplegia (SPG35). *Hum. Mutat.* **31**, E1251-1260 (2010).
- 24 Dickson, R. C., Sumanasekera, C. & Lester, R. L. Functions and metabolism of sphingolipids in *Saccharomyces cerevisiae*. *Prog. Lipid Res.* **45**, 447-465 (2006).
- 25 Skrzypek, M. S., Nagiec, M. M., Lester, R. L. & Dickson, R. C. Analysis of phosphorylated sphingolipid long-chain bases reveals potential roles in heat stress and growth control in *Saccharomyces*. *J. Bacteriol.* **181**, 1134-1140 (1999).
- 26 Mandala, S. M. *et al.* Sphingoid base 1-phosphate phosphatase: a key regulator of sphingolipid metabolism and stress response. *Proc. Natl. Acad. Sci. USA* **95**, 150-155 (1998).
- 27 Oshiro, J., Han, G. S. & Carman, G. M. Diacylglycerol pyrophosphate phosphatase in *Saccharomyces cerevisiae*. *Biochim. Biophys. Acta* **1635**, 1-9 (2003).
- 28 Kihara, A., Sano, T., Iwaki, S. & Igarashi, Y. Transmembrane topology of sphingoid long-chain base-1-phosphate phosphatase, Lcb3p. *Genes Cells* **8**, 525-535 (2003).
- 29 Huh, W. K. *et al.* Global analysis of protein localization in budding yeast. *Nature* **425**,

- 686-691 (2003).
- 30 Hansen, R. P., Shorland, F. B. & Cooke, N. J. The occurrence of n-heptadecanoic acid (margaric acid) in hydrogenated mutton fat. *Biochem. J.* **58**, 513-515 (1954).
- 31 Wheatley, V. R. & James, A. T. Studies of sebum. 7. The composition of the sebum of some common rodents. *Biochem. J.* **65**, 36-42 (1957).
- 32 Tove, S. B. Production of odd-numbered carbon fatty acids from propionate by mice. *Nature* **184**, 1647-1648 (1959).
- 33 Demarne, Y., Sacquet, E., Lecourtier, M. J. & Flanzy, J. Comparative study of endogenous fecal fatty acids in germ-free and conventional rats. *Am. J. Clin. Nutr.* **32**, 2027-2032 (1979).
- 34 Wanders, R. J. Metabolic functions of peroxisomes in health and disease. *Biochimie* **98**, 36-44 (2014).
- 35 Motta, S. *et al.* Ceramide composition of the psoriatic scale. *Biochim. Biophys. Acta* **1182**, 147-151 (1993).
- 36 Schulze, H. & Sandhoff, K. Sphingolipids and lysosomal pathologies. *Biochim. Biophys. Acta* **1841**, 799-810 (2014).
- 37 McCarthy, A. A. *et al.* Expression, crystallization and preliminary characterization of methylmalonyl coenzyme A epimerase from *Propionibacterium shermanii*. *Acta Crystallogr. D Biol. Crystallogr.* **57**, 706-708 (2001).
- 38 Brachmann, C. B. *et al.* Designer deletion strains derived from *Saccharomyces cerevisiae* S288C: a useful set of strains and plasmids for PCR-mediated gene disruption and other applications. *Yeast* **14**, 115-132 (1998).
- 39 Longtine, M. S. *et al.* Additional modules for versatile and economical PCR-based gene deletion and modification in *Saccharomyces cerevisiae*. *Yeast* **14**, 953-961 (1998).
- 40 Kihara, A. *et al.* Sphingosine-1-phosphate lyase is involved in the differentiation of F9

- embryonal carcinoma cells to primitive endoderm. *J. Biol. Chem.* **278**, 14578-14585 (2003).
- 41 Lester, R. L., Withers, B. R., Schultz, M. A. & Dickson, R. C. Iron, glucose and intrinsic factors alter sphingolipid composition as yeast cells enter stationary phase. *Biochim. Biophys. Acta* **1831**, 726-736 (2013).
- 42 Kihara, A., Kurotsu, F., Sano, T., Iwaki, S. & Igarashi, Y. Long-chain base kinase Lcb4 is anchored to the membrane through its palmitoylation by Akr1. *Mol. Cell. Biol.* **25**, 9189-9197 (2005).
- 43 Kihara, A. & Igarashi, Y. Identification and characterization of a *Saccharomyces cerevisiae* gene, *RSB1*, involved in sphingoid long-chain base release. *J. Biol. Chem.* **277**, 30048-30054 (2002).
- 44 Kyte, J. & Doolittle, R. F. A simple method for displaying the hydropathic character of a protein. *J. Mol. Biol.* **157**, 105-132 (1982).

Acknowledgements

This work was supported by a Grant-in-Aid for Scientific Research on Innovative Areas (25116701) from the Ministry of Education, Culture, Sports, Sciences and Technology of Japan. We are grateful to Professor Dr. J. H. Hagemann (Heinrich-Heine University) and Dr. T. Toyokuni for providing the pUG36 plasmid and for scientific editing of the manuscript, respectively. We also thank Dr. K. Obara and Ms. K. Nakahara for technical support.

Author contributions

A.K. planned the project, designed and performed experiments, analyzed the data, and wrote the manuscript. N.K. and Y.O. performed experiments and analyzed the data. T.O. and N.S. performed experiments. M.Y., T.K., and T.N. provided scientific advice.

Additional information

Supplementary information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interests.

FIGURE LEGENDS

Figure 1 | PHS is metabolized to both sphingolipids and glycerophospholipids. (a) Shown are pathways of sphingolipid biosynthesis and degradation as well as the enzymes involved in each step. In the *de novo* sphingolipid biosynthetic pathway, acyl-CoA and serine are condensed to 3-ketodihydrosphingosine (KDS), which is then reduced to DHS. DHS or its hydroxylated product PHS is acylated to generate ceramide (Cer). Cer is converted to complex sphingolipids (IPC, MIPC, and M(IP)₂C) by receiving phosphoinositol, mannose, and second phosphoinositol, successively. Some DHS and PHS are phosphorylated by LCB kinase to DHS1P and PHS1P, respectively. These LCBPs are either dephosphorylated by LCBP phosphatases to LCBs or cleaved between the 2- and 3-positions by the LCBP lyase Dpl1, generating fatty aldehydes (hexadecanal and 2-hydroxyhexadecanal, respectively) and phosphoethanolamine (P-Etn). Hexadecanal is converted to glycerophospholipids (PC, PE, PI, and PS) via palmitic acid and palmitoyl-CoA. However, the metabolism of 2-hydroxyhexadecanal is unclear and is the subject of this study. (b) [11,12-³H]PHS prepared in this study and [4,5-³H]DHS purchased from American Radiolabeled Chemicals, each at 0.1 μ Ci, together with cold DHS and PHS, each at 10 nmol, were separated by normal-phase TLC with chloroform/methanol/4.2 N ammonia (9:7:2, v/v). Labeled and cold lipids were detected by autoradiography (left panel) and ninhydrin staining (right panel), respectively. (c) [³H]PHS (8 nCi) was incubated with or without purified 3xFLAG-Lcb4 (250 ng) in the presence of 5 mM ATP at 37 °C for 1 hr. Lipids were extracted, separated by normal-phase TLC with chloroform/methanol/4.2 N ammonia (9:7:2, v/v), and detected by autoradiography. The asterisk indicates an undetermined PHS1P derivative. (d) BY4741 (wild-type) cells were labeled with 0.1 μ Ci [³H]DHS or [³H]PHS at 30 °C for 3 hr. Lipids were extracted, separated by normal-phase TLC with chloroform/methanol/4.2 N ammonia (9:7:2, v/v), and detected by autoradiography. (e) The radioactivity associated with each lipid in (d) was quantified, and the

data are expressed as the mean $\pm$ S.D. relative to the radioactivity in the total lipids from three independent experiments. Statistically significant differences are indicated (Student's *t*-test; *, $p < 0.05$; **, $p < 0.01$).

Figure 2 | PHS is metabolized to odd-numbered FAs in yeast. (a,b) BY4741 (wild-type) cells were labeled with 0.2 μ Ci [3 H]DHS or [3 H]PHS at 30 °C for 3 hr. Lipids were extracted, treated with alkaline solution, and separated by normal-phase TLC with chloroform/methanol/4.2 N ammonia (9:7:2, v/v). FAs were scraped from TLC, eluted from TLC silica, and methyl-esterified. (a) FAMES were separated by reverse-phase TLC with chloroform/methanol/water (5:15:1, v/v) and detected by autoradiography. The standard containing C14:0-, C15:0-, C16:0-, and C18:0 FAMES was a mixture of FA elongation products prepared in Supplementary Fig. 1. (b) FAMES were treated with/without aqueous KMnO₄/KIO₄, extracted with hexane, and separated by reverse-phase TLC with chloroform/methanol/water (5:15:1, v/v). (c-e) BY4741 and 3653 ($\Delta dpl1$) cells were grown in SC medium at 30 °C. Lipids were extracted from these cells and separated by normal-phase TLC with chloroform/methanol/4.2 N ammonia (9:7:2, v/v). PC was collected from TLC plates, and its composition was analyzed by UPLC-MS with a Xevo G2 QTof LC/MS system. The data were analyzed with MassLynx software. (c,e) Each bar represents the amount of each PC species (c, odd-numbered PC; e, even-numbered PC) and the mean $\pm$ S.D. from three independent experiments. (d) Values represent percent of the sum amounts of C15 PC species in (c) relative to those in wild-type cells. Statistically significant differences are indicated (Student's *t*-test; * $p < 0.05$, ** $p < 0.01$).

Figure 3 | PHS1P is more efficiently metabolized by LCBP phosphatases than LCBP lyase. (a,b) [32 P]DHS1P/[32 P]PHS1P (25 nCi, 0.5 μ M) was incubated with buffer alone (mock) or a 5

μg total membrane fraction prepared from SEY6210 (wild-type), KHY13 ($\Delta dpl1$), KHY20 ($\Delta lcb3$), and KHY49 ($\Delta lcb3 \Delta ysr3 \Delta lpp1 \Delta dpp1$) cells at 37 °C for 10 min. **(a)** The reaction products were separated by normal-phase TLC with 1-butanol/acetic acid/water (2:1:1, v/v) and detected by autoradiography. **(b)** The radioactivity associated with [^{32}P]phosphoethanolamine and [^{32}P]phosphate in the wild-type membrane in **(a)** was quantified using a bioimaging analyzer BAS-2500. Values represent the mean $\pm$ S.D. of the ratio of phosphoethanolamine to phosphate from three independent experiments. Statistically significant differences are indicated (Student's *t*-test; ***p* < 0.01). **(c)** Total membrane fractions (0.8 μg) prepared from KHY80 ($\Delta dpl1 \Delta lcb3 \Delta lpp1 \Delta dpp1 \Delta lcb4$) cells harboring the pAN9 (3xFLAG-DPL1) or pAN13 (3xFLAG-LCB3) plasmids were separated by SDS-PAGE, followed by immunoblotting with anti-FLAG antibody. **(d,e)** Total membrane fractions (0.5 μg) prepared from KHY80/pAN9 **(d)** and KHY80/pAN13 **(e)** were incubated with 25 nCi ^{32}P -labeled DHS1P or PHS1P at 0.3, 1, 3, and 10 μM at 37 °C for 10 min. After termination of the reaction, reaction products were separated by normal-phase TLC, followed by detection and quantification using a bioimaging analyzer BAS-2500. The radioactivity associated with the reaction product phosphoethanolamine **(d)** or phosphate **(e)** was quantified. The 3xFLAG-Dpl1 and 3xFLAG-Lcb3 levels in the membrane fractions used were determined relative to the level of the 3xFLAG-maltose binding protein standard and used for the calculation of the specific activity. We estimated that 1 μg of the total membrane fraction contained 3.2 ng Dpl1 and 8.9 ng Lcb3. The obtained values were expressed in a Lineweaver-Burk plot with a linear regression line. R indicates the correlation coefficient.

Figure 4 | Pentadecanoic acid is accumulated in the medium of acyl-CoA synthetase-deficient cells. **(a)** BY4741 (wild-type; wt), 2427 ($\Delta lcb4$), 3653 ($\Delta dpl1$), 6550 ($\Delta hfd1$), and AOY13 ($\Delta faa1 \Delta faa4$; $\Delta \Delta faa$) cells were incubated with 0.2 μCi [^3H]PHS at

30 °C for 3 hr. Lipids were extracted and separated by normal-phase TLC with chloroform/methanol/4.2 N ammonia (9:7:2, v/v). **(b)** BY4741 and AOY13 cells were incubated with 0.2 µCi [³H]DHS/[³H]PHS at 30 °C for 3 hr. Lipids were extracted from the cell pellets and cell culture medium, separated by normal-phase TLC with hexane/diethyl ether/acetic acid (30:70:1, v/v), and detected by autoradiography. Asterisk indicates unidentified lipid. 2-hydroxypalmitic acid (2-OH PA) was prepared by incubating [³H]PHS1P with purified S1P lyase and fatty aldehyde dehydrogenase. PA, palmitic acid. **(c)** AOY13 cells were incubated with 0.4 µCi [³H]DHS/[³H]PHS at 30 °C for 3 hr. Lipids were extracted from culture medium, saponified, and methyl-esterified. The resulting FAMES were separated by reverse-phase TLC.

Figure 5 | *MPO1* is involved in PHS-to-glycerophospholipid metabolism. **(a,b)** BY4741 (wild-type) and 4378 ($\Delta mpo1$) cells were incubated with 0.1 µCi [³H]DHS/[³H]PHS at 30 °C for 3 hr. Lipids were extracted from the cells **(a,b)** and cell culture medium **(b)**, separated by normal-phase TLC with chloroform/methanol/4.2 N ammonia (9:7:2, v/v) **(a)** and hexane/diethyl ether/acetic acid (30:70:1, v/v) **(b)**, and detected by autoradiography. PA, palmitic acid; 2-OH PA, 2-hydroxypalmitic acid. **(c)** A hydrophilicity plot of the Mpo1 protein was generated using MacVector software (MacVector, Inc, Cary, NC) with Kyte and Doolittle hydrophilicity values⁴⁴ (window size of 15). **(d)** Amino acid sequence of the Mpo1 protein. The underlined region shares a weak sequence similarity with the NAD⁺-binding domain of Sir2. **(e)** YOY158 (*SEC61-mCherry*) cells bearing a pNK47 (*GFP-MPO1*) plasmid were grown in SC medium lacking uracil at 30 °C. Expression of GFP-Mpo1 was induced by transferring cells to SC medium lacking uracil and methionine for 30 min. Cells were analyzed by fluorescence microscopy using a DM5000B fluorescence microscope (Leica Microsystems, Wetzlar, Germany). Bar, 5 µm.

Figure 6 | PHS is metabolized to odd-numbered FAs in mammals. (a-e) CHO-K1 (a,b), HEK 293T (c,d), and mouse embryonic carcinoma F9 (*Sp1*^{+/+}) and F9-2 (*Sp1*^{-/-}) cells (e) were labeled with 0.1 μ Ci (2.25 nmol) [³H]DHS or [³H]PHS at 30 °C for 5 hr. (a,c,e) Lipids were extracted, treated with or without alkaline solution, and separated by normal-phase TLC with 1-butanol/acetic acid/water (3:1:1, v/v). HexCer, monohexosylceramide; LacCer, lactosylceramide; GSL, glycosphingolipid; SM, sphingomyelin. (b,d) Lipids were extracted and treated with alkaline solution. The released FAs were methyl-esterified, and the resulting FAMES were incubated with or without KMnO₄/KIO₄ solution, extracted with hexane, and separated by reverse-phase TLC with chloroform/methanol/water (5:15:1, v/v). The standard containing C14:0-, C15:0-, C16:0-, and C18:0-FAMES was a mixture of FA elongation products prepared as in Supplementary Fig. 1.

Figure 7 | Proposed metabolic pathway of PHS to pentadecanoyl-CoA. The proposed pathway and enzymes responsible for PHS metabolism in yeast are shown. PHS is first phosphorylated by LCBP kinase Lcb4. The resulting PHS1P is cleaved to 2-hydroxyhexadecanal by LCBP lyase Dpl1. The fatty aldehyde dehydrogenase Hfd1 oxidizes 2-hydroxyhexadecanal to 2-hydroxypalmitic acid. Faa1, Faa4, and probably other acyl-CoA synthetases may catalyze conversion of 2-hydroxypalmitic acid to 2-hydroxypalmitoyl-CoA. Mpo1 may convert 2-hydroxypalmitoyl-CoA to pentadecanoic acid or another metabolite leading to pentadecanoic acid. Finally, pentadecanoic acid is activated to pentadecanoyl-CoA by the acyl-CoA synthetases Faa1 and Faa4 and incorporated into glycerophospholipids. Open arrows represent reactions that are yet to be fully elucidated.

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7

