



HOKKAIDO UNIVERSITY

Title	Hangeshashinto Improves the Completion Rate of Chemoradiotherapy and the Nutritional Status in Patients with Head and Neck Cancer
Author(s)	Hatakeyama, Hiromitsu; Takahashi, Hiroki; Oridate, Nobuhiko et al.
Citation	ORL-journal for oto-rhino-laryngology head and neck surgery, 77(2), 100-108 https://doi.org/10.1159/000381026
Issue Date	2015-05
Doc URL	https://hdl.handle.net/2115/59425
Type	journal article
File Information	manuscript.pdf



Hangeshashinto improves the completion rate of chemoradiotherapy and nutritional status in head and neck cancer patients.

Hiromitsu Hatakeyama M.D., Ph.D.¹⁾; Hiroki Takahashi M.D.¹⁾, Nobuhiko Oridate M.D., Ph.D.²⁾; Rinnosuke Kuramoto M.D.¹⁾; Keishi Fujiwara M.D., Ph. D.¹⁾; Tomohiro Sakashita M.D., Ph. D.¹⁾; Satoshi Kano M.D., Ph. D.¹⁾; Takatsugu Mizumachi M.D., Ph. D.¹⁾ Akihiro Homma M.D., Ph.D.¹⁾; Hiroshi Takeda M.D., Ph.D.³⁾ and Satoshi Fukuda M.D., Ph.D.¹⁾

1) Department of Otolaryngology-Head and Neck Surgery, Hokkaido University Graduate School of Medicine, Sapporo, Japan

2) Department of Otorhinolaryngology and Head and Neck Surgery, Yokohama City University School of Medicine, Yokohama, Japan

3) Department of Pathophysiology and Therapeutics, Faculty of Pharmaceutical Sciences, Hokkaido University, Sapporo, Japan

Corresponding author: Hiromitsu Hatakeyama

Mailing address: Department of Otolaryngology-Head and Neck Surgery, Hokkaido University Graduate School of Medicine, Kita 15, Nishi 7, Kita-ku, Sapporo 060-8638, Japan

- 1 Phone: +81-11-706-5958
- 2 Fax: +81-11-717-7566
- 3 E-mail: htkym@med.hokudai.ac.jp
- 4 Keywords: Mucositis, Hangeshashinto (TJ-14), Chemoradiotherapy, Head and Neck Cancer,
- 5 Nutrition
- 6 Short title: TJ-14 improves the completion rate of CRT in HNC
- 7 COI : The authors declare that they have no conflict of interests
- 8
- 9 Funding : H.H. and A.H. were supported, in part, by a Grant-in-Aid for Scientific Research from the
- 10 Ministry of Education, Science and Culture of Japan

Abstract

Purpose: Severe oral and pharyngeal mucositis is one of the most critical toxicities known to lead to the discontinuation of chemoradiotherapy (CRT) for head and neck cancer (HNC). Hangeshashinto (TJ-14) is a Kampo medicine that relieves chemotherapy-induced oral mucositis. We investigated the effect of TJ-14 on mucositis, nutritional status and completion rate of CRT.

Methods: The study group comprised patients with advanced HNC who treated with concomitant weekly cisplatin administration and 70 Gy of radiotherapy. The primary end point was the completion rate of chemotherapy, and the secondary end points were the grade of mucositis, and nutritional status.

Results: Total 57 patients were included in this study. The completion rate of CRT among patients who treated with TJ-14 was 91.4%. There was a significant difference in the completion rate of CRT between the groups with and without TJ-14. ($p=0.0452$) The reduction in body weight was significantly improved from 10.89% to 5.89 % with TJ-14 administration ($p=0.003$) and the reduction in serum albumin was also significantly decreased from 17.37% to 8.73 %. ($p=0.024$)

Conclusion: This therapy allowed a high completion rate of CRT as well as significant benefits in terms of nutritional status. We plan to carry out a further large-scale study of TJ-14.

Introduction

Concomitant chemotherapy with radiation for HNC provides a favorable prognosis as well as a significantly higher rate of organ preservation in comparison with radiation alone. [1] [2] Radiation with high-dose cisplatin every 3 weeks remains the most standard regimen in US Cooperative Group clinical trials. However, the toxicity of this regimen is too severe for most Japanese patients to complete; therefore, our institute has introduced a regimen of radiation with weekly 40 mg/m² of cisplatin. [3] We have to date achieved good clinical outcomes with acceptable toxicity in terms of neutropenia and renal failure. [4] However, the completion rate of this regimen remains around 60 % and severe oral and pharyngeal mucositis sometimes causes the discontinuation of the CRT.

Hangeshashinto (TJ-14) is a Japanese traditional medicine (Kampo) that includes 7 natural extracts: *Pinelliae Tuber*, *Scutellariae Radix*, *Zingiberis Rhizoma*, *Ginseng Radix*, *Glycyrrhizae Radix*, *Zizyphi Fructus*, and *Coptidis Rhizoma*. TJ-14 has been reported to regulate prostaglandin E₂ in in vivo colitis models and to reduce chemotherapy-induced severe oral mucositis in gastric cancer patients. [5] [6] [7] The degree of severity of oral and pharyngeal mucositis varies during CRT for pharyngeal cancer, and new agents are needed to improve completion rates and to limit pain in these patients.

This study evaluates the effect of TJ-14 on the completion rate of CRT in patients with cancers of the oropharynx and hypopharynx through the regulation of CRT-induced oral and pharyngeal mucositis.

This is the first trial to use TJ-14 as a gargle to prevent mucositis induced by chemoradiation. In this study we investigated the compliance of TJ-14 for CRT in oro- and hypopharyngeal cancer and the complete rate. Furthermore, nutritional status, grade of mucositis and pain were estimated in comparison with the patient treated without TJ-14.

Patients and Methods

Patients

The study group comprised Japanese patients who were diagnosed with advanced oropharyngeal and hypopharyngeal squamous cell carcinoma and treated with CRT at Hokkaido University Hospital, Japan. Written informed consent was obtained from all patients before entry into the study. This study was approved by the Institutional Review Board of Hokkaido University Hospital (UMIN000006461).

CRT and TJ-14 administration

All patients were treated with a total dose of 70 Gy of radiation in 35 daily fractions. After the initial dose of 40 Gy had been administered, an additional dose of 30 Gy was given with a shrunken field in 15 fractions over 3 weeks.

Weekly cisplatin was administered at a dose of 40mg/m² on week 1, 2, 3, 4, 6 and 7 of radiation

therapy. The intended maximum total dose of cisplatin was 240mg/m².

Patients received 2.5g of TJ-14 as a gargle three times a day as follows: 1) 2.5g of TJ-14 was dissolved in 40ml of drinking water and the patient rinsed the oral cavity and pharynx with the solution, without swallowing, for more than 5 seconds, 2) TJ-14 was expelled out after rinsing and was not to be swallowed, and 3) patient were advised not to consume any food or drink within 30 minutes after rinsing with TJ-14. CRT and TJ-14 administration schedules are shown in Figure 1.

Evaluation of mucositis, pain, and nutritional status

Oral and pharyngeal mucositis were graded as according to CTCAE v4.0. The maximum dose of morphine was defined as that on day 57. Body composition of all patients was estimated by bioelectric impedance analysis using an InBody S20 body composition analyzer with InBody 3.0 software (BioSpace) on Day 0 and Day 57. [8] Nutritional intake was assessed by the Nutrition Support Team of Hokkaido University Hospital, by measuring the weight of each dish before and after every meal. Furthermore, the calories of all additional foods, snacks, and drinks were added to the measured food intake. Total oral intake of calories and protein were recorded as the average of 3 days on 1st, 3rd, 6th and 8th week. The serum levels of albumin, pre-albumin and retinol binding protein were used as nutritional status markers and were estimated on day 1, 22, 43, 57.

Statistical Considerations

The primary endpoint was the completion rate of CRT. Complete chemoradiation treatment delivery was defined as the administration of full dose of 70 Gy radiation within 63 days, and the completion of at least five of six courses of cisplatin. Complete administration of TJ-14 was defined as more than 7 weeks treatment without interruption. The secondary endpoints were oral and pharyngeal mucositis grade, nutritional status and maximum dose of morphine.

Results

Patient characteristics

Demographic and clinicopathological data, including age, gender, tumor site, TNM stage, smoking and alcohol abuse of all patients are shown in Table 1. Twenty three patients with oropharyngeal cancer and 34 patients with hypopharyngeal cancer were included in this study. Forty patients had Stage IV disease, 6 had Stage III disease and 10 had Stage II. Of 57 patients, twelve patients (11 male and 1 female) who received TJ-14 treatment were enrolled in the study and were evaluated from January 2012 through September 2013. Other 45 patients treated without TJ-14 from June 2006 through January 2011. The patients treated with TJ-14 ranged in age from 40 to 70 years (median = 60 years) and the patients treated without TJ-14 ranged from 40 to 75 years (median 59 years). All patients received oral care from the Dental Care Team of Hokkaido University Hospital during

CRT.

TJ-14 compliance

A total 9 of the 12 patients completed 7 or 8 weeks of treatment with TJ-14. Three patients could not continue to use TJ-14, but they could complete the CRT. One patient discontinued TJ-14 on day 6 due to its bitter taste. One patient could not keep himself from eating or drinking for 30 minutes after gargling with TJ-14, and stopped the study on day 16. The third patient discontinued TJ-14 on day 38 due to vomiting and nausea. No patients interrupted TJ-14 treatment due to oral and pharyngeal mucositis.

Chemoradiotherapy completion rate

Completion rate of chemoradiotherapy were summarized in Table 2. All patients who treated with TJ-14 received the full dose of RT (70Gy) without interruption. A total of 11 (91.6%) patients completed five (5 patients) or six (6 patients) courses of chemotherapy. One patient could not receive a 5th course of chemotherapy due to fever. Eight of 9 (88.9%) patients who completed treatment with TJ-14 also completed CRT. Eighteen of 45 patients treated without TJ-14 interrupted the CRT. The cause of Chemotherapy interruption were neutropenia in 4 cases, fever in 4, renal failure in 3, liver dysfunction in 1, mucositis in 1, tumor bleeding in 1, and the rejection by 2 patients. The

radiation course was extended in 2 cases by holidays and machine maintenance. [9]

There was a significant difference in the completion rate of CRT between the groups with and without TJ-14. ($p=0.0452$, Fisher's exact test)

Mucositis and Pain

There were no significant differences in the maximum grade of mucositis and the maximum dose of morphine between with and without TJ-14. (Table 3). Grade 3 mucositis were developed in 5 (41.7%) patients and Grade 4 mucositis was absent in the TJ-14 administration group, whereas Grade 3 mucositis were developed in 25 (57.8%) patients and one of Grade 4 mucositis occurred in TJ-14 non-administration group. Significant difference was not observed in mean maximum dose of morphine between Grade 1-2 mucositis and Grade 3-4.

Body composition analysis and nutritional status

We investigated the body composition index of the patients treated with TJ-14 to reveal their nutritional status and the results were summarized in Table 4. The median body weight of the patients was significantly decreased after chemoradiation ($p=0.006$) with a median decrease in body weight of 5.89 %. (Maximum 11.7% and minimum -2.8%). Median fat and bone mass did not change during CRT, but the muscle mass and total body water were significantly decreased by 5.7%

and 5.5%, respectively (Muscle mass: $p=0.045$, Body water: $p=0.049$). The median metabolic rate also decreased by about 4.0% during CRT. Table 5 shows the changes in median oral intake calories and nutritional status markers in the serum. The median total oral intake of calories drastically decreased after the 4th week. Seven patients had used PEG before CRT, and 4 of the 7 patients started to use PEG due to oral feeding difficulties on day 21, 36, 45 and 47, respectively. The median level of albumin and pre-albumin in the serum gradually decreased throughout the course of CRT, but the changes were not statically significant. The mean serum concentration of retinoid-binding protein increased on the 3rd week, but decreased again after the 6th week.

The median reduction in body weight during CRT without TJ-14 was 10.72% and the reduction in serum albumin was 17.37%, whereas the reduction in body weight was significantly improved up to 5.89 % with TJ-14 administration ($p=0.003$) and the reduction in serum albumin was also significantly decreased to 8.73 % in this study. ($p=0.024$) (Figure 2)

Discussion

Toxicities with CRT often lead to treatment interruptions that are invariably associated with a poorer outcome.[10] [11]To improve the completion rate of CRT for HNC, we modified the regimen and improved the management program in terms of the hydration protocol and the adequate administration of G-CSF to avoid severe leucopenia, neutropenia and renal failure.[4]Although

1 nausea and vomiting were one of the most severe side effects of cisplatin administration, it was
2 dramatically improved with the administration of dexamthasthon and serotonin 5-HT3 receptor
3 antagonists. [12] On the other hand, the pain and the dysphagia associated with severe oral and
4 pharyngeal mucositis have been a critical factor in completion rate. Most patients with
5 oropharyngeal and hypopharyngeal cancer undergoing CRT receive direct irradiation to oral cavity
6 and pharynx, so it is easy to assume that therapy-induced mucositis is more severe than in other
7 digestive cancers. Severe direct irradiation-induced mucositis led us to apply TJ-14 for
8 oropharyngeal and hypopharyngeal cancers at first, with the gargle therapy involving TJ-14
9 achieving good compliance with a 75% completion rate. Only one patient had an early interruption
10 of treatment with TJ-14 due to its bitter taste, and no patient discontinued treatment due to oral
11 mucositis or pain.

12 The pilot study of TJ-14 for chemotherapy-induced oral mucositis in gastric cancer showed a
13 significant effect. [13] However, a double-blind, placebo-controlled, randomized phase II study of
14 TJ-14 did not show any difference in oral mucositis grade, but a tendency for TJ-14 to reduce the
15 duration of chemotherapy-induced oral mucositis from 17 days to 9 days was observed. [7] We
16 previously reported that the rate of grade 3-4 mucositis induced by the same regimens of CRT for
17 HNC were 39.6%. [4] If limited to oropharyngeal and hypopharyngeal cancers, the rate of grade 3-4
18 mucositis was 51.6%. [9] In this study, no patients had grade 4 mucositis, and the rate of grade 3

1 mucositis decreased to 41.6%. Unlike the study on gastric cancer, no statistical significant effect of
2 TJ-14 in terms of mucositis grade could be observed. However, no patient interrupted CRT due to
3 mucositis and almost grade 2-3 mucositis were occurred at late phase of CRT in TJ-14 administrated
4 group. These facts might be induced the beneficial change in the complete rate of CRT and the
5 nutrition status. No statistical difference in oral mucositis might be caused from the fuzzy grading
6 system. The grading in oral mucositis of CTCAE v4.0 has only 5 grades and lacks the clarity, and the
7 real extent of mucositis was different in the same grade. Therefore, a more extensive method of
8 evaluation mucositis in place of CTCAE v4.0 is needed.

9 Molecular targeted therapy using cetuximab, an epidermal growth factor receptor monoclonal
10 antibody, has recently been investigated in conjunction with radiation therapy for advanced HNC
11 patients, and has shown promising results. [14] [15] Radiation therapy with cetuximab does not cause
12 toxicity leading to leucopenia, neutropenia or renal failure, and is relatively-safe compared to
13 platinum-based CRT. However, severe mucositis occurs more often during radiation therapy with
14 cetuximab, so a new agent to prevent radiation-induced mucositis is needed. TJ-14 may be a
15 promising option for use in combination with cetuximab treatment for HNC.

16 One of the reasons for the improvement in completion rate of CRT with TJ-14 is the good outcome
17 in terms of nutritional status. Recently, several studies have suggested that early and intensive
18 nutritional support improved the quality of life and outcome of patient with HNC undergoing RT or

CRT. [16, 17] [18] [19, 20] Capuano et.al. showed that a weight reduction of more than 20% significantly increased CRT-induced toxicity, risk of early mortality and hospital readmission rate. [21]To improve nutrition, we introduced oral care, a PEG prior to treatment and a pain management program based on the combined efforts of dentists and supportive-care staff. Despite the introduction of a PEG and oral care, the median reduction in body weight during CRT remained around 10 %. It is interesting to note that most of the body weight loss involved the loss of body water and muscle mass, so that the control of dehydration and muscle maintenance may improve nutritional status. Although this study was a single-arm trial, not a controlled randomized trial with a placebo, these results suggest that improved nutritional status might contribute to a higher completion rate of CRT. TJ-14 has been known as an agent for enhancing digestive activity, and it was originally used for acute and chronic gastritis, heartburn, and diarrhea. Patients often swallowed TJ-14 after gargling, so the possibility that it had an effect on digestive activity cannot be totally excluded. The anti-inflammatory effect of TJ-14 has been explained by its direct regulation of PGE2 and COX-2, but the mechanism by which it enhances digestive activity remains to be clarified. [5, 6] The effects of Kampo drugs has been known historically and confirmed empirically, and their mechanisms have been gradually revealed both biologically and pharmacologically. One of the advantages of Kampo drugs is their limited side effects, and several Kampo drugs have been used for supportive treatment for various cancers. [22-24] Although most Kampo drugs are composed and used according to

1 traditional strategies, we should accumulate the cases and provide clinical evidence for their
2 effective utilization. Seven components are included in TJ-14 and these interact with each other.
3 Therefore, to reveal the mechanism of TJ-14, we need to investigate not only the function of all
4 components, but also their cross-reactions.

5 In conclusion, the feasibility of TJ-14 for HNC CRT-induced oral mucositis was confirmed.
6 Treatment with TJ-14 was shown to benefit the completion rate of CRT as well as nutritional status.
7 Next, a large-scale, randomized prospective controlled study is needed to define the effects of TJ-14
8 on the completion rate of CRT, nutritional status, and survival benefits for patients with HNC.

9
10 Acknowledgement : This work was supported by Nutrition Support Team of Hokkaido University
11 Hospital and a Health and Labour Sciences Research Grant for Clinical Cancer Research (H26-141)
12 from the Ministry of Health, Labour and Welfare of Japan and the National Cancer Center Research
13 and Development Fund (26-A-4) of Japan.

Figure Legends

Figure 1 Chemoradiation and TJ-14 administration schedules

Figure 2 Comparison between TJ-14 administration group and non-administration group

A, Reduction in total body weight

B, Reduction in the level of serum albumin

Table 1 Clinical characteristics of all patients.

Table 2 Completion rate of chemoradiotherapy

Numbers shown in parentheses are the number of patients who completed the treatment of TJ-14 and the completion rates

Table 3 Mucositis grade and maximum dose of morphine

Numbers shown in parentheses are the number of patients who interrupted the TJ-14 treatment

Table 4 Changes in total body weight and body composition during chemoradiotherapy in TJ-14 administration group

Table 5 Changes in oral intake and nutritional status markers in TJ-14 administration group

1

2

3

4

1 Forastiere AA, Goepfert H, Maor M, Pajak TF, Weber RS, Morrison W, Glisson B, Trotti A, Ridge JA, Chao C, Peters G, Lee D-J, Leaf A, Ensley J, Cooper J: Concurrent chemotherapy and radiotherapy for organ preservation in advanced laryngeal cancer. The New England Journal of Medicine 2003;349:2091-2098.

2 Forastiere AA, Trotti A, Pfister DG, Grandis JR: Head and neck cancer: Recent advances and new standards of care. J Clin Oncol 2006;24:2603-2605.

3 Taki S, Homma A, Oridate N, Suzuki S, Suzuki F, Sakashita T, Fukuda S: Salvage surgery for local recurrence after chemoradiotherapy or radiotherapy in hypopharyngeal cancer patients. European archives of oto-rhino-laryngology : official journal of the European Federation of Oto-Rhino-Laryngological Societies 2010;267:1765-1769.

4 Homma A, Inamura N, Oridate N, Suzuki S, Hatakeyama H, Mizumachi T, Kano S, Sakashita T, Onimaru R, Yasuda K, Shirato H, Fukuda S: Concomitant weekly cisplatin and radiotherapy for head and neck cancer. Jpn J Clin Oncol 2011;41:980-986.

5 Kase Y, Saitoh K, Ishige A, Komatsu Y: Mechanisms by which hange-shashin-to reduces prostaglandin e2 levels. Biol Pharm Bull 1998;21:1277-1281.

6 Kase Y, Saitoh K, Makino B, Hashimoto K, Ishige A, Komatsu Y: Relationship between the antidiarrhoeal effects of hange-shashin-to and its active components. Phytotherapy research : PTR 1999;13:468-473.

7 Aoyama T, Nishikawa K, Takiguchi N, Tanabe K, Imano M, Fukushima R, Sakamoto J, Oba MS, Morita S, Kono T, Tsuburaya A: Double-blind, placebo-controlled, randomized phase ii study of tj-14 (hangeshashinto) for gastric cancer chemotherapy-induced oral mucositis. Cancer Chemother Pharmacol 2014

8 Torimoto K, Hirayama A, Samma S, Yoshida K, Fujimoto K, Hirao Y: The relationship between nocturnal polyuria and the distribution of body fluid: Assessment by bioelectric impedance analysis. J Urol 2009;181:219-224; discussion 224.

9 Kuramoto R, Oridate N, Homma A, Oikawa K, Fujita K, Suzuki S, Hatakeyama H, Kano S, Mizumachi T, Sakashita T, Fukuda S: The role of tube feeding for chemoradiotherapy in head and neck cancer. Deglutition 2012;1:359-363.

10 Bernier J, Domenge C, Ozsahin M, Matuszewska K, Lefebvre JL, Greiner RH, Giralt J, Maingon P, Rolland F, Bolla M, Cognetti F, Bourhis J, Kirkpatrick A, van Glabbeke M: Postoperative irradiation with or without concomitant chemotherapy for locally advanced head and neck cancer. *N Engl J Med* 2004;350:1945-1952.

11 Cooper JS, Pajak TF, Forastiere AA, Jacobs J, Campbell BH, Saxman SB, Kish JA, Kim HE, Cmelak AJ, Rotman M, Machtay M, Ensley JF, Chao KS, Schultz CJ, Lee N, Fu KK: Postoperative concurrent radiotherapy and chemotherapy for high-risk squamous-cell carcinoma of the head and neck. *N Engl J Med* 2004;350:1937-1944.

12 Latreille J, Pater J, Johnston D, Laberge F, Stewart D, Rusthoven J, Hoskins P, Findlay B, McMurtrie E, Yelle L, Williams C, Walde D, Ernst S, Dhaliwal H, Warr D, Shepherd F, Mee D, Nishimura L, Osoba D, Zee B: Use of dexamethasone and granisetron in the control of delayed emesis for patients who receive highly emetogenic chemotherapy. National cancer institute of canada clinical trials group. *J Clin Oncol* 1998;16:1174-1178.

13 Kono T, Kaneko A, Matsumoto C, Miyagi C, Ohbuchi K, Mizuhara Y, Miyano K, Uezono Y: Multitargeted effects of hangeshashinto for treatment of chemotherapy-induced oral mucositis on inducible prostaglandin e2 production in human oral keratinocytes. *Integrative cancer therapies* 2014

14 Bonner JA, Harari PM, Giralt J, Cohen RB, Jones CU, Sur RK, Raben D, Baselga J, Spencer SA, Zhu J, Youssoufian H, Rowinsky EK, Ang KK: Radiotherapy plus cetuximab for locoregionally advanced head and neck cancer: 5-year survival data from a phase 3 randomised trial, and relation between cetuximab-induced rash and survival. *Lancet Oncol* 2010;11:21-28.

15 Pfister DG, Su YB, Kraus DH, Wolden SL, Lis E, Aliff TB, Zahalsky AJ, Lake S, Needle MN, Shaha AR, Shah JP, Zelefsky MJ: Concurrent cetuximab, cisplatin, and concomitant boost radiotherapy for locoregionally advanced, squamous cell head and neck cancer: A pilot phase ii study of a new combined-modality paradigm. *J Clin Oncol* 2006;24:1072-1078.

16 List MA, Mumby P, Haraf D, Siston A, Mick R, MacCracken E, Vokes E: Performance and quality of life outcome in patients completing concomitant chemoradiotherapy protocols for head and neck cancer. *Quality of life research : an international journal of quality of life aspects of treatment, care and rehabilitation* 1997;6:274-284.

17 List MA, Siston A, Haraf D, Schumm P, Kies M, Stenson K, Vokes EE: Quality of life and performance in advanced head and neck cancer patients on concomitant chemoradiotherapy: A prospective examination. *J Clin Oncol* 1999;17:1020-1028.

18 Valentini V, Marazzi F, Bossola M, Micciche F, Nardone L, Balducci M, Dinapoli N,

- 1 Bonomo P, Autorino R, Silipigni S, Giuliani F, Tamanti C, Mele MC, Martorana GE:
2 Nutritional counselling and oral nutritional supplements in head and neck cancer patients
3 undergoing chemoradiotherapy. *Journal of human nutrition and dietetics : the official*
4 *journal of the British Dietetic Association* 2012;25:201-208.
- 5 19 Paccagnella A, Morello M, Da Mosto MC, Baruffi C, Marcon ML, Gava A, Baggio V,
6 Lamom S, Babare R, Rosti G, Giometto M, Boscolo-Rizzo P, Kiwanuka E, Tessarin M,
7 Caregaro L, Marchiori C: Early nutritional intervention improves treatment tolerance and
8 outcomes in head and neck cancer patients undergoing concurrent chemoradiotherapy.
9 *Supportive care in cancer : official journal of the Multinational Association of Supportive*
10 *Care in Cancer* 2010;18:837-845.
- 11 20 Ravasco P, Monteiro-Grillo I, Marques Vidal P, Camilo ME: Impact of nutrition on
12 outcome: A prospective randomized controlled trial in patients with head and neck cancer
13 undergoing radiotherapy. *Head Neck* 2005;27:659-668.
- 14 21 Capuano G, Grosso A, Gentile PC, Battista M, Bianciardi F, Di Palma A, Pavese I,
15 Satta F, Tosti M, Palladino A, Coiro G, Di Palma M: Influence of weight loss on outcomes in
16 patients with head and neck cancer undergoing concomitant chemoradiotherapy. *Head Neck*
17 2008;30:503-508.
- 18 22 Hosokawa A, Ogawa K, Ando T, Suzuki N, Ueda A, Kajiura S, Kobayashi Y,
19 Tsukioka Y, Horikawa N, Yabushita K, Fukuoka J, Sugiyama T: Preventive effect of
20 traditional japanese medicine on neurotoxicity of folfox for metastatic colorectal cancer: A
21 multicenter retrospective study. *Anticancer Res* 2012;32:2545-2550.
- 22 23 Mori K, Kondo T, Kamiyama Y, Kano Y, Tominaga K: Preventive effect of kampo
23 medicine (hangeshashin-to) against irinotecan-induced diarrhea in advanced non-small-cell
24 lung cancer. *Cancer Chemother Pharmacol* 2003;51:403-406.
- 25 24 Endo S, Nishida T, Nishikawa K, Nakajima K, Hasegawa J, Kitagawa T, Ito T,
26 Matsuda H: Dai-kenchu-to, a chinese herbal medicine, improves stasis of patients with total
27 gastrectomy and jejunal pouch interposition. *Am J Surg* 2006;192:9-13.

Figure. 1





Figure. 2

A



B




Table 1			
Clinical Characteristics		TJ-14(n=12)	No TJ-14(n=45)
Age (years)			
	Range	40-70	40-75
	Median	60	59
Sex ratio(M/F)			
	Male	11	42
	Female	1	3
Tumor Site			
	Oropharynx	3	20
	Hypopharynx	9	25
Tumor Stage			
	II	0	10
	III	3	3
	IV	9	32
Perfomance Status			
	0	10	44
	1	1	1
	2	1	0
Heavy use of Alcohol			
	Yes	5	21
	No	7	24
Smoker			
	Yes	12	40
	No	0	5
PEG			
	Yes	7	32
	No	5	13

Table 2. Completion Rate of CRT			
	TJ-14	No TJ-14	<i>p</i>
CRT Complete	11(8)	27	
CRT Failure	1(1)	18	
Completion Rate	91.6% (88.9%)	60.00%	0.0452

*Numbers shown in parentheses are the number of patients who completed the treatment of TJ-14 and the completion rates

Table 3			
Mucositis Grade and Maximum dose of morphine			
	TJ-14	No TJ-14	<i>p</i>
Grade 1	0	4	
Grade 2	7 (2)	15	
Grade 3	5 (1)	25	
Grade 4	0	1	n.s.
Maximum dose of morphine (mg/day)	88.5±65.4	42.1±42.52	n.s.

*Numbers shown in parentheses are the number of patients who interrupted the TJ-14 treatment

Table 4.				
Body weight and Body composition pf the patients treated with TJ-14				
	1st week	8th week	<i>p</i>	Change
Median Body Weghit (kg)	60.8±13.58	57.06±9.97	0.006	-5.89%
Body Fat Percentage(%)	19.05±6.34	20.62±5.74	0.494	8.20%
Total Body Muscle (kg)	46.13±8.81	43.52±6.35	0.045	-5.65%
Total Body Water (kg)	36.07±6.88	34.07±4.70	0.049	-5.54%
Total Body Bone (kg)	2.59±0.438	2.53±0.284	0.103	-2.32%
Basal Metabolic Rate(kcal)	1422.33±199.5	1364.33±137.16	0.046	-4.08%

Table 5. Nutritional Status

	1st week	4th week	6th week	8th week	<i>p</i>
Total oral intake (kcal/day)	2051.66±318.64	1701.3±676.96	1181.1±844.97	1108.71±671.51	0.013
Oral protein intake (g/day)	71.70±11.25	61.07±23.45	39.87±28.91	38.74±26.06	0.024
Albumin (g/dl)	3.94±0.41	3.70±0.28	3.66±0.39	3.63±0.45	0.056
Pre-albumin (mg/dl)	29.76±6.85	27.94±5.34	25.94±6.94	23.32±6.89	0.053
Retinol binding protein (mg/dl)	3.29±0.98	3.54±0.88	3.08±0.58	2.96±0.91	0.284

Changes in oral intake and nutritional status markers.