



HOKKAIDO UNIVERSITY

Title	北海道植物志料 V
Author(s)	宮部, 金吾; 工藤, 祐舜
Citation	札幌博物学会会報, 6(1), 1-9
Issue Date	1915-12-01
Doc URL	https://hdl.handle.net/2115/61256
Type	journal article
File Information	Vol.6No.1_001.pdf



MATERIALS FOR A FLORA OF HOKKAIDO. V.

By

KINGO MIYABE

and

YŪSHUN KUDŌ.

北海道植物志料 V.

宮 部 金 吾

工 藤 祐 舜

45. **Hydrilla verticillata** Royle, Ill. Bot. Himal. t. 376 (1839); Caspary, in Pringsh. Jahrb. I. p. 494; Hook. f. Fl. Brit. Ind. V. p. 659; Forb. et Hemsl. Ind. Fl. Sin. III. p. 1; Aschers. u. Gürke, in Engl. u. Prantl, Nat. Pfl.-fam. II. 1. p. 250. f. 184. a, b; Korsch. Acta Hort. Peterop. XII. p. 394; Aschers. u. Graebn. Syn. Mitteleurop. Fl. I. p. 398; Kawakami, Bot. Mag. Tokyo XII. p. (225); Diels, in Engl. Bot. Jahrb. XXIX. p. 220; Kom. Fl. Mansh. I. p. 237; Matsum. et Hayata, Enum. Pl. Formos. p. 405; Nakai, Fl. Kor. II. p. 216.

Serpicula verticillata L. f. Suppl. p. 416.—*Hydrilla verticillata*, var. *Roxburghii* Casp. l. c.; Miq. Prol. Fl. Jap. p. 159; Fr. et Sav. Enum. Pl. Jap. II. p. 17; Matsum. Ind. Pl. Jap. II. 1. p. 32.—*Hydrilla japonica* Miq. l. c.; Fr. et Sav. l. c. p. 18.

NOM. JAP. *Kuromo*.

HAB. *Yezo*. Prov. Iburi : Lake Osatsu (Y. Tokubuchi!¹⁾ 1893).—Prov. Kushiro : Lake Akan (Faurie! n. 10722, 1893; T. Kawakami!²⁾ 1897; J. Hanzawa!³⁾ 1904); Lake Tōro (J. Hanzawa! 1904).—Prov. Kitami : Lake Abashiri (Faurie! 1899).

DISTRIB. *Yezo*, Honshū, Kiushū, Riukiu, Formosa, Korea, Amur-region, Manchuria, China, Tropical Asia, Central Europe, Mauritius, Madagaskar and Australia.

As to the distribution of the species in the Japanese Archipelago, the Lake Abashiri is probably its northernmost locality.

46. **Vallisneria spiralis** L. Spec. Pl. ed. 1. p. 1015; Fr. et Sav. Enum. Pl.

1) 徳淵永次郎。 2) 川上瀧彌。 3) 半澤 洵。

Jap. II. p. 18; Hook. f. Fl. Brit. Ind. V. p. 660; Aschers. u. Gürke, in Engl. Nat. Pfl.-fam. II. 1. p. 252. f. 185; Aschers. u. Graebn. Syn. Mitteleurop. Fl. I. p. 405; Britton and Brown, Ill. Fl. I. p. 93; Forb. et Hemsl. Ind. Fl. Sin. III. p. 2; Hegi, Ill. Fl. Mitteleurop. I. p. 161. f. 75; Matsum. Ind. Pl. Jap. II. 1. p. 33; Nakai, Fl. Kor. II. p. 217.

NOM. JAP. *Itomo*, *Sekishōmo*.

HAB. *Yezo*. Prov. Oshima: Ōnuma (Y. Tokubuchi! 1888); Junsainuma (K. Miyabe!¹⁾ 1899; S. Nishida!²⁾ 1913).—Prov. Ishikari: Lake Maoi (G. Yamada!³⁾ 1899).—Prov. Iburi: Lake Osatsu (Y. Tokubuchi! 1893).—Prov. Kushiro: Lake Tōro (S. Nozawa!⁴⁾ 1887); Lake Harutoru (K. Miyabe! 1894); Lake Akan (T. Kawakami! 1897; J. Hanzawa! 1904); Lake Kutcharo (K. Miyabe! 1915).

DISTRIB. *Yezo*, Honshū, Kiushū, Korea and widely distributed throughout Asia, Europe, Africa and North America.

47. *Calanthe nipponica* Makino, Bot. Mag. Tokyo XIII. (1899) p. 128; Matsum. Ind. Pl. Jap. II. 1. p. 237.

C. trulliformis King et Pantl. var. *hastata* A. Finet, Bull. Soc. Bot. de France XLVII. (1900) p. 267.

NOM. JAP. *Kinsei-ran*, *Kinsei-ebine*.

HAB. *Yezo*. Prov. Ishikari: Sapporo (T. Hasegawa!⁵⁾ 1882); Nopporo (K. Miyabe! 1914); Manju (F. C. Greatrix! 1915).—Prov. Kushiro: Kutcharo, Huptainoshiri (K. Miyabe! 1915); Mt. Oakan (T. Kawakami! 1897).

DISTRIB. *Yezo*, Honshū, Shikoku, Kiushū.

48. *Cymbidium virens* Lindl. Bot. Reg. (1853) p. 24; Miq. Prol. Fl. Jap. p. 137; Fr. et Sav. Enum. Pl. Jap. II. p. 27; Yabe, Bot. Mag. Tokyo XVII. p. 143; Matsum. Ind. Pl. Jap. II. 1. p. 241.

C. virescens Lindl. Bot. Mag. Misc. p. 37; Iinuma et Makino, Zōtei Somoku Zusetsu IV. pp. 1067 et 1186.—*C. Goeringii* Rchb. f. in Walp. Ann. III. p. 547.

NOM. JAP. *Shunran*, *Hokuro*.

HAB. *Yezo*. Prov. Shiribeshi: Isl. Okushiri (T. Kawakami! 1900; C. Suzuki!⁶⁾ 1914).

DISTRIB. Okushiri Island, Honshū, Shikoku, Kiushū.

49. *Dactylostalix ringens* Rchb. f. Bot. Zeit. XXXVI. (1878) p. 74;

1) 宮部金吾。 2) 西田彰三。 3) 山田玄太郎。 4) 野澤俊次郎。
5) 長谷川豊太郎。 6) 須崎忠助。

Pfitzer, in Engl. u. Prantl, Nat. Pfl.-fam. II. 6. p. 158.

Pergamena uniflora Finet, Bull. Soc. Bot. de France XLVII. (1900) p. 263 t. 8; Takeda, Hakubutsu-no-tomo VIII. n. 4. p. 16.

NOM. JAP. *Hime-hitoharan* (nov.).

HAB. *Yezo*. Prov. Ishikari : Mt. Nutakkanuishpe (A. Andō,¹⁾ 1907; Hideo Koizumi!²⁾ 1911).

DISTRIB. *Yezo*, Honshū.

50. *Dactylostalix maculosa* Miyabe et Kudo, sp. nov.

Dactylostalix ringens Makino, Bot. Mag. Tokyo XI. p. (413); Matsum. Ind. Pl. Jap. II. 1. p. 242? (non Rchb. f.)—*Pergamena uniflora* Makino, in Iinuma et Makino, Zōtei Somoku Zusetsu IV. pp. 1096 et 1210; Miyoshi and Makino, Atl. Alp. Pl. Jap. I. pl. 5. f. 28 (non Finet).

Rhizome horizontal, producing every year a tuber, a scape and a leaf, bearing a small bud at the end. Tuber almost perpendicular, slenderly elongate, tapering toward the end, villose throughout. Scape 16–20 cm in height, erect but ascending at the base, glabrous, usually with two cataphylls. Leaf one, subradical, petiolate, ovate or elliptical, 4–6 cm long, 3–4 cm broad, obtuse at the apex, round or obtuse at the base, membranaceous, 5–6 nerved on each side; petiole broadly winged, 1.2–2.5 cm long, 5–6 mm broad, many nerved. Flower solitary on the top of the scape. Outer perianth lineae-oblong, obtuse or acutish at the apex, 3-nerved, greenish in color, 18–20 mm long, 2.5 mm broad, the upper one a little broader and thicker. Inner perianth slightly shorter than the outer, linear-oblong, 3-nerved, acutish at the apex, greenish. Labellum shorter than the inner perianth, 14 mm long, obovate-orbicular, 3-lobed, whitish with purple dots; lateral lobes small, almost orbicular, tinged with a large dot; middle one largest, codate, round at the apex, a little crisped on the margin, 5-nerved, with 2 small dots at the base, 7 mm long, 10 mm broad. Gynostemium oblong, 1 cm long, 3.5 mm broad, 3-nerved, winged on the margin.

NOM. JAP. *Hitoharan*, *Ichiyōran*.

HAB. *Yezo*. Prov. Oshima : Moheji (S. Ninouye!³⁾ 1906 and 1907).

DISTRIB. Honshū and Southern part of Hokkaidō.

It is near to *Dactylostalix ringens* Rchb. f. from which it differs distinctly by its stouter stem and larger flowers, which have roundish labellum with conspicuous purple dots.

1) 安藤秋三郎。 2) 小泉秀雄。 3) 二ノ上七五三次郎。

51. **Liparis Krameri** Fr. et Sav. Enum. Pl. Jap. II. pp. 22 et 509 : Makino, Bot. Mag. Tokyo III. p. 7 ; Finet, Bull. Soc. Bot. de France XLVII. p. 262 ; Matsum. Ind. Pl. Jap. II. 1. p. 252 ; Nakai, Fl. Kor. II. p. 224.

NOM. JAP. *Jigabachisō*.

HAB. *Yezo*. Prov. Ishikari : Nopporo (K. Miyabe ! 1914).—Prov. Iburi : Mt. Eniwa (K. Miyabe and S. Arimoto ! ¹⁾ 1912 ; M. Majima ! ²⁾ 1906).

DISTRIB. *Yezo*, Honshū, Shikoku and Korea.

52. **Orchis cyclochila** Maxim. Mél. Biol. XII. (1886) p. 548 : Finet, Bull. Soc. Bot. de France XLVII. p. 267 : Kom. Fl. Mansh. I. p. 511 : Nakai, Fl. Kor. II. p. 223.

Habenaria cyclochila Fr. et Sav. Enum. Pl. Jap. II. p. 516.—*Gymnadenia spec.* Miq. Prol. Fl. Jap. p. 139.—*Gymnadenia cyclochila* Korsh. Acta Hort. Peterop. XII. p. 396 ; Makino Bot. Mag. Tokyo, XI. p. (415) ; Inuma et Makino, Zōtei Somoku Zusetsu IV. pp. 1097 et 1121 ; Matsum. Ind. Pl. Jap. II. 1. p. 249 ; Makino and Miyoshi, Atl. Alp. Pl. Jap. I. pl. 7. f. 41.

NOM. JAP. *Kamomesō*, *Kamome-ran*, *Ichiyō-chidori*.

HAB. *Yezo*. Prov. Kushiro : Shoro (K. Nishikawa ! ³⁾ 1914).

DISTRIB. *Yezo*, Honshū, Korea, Manchuria.

53. **Thesium repens** Ledeb. Fl. Alt. I. p. 274 et Fl. Ross. III. p. 539 ; DC. Prodr. XIV. p. 648 ; Fr. Schm. Fl. Sachal. p. 170 ; H. Takeda, Fl. Shikotan p. 484 ; Miyabe and Miyake, Fl. Saghal. p. 401.

NOM. JAP. *Kamayarisō*, *Hai-kanabikisō*.

HAB. *Yezo*. Prov. Hidaka : Mt. Apoi, Samani (K. Kondo ! ⁴⁾ 1912)

Kuriles. Isl. Shikotan : Shakotan (H. Takeda ! ⁵⁾ 1909).

Saghalin. Isl. Todomoshiri : Dainan (T. Miyake ! ⁶⁾ 1906) ; Kotan (T. Miyake ! 1906).—Kusunnai-Subdistrict : Kusunnai (Miyabe and Miyagi ! ⁷⁾ 1906).—Usutomanai-Subdistrict : Usutomanai (Miyabe and Miyagi ! 1906).—Toninai-Subdistrict : Chishinai (Miyake ! 1908).—Motodomari-Subdistrict : Mt. Nupuripo (Miyake ! 1907).

DISTRIB. *Yezo*, Southern Kuriles, Saghalin, Dahuria, Baical-region and Altai.

54. **Nuphar pumilum** Smith, Engl. Bot. t. 2292 ; DC. Prodr. I. p. 116 ;

1) 有元新太郎。 2) 眞島政吉。 3) 西川冠次郎。 4) 近藤金吾。
5) 武田久吉。 6) 三宅 勉。 7) 宮城鐵夫。

Ledeb. Fl. Alt. II. p. 274 et Fl. Ross. I. p. 85; Regel, Fl. Ussur. p. 18 et Pl. Radd. Polyp. p. 126; Korsh. Acta Hort. Peterop. XII. p. 304; Kawakami, Bot. Mag. Tokyo, XII. p. (224); Kom. Fl. Mansh. II. p. 219.

Nuphar japonicum DC. var. *subintegerrimum* Matsum Ind. Pl. Jap. II. 2. p. 92 (non Casp).

NOM. JAP. *Yezo-kōhone*, *Nemuro-kōhone*.

HAB. *Yezo*. Prov. Kushiro: Lake Akan (T. Kawakami! 1897).—Prov. Nemuro: Otsuishi (K. Miyabe! 1884); Tomoshiri (K. Miyabe! 1884; S. Yamane!¹⁾ 1894); Shibetsu (K. Miyabe! 1884).

Kuriles. Isl. Kunashiri: Tōbutsu-mura, Kenitaushi (C. Yendo!²⁾ 1894).—Isl. Etorofu: Toshimoi (T. Kawakami! 1898).

DISTRIB. *Yezo*, Manchuria, Amur-region, Siberia and Europe.

55. *Cardamine Fauriei* Franch. Bull. Soc. Phil. Paris (1888) p. 85; H. Boiss. in Bull. Herb. Boiss. VII. p. 792; Matsum. Ind. Pl. Jap. II. 2. p. 152.

C. geumifolia Koidz. Bot. Mag. Tokyo XXVII. p. (564).—*C. geifolia* Koidz. in Matsum. Ic. Pl. Koishi. II, p. 25. Pl. 97.

NOM. JAP. *Yezo-wasabi*, *Mitsuba-tanetsukebana*.

HAB. *Yezo*. Prov. Oshima: Ichinowatari near Shiriuchi (Miyabe and Tokubuchi! 1890).—Prov. Shiribeshi: Isl. Okushiri, Aonai (S. Yokoyama!³⁾ 1891); R. Yoichi at Nakanosawa (T. Ishikawa!⁴⁾ 1893); Inaotoge (Y. Takenobu and K. Mimashi!⁵⁾ 1883; Y. Tokubuchi! 1888); Kumaishi (Miyabe and Tokubuchi! 1890).—Prov. Ishikari: Asahikawa (Hideo Koizumi! 1913); Kamuikotan (K. Miyabe! 1891); Yūbari (Y. Tokubuchi! 1893); Utashinai (Y. Tokubuchi! 1892).—Prov. Teshio: Oketo (T. Ishikawa! 1892); Uyenbetsu (T. Ishikawa! 1891).—Prov. Iburi: Shitsukari (T. Kawakami! 1892).—Prov. Kitami: Nayoro (S. Yokoyama! 1892); Okoppe (S. Yokoyama! 1892); Sarubutsu (T. Ishikawa! 1892).

DISTRIB. *Yezo* and Honshū.

56. *Cardamine nasturtiifolia* Boiss. in Bull. Herb. Boiss. VII. (1899) p. 793; Matsum. Ind. Pl. Jap. II. 2. p. 153.

C. yezoensis Maxim. in litt.; Matsum. Bot. Mag. Tokyo IV. p. 11.—*C. Miyahei* Matsum. Ind. Pl. Jap. II. 2. p. 153.

NOM. JAP. *Yezo-no-janinjin*.

1) 山根 榮。 2) 遠藤千尋。 3) 横山壯次郎。 4) 石川貞治。
5) 武信由太郎。 三増久米吉。

HAB. *Yezo*. Prov. Hidaka : Saruru-sandō (K. Miyabe ! 1884 : Tokubuchi ! 1892); Saruru (Y. Tokubuchi ! 1892 : Faurie, 1893) : Shoya (Y. Tokubuchi ! 1892).

DISTRIB. An edemic species.

57. *Cardamine tenuifolia* Turcz. Fl. Baic.-Dahur. I. p. 120 : Maxim. Mém. Biol. IX. p. 4; Forb. et Hemsl. Ind. Fl. Sin. I. p. 44; Korsh. Acta Hort. Peterop. XII. p. 307; Palib. Fl. Kor. I. p. 27; Kom. Fl. Mansh. II. p. 362; Nakai, Fl. Kor. I. p. 56; Miyabe and Miyake, Fl. Saghal. p. 45.

Dentaria tenuifolia Ledeb. Fl. Alt. III. p. 43 et Fl. Ross. I. p. 130; DC. Prodr. I. 156 : Trautv. et Mey. Fl. Ochot. p. 16; Rgl. et Tiling, Fl. Ajan. p. 47; Maxim. Prim. Fl. Amur. p. 45; Fr. Schm. Fl. Sachal. p. 112.—*Dentaria tenella* Rgl. Fl. Ussur. p. 22, et Pl. Radd. Polyp. p. 176 (non Pursh.).—*Cardamine* Gmelin, Fl. Sib. III. p. 272. n. 41 t. 65.

NOM. JAP. *Hosoba-konronsō*.

HAB. *Yezo*. Prov. Kitami : Minami-sōya (Yanagisawa ! ¹⁾ 1911).

Saghalin. Mauka-Subdistrict : Mt. Mauka (T. Miyake ! 1907).—Tonnai-Subdistrict; Kimunai (T. Miyake ! 1908).—Russian Saghalin : Between Nuto and Chaiwo (T. Ishikawa ! 1912) : Dui and Arakai (Glehn 1861).

DISTRIB. Northern Yezo, Saghalin, Korea, Manchuria, Amur-region, Dahuria, and Siberia.

New to the Flora of Hokkaidō.

58. *Berchemia racemosa* Sieb. et Zucc. in Abh. Akad. Muench. iv. II. (1843) p. 147; A. Gray, Bot. Jap. p. 384; Benth Fl. Hongk. p. 67; Maxim. Rhamn. Or. p. 5; Miq. Prol. Fl. Jap. p. 219; Fr. et Sav. Enum. Pl. Jap. I. p. 81; Forb. et Hemsl. Ind. Fl. Sin. I. p. 127; Werberbauer Rhamn. in Engl. u. Prantl, Nat. Pfl.-fam. III. 5. p. 406 f. 199 A-C; Diels, Fl. Centr. China p. 458; Matsum. et Hayata, Enum. Pl. Formos. p. 87; Matsum. Ind. Pl. Jap. II. 2. p. 337.

NOM. JAP. *Kumayanagi*.

HAB. *Yezo*. Prov. Iburi : Tomakomai (Y. Kudō and T. Yoshimi ! ²⁾ 1915).

DISTRIB. Yezo, Honshū, Kiushū, Formosa, and China.

New to the Flora of Hokkaidō.

59. *Eritrichium nipponicum* Makino, Bot. Mag. Tokyo XVII. p. 51; Matsum. Ind. Pl. Jap. II. 2. p. 525; Miyabe and Miyake, Fl. Saghal. p. 334. pl. X. f.

1) 柳澤 某。 2) 工藤祐舜。吉見辰三郎。

3-6.

NOM. JAP. *Miyama-murasaki*.HAB. *Yezo*. Prov. Hidaka. : Mt. Apoi, Samani (K. Kondo ! 1912).*Saghalin*. Motodomari-Subdistrict : Chikaporonai (T. Miyake ! 1907) : Mt. Nupuripo (T. Miyake ! 1907).DISTRIB. *Saghalin*, *Yezo* and *Honshū*.

60. ***Lonicera strophophora*** Franch. in Bull. Soc. Philomath. Paris, Sér. 7. (1886) p. 142 ; A. Rehder, Syn. Lonic. in 14th An. Rep. of Mo. Bot. Gard. p. 95 ; Makino, Bot. Mag. Tokyo, XIX. p. 148 : C. K. Schn. Laubholzk. II. p. 703 ; Matsum. Ind. Pl. Jap. II. 2. p. 601.

Lonicera pilosa Maxim. Mém. Biol. X. p. 73 ; Fr. et Sav. Enum. Pl. Jap. II. p. 653 ; Kawakami, Dainippon Sanrinkwaihō. no. 219 (1901) p. 26, (non Willd.).—*Lonicera Amherstii* Dippel, Handb. Laubholzk. I. p. 263 f. 175 e. p. ; Makino, Bot. Mag. Tokyo XVII. p. 208.—*Caprifolium Amherstii* O. Kuntze, Rev. Gen. Pl. I. p. 274. e. p.

NOM. JAP. *Arage-hyotanboku*, *Ohba-hyotanboku*.HAB. *Yezo*. Prov. Shiribeshi : Isl. Okushiri (Miyabe and Tokubuchi ! 1890 ; T. Kawakami ! 1900).DISTRIB. Island of Okushiri and Northern *Honshū*.

摘 要

45. *Hydrilla verticillata* Royle. くろも。

膽振國長都湖、釧路國阿寒湖及塘路湖、北見國網走湖等に産す。

46. *Vallisneria spiralis* L. いとも、せきしやうも。

渡島、石狩、膽振、釧路等の湖沼中に生ず。

47. *Calanthe nipponica* Makino. きんせいらん、きんせいえびね。

石狩、釧路、兩國に産す。

48. *Cymbium virens* Lindl. しゆんらん、ほくろ。

本道に於ては奥尻島のみに産す。圖は増訂草木圖説第四輯千六十七頁にあり。

49. *Dactylostalix ringens* Rchb. f. ひめひとばらん。(新稱)

根莖は匍匐し先端に大形なる芽を生ず。一莖一葉一花。莖は細纖、上部に於て益細し、花も亦比較的小形なり。外花被片三個、膜質、三脈を有し、内花被片は二個、一乃至三脈を有し、膜質、唇瓣は三淺裂し、兩側裂片は極めて小形、中央裂片は大形にして卵狀長橢圓形又は長橢圓狀披針形、鈍頭、數脈を有し、膜質、斑点を欠く。藥柱は中央部甚だ厚く兩側は薄くして翼狀をなす。本種は本道に於てはヌタクカムシユベ山に産す。

50. *Dactylostalix maculosa* Miyabe et Kudo. ひとはらん、いちえふらん。

根莖は匍匐し先端に小形なる芽を生ず。一莖一葉一花。莖は比較的太くして上部に至るも甚だしく細形とならず。花は前者より大形なり。外花被片三個、少しく厚くして綠色を呈し、三脈を有す。内花被片二個、厚くして綠色を呈し、三脈を有す、唇瓣は大形、三淺裂し左右の裂片は小形、殆ど圓形、白色、内に二個の紫色の斑点を有す、中央裂片は大形、心臟形、圓頭、白色にして基部に二個の紫色の斑点を有す。藥柱は扁平、其縁部は膜質、三脈を有す。本種は本道にては渡島國上磯郡茂邊地村に産す。圖は増訂草木圖説第四輯千九十六頁及び高山植物圖譜第一卷第五圖版(二十八號)にあり。

51. *Liparis Kramerii* Fr. et Sav. じがばちさう。

石狩國及び膽振國に産す。

52. *Orchis cyclochila* Maxim. かもめさう、かもめらん、いちえふちどり。

釧路國に産す。圖は増訂草木圖説第四輯千〇九七頁にあり。

53. *Thesium repens* Ledeb. かまやりさう、はひかなびきさう。

北海道本島にては日高國アボイ山に産し、又千島にては支古丹島に産す又樺太にも生ず。

54. *Nuphar pumilum* Smith. えぞかはほね、ねむろかはほね。

釧路國及び根室國に産す。

55. *Cardamine Fauriei* Franch. えぞわさび、みつはたねつけはな。

渡島、後志、石狩、天鹽、北見等に於て通常に之れを産す。小泉氏の *Cardamine geifolia* と命ぜしは本種に外ならず。圖は植物圖編第二編第二集第九十七圖なり。

56. *Cardamine nasturtiifolia* Boiss. えぞのじやにんじん。

日高國、猿留及び庶野地方に限り本種を産す。

57. *Cardamine tenuifolia* Turcz. ほそばこんろんさう。

北見國南宗谷に産し又樺太に生ず。

58. *Berchemia racemosa* Sieb. et Zucc. くまやなぎ。

膽振國苫小牧、農科大學演習林内に自生す。

59. *Eritrichium nipponicum* Makino. みやまむらさき。

北海道に於ては日高國アポイ山に産し、又樺太にも之れを産す。圖は樺太植物誌第十圖版にあり。

60. *Lonicera strophophora* Franch. あらびへうたんぽく、おほへうたんぽく。

後志國奥尻島に産す。

