



HOKKAIDO UNIVERSITY

Title	Osteonecrosis and panniculitis as life-threatening signs
Author(s)	Kuwatani, Masaki; Kawakami, Hiroshi; Yamada, Yosuke
Citation	Clinical Gastroenterology and Hepatology, 8(5), e52-e53 https://doi.org/10.1016/j.cgh.2009.12.026
Issue Date	2010-05
Doc URL	https://hdl.handle.net/2115/61448
Rights	©2010. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	https://creativecommons.org/licenses/by-nc-nd/4.0/
Type	journal article
File Information	PPP_syndrome-CGH-revision2.pdf



Osteonecrosis and panniculitis as life-threatening signs

Masaki Kuwatani^{1,*}, Hiroshi Kawakami¹, Yosuke Yamada²

¹Department of Gastroenterology, Hokkaido University Graduate School of Medicine, North 15, West 7, Kita-ku, Sapporo 060-8638, Japan

²Department of Surgical pathology, Hokkaido University Hospital, North 15, West 7, Kita-ku, Sapporo 060-8638, Japan

*Address correspondence to: Dr. Masaki Kuwatani, Department of Gastroenterology, Hokkaido University Graduate School of Medicine, North 15, West 7, Kita-ku, Sapporo 060-8638, Japan

E-mail: mkuwatan@med.hokudai.ac.jp (M. Kuwatani)

Telephone: +81-11-716-1161(ext. 5920). FAX: +81-11-706-7867.

No conflicts of interest exist for all authors.

A 58-year-old male presented with polyarthritis in the bilateral wrists and ankles with elevated serum pancreatic enzymes (amylase, 4292 IU/l; lipase, 6000 IU/l). He had had several episodes of acute pancreatitis, and undergone cholangiojejunostomy for alcoholic chronic pancreatitis with a pseudocyst and stricture of the lower bile duct 6 years ago. Although pancreatic enzymes were temporarily normalized, roentgenograms exhibited osteonecrosis in four extremities one year after the development of polyarthritis (Figure A), and examination of a biopsy specimen of the left femur revealed intraosseous fat necrosis. CT just before admission to our department revealed slight dilatation of the main pancreatic duct and a cystic lesion adjacent to the superior mesenteric vein (SMV) in the pancreas head (Figure B). On admission, serological examinations indicated liver and renal dysfunction (elevated serum levels of AST [238 IU/l], ALT [167 IU/l], γ -GT [101 IU/l], BUN [49mg/dl], and creatinine [1.9mg/dl]) and elevated levels of pancreatic enzymes (amylase, 7966 IU/l; lipase, 12904 IU/l). Physical findings were erythema in the abdomen (Figure C) and purpura in bilateral lower extremities. Renal dysfunction rapidly exacerbated and he died 5 days after admission to our department. After informed consent from his family, autopsy was done. Gross findings of specimens of the pancreas head and the peripancreatic tissue showed a fistula between the cystic lesion in the pancreas head and SMV and exposure of debris from the cystic lesion. Microscopically, the cystic lesion was a pseudocyst connecting to the SMV, and the debris was protein plaque. The pancreas tissue surrounding the lower bile duct showed fibrous change and atrophy of the acini. In the kidneys, glomerular crescent formation was diffusely found, and a part of basement membrane was necrotic. In the skin, fibrinoid changes and infiltration of inflammatory cells were found in the walls of small arteries, and there was nodular fat necrosis with calcification (Figure D), which was also found in other organs including gastric subserosa and bone marrow. Finally, we made a diagnosis of PPP syndrome caused by pancreatic pseudocyst-portal vein fistula.

A case of pancreatitis with panniculitis and polyarthrits was reported in 1968 for the first time¹. PPP syndrome as terminology was suggested as a new entity in several reports, but it is a not well-known category of disease. This syndrome is frequently caused by alcoholic pancreatitis with a

pseudocyst and occasionally by pancreatic neoplasia². Although the mechanism remains to be determined, two possible mechanisms have been proposed: one is lysis of fat cells by high levels of pancreatic enzymes and the other is lysis of fat cells by autoimmune phenomena^{2, 3}. The main treatment for PPP syndrome is control of pancreatitis or neoplasm and drainage or preservative surgical intervention for a pseudocyst if it is present. However, the prognosis is poor, a review of the literature showing its mortality rates of 75-85%². Attention should be given to the possibility of PPP syndrome if polyarthrititis or osteonecrosis and panniculitis of unknown origin are detected. Delay in diagnosis of this syndrome can result in a negative chain reaction leading to death.

References

1. Mullin GT, Caperton EM, Crespin SR, et al. Arthritis and skin lesions resembling erythema nodosum in pancreatic disease. *Ann Intern Med* 1968;68:75–78
2. Carasso S, Oren I, Alroy G, et al. Disseminated fat necrosis with asymptomatic pancreatitis: a case report and review of the literature. *Am J Med Sci* 2000;319:68-72.
3. Mustafa KN, Hadidy A, Shoumaf M, Razzuki SA. Razzuki Polyarthritis with chondronecrosis associated with osteonecrosis, panniculitis and pancreatitis. *Rheumatol Int* 2009; epub

Figure A



Figure B




Figure C




Figure D

