




HOKKAIDO UNIVERSITY

Title	Virulence of tick-borne encephalitis virus is associated with intact conformational viral RNA structures in the variable region of the 3'-UTR
Author(s)	Sakai, Mizuki; Muto, Memi; Hirano, Minato et al.
Citation	Virus research, 203, 36-40 https://doi.org/10.1016/j.virusres.2015.03.006
Issue Date	2015-05-04
Doc URL	https://hdl.handle.net/2115/61469
Rights	© 2015, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International http://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	https://creativecommons.org/licenses/by-nc-nd/4.0/
Type	journal article
File Information	figure supplementary.pdf



Oshima-IC-pt

Supplementary Figure 1 (S1). RNA secondary structures of the 3'-UTR of viruses used in this study.
 The RNA secondary structures were predicted by MFOLD (<http://mfold.rna.albany.edu/?q=mfold/RNA-Folding-Form>).

Fig.S1

Sofjin-IC-pt




Fig.S1

Oshima dSL3/4/5




Fig.S1

Oshima dSL3/4




Fig.S1

Oshima dSL5




Fig.S1

Oshima_polyA

