



HOKKAIDO UNIVERSITY

Title	Studies of transcription factor FMBP-1 derived from silkworm Bombyx mori by using fluorescence correlation spectroscopy
Author(s)	堤, 元佐
Degree Grantor	北海道大学
Degree Name	博士(生命科学)
Dissertation Number	甲第12266号
Issue Date	2016-03-24
DOI	https://doi.org/10.14943/doctoral.k12266
Doc URL	https://hdl.handle.net/2115/61718
Type	doctoral thesis
File Information	Motosuke_Tsutsumi.pdf



Studies of transcription factor FMBP-1 derived from silkworm

Bombyx mori by using fluorescence correlation spectroscopy

(蛍光相関分光法を用いたカイコ *Bombyx mori* 由来転写因子 FMBP-1 の研究)

Laboratory of Protein Science

Graduate School of Life Science

Hokkaido University

2016

Motosuke Tsutsumi

Contents

Abbreviations	4
Abstract	5
Chapter 1: General Introduction	6
Chapter 2: <i>In vivo</i> fluorescence-correlation spectroscopy analyses of FMBP-1, a silkworm transcription factor	19
2-1 Introduction	20
2-2 Results	23
2-3 Discussion	47
2-4 Materials and Methods	58
2-5 References	66
2-6 Supporting Tables	69
Acknowledgements	79

Abbreviations

ACF	Autocorrelation function
AIC	Akaike's information criteria
AR	Acidic region
<i>B. mori</i>	<i>Bombyx mori</i>
EGFP	Enhanced green fluorescent protein
FCS	Fluorescence correlation spectroscopy
FMBP-1	Fibroin modulator binding protein 1
FRAP	Fluorescence recovery after photobleaching
HBR	Hyper basic region
LSM	Laser scanning microscopy
PSG	Posterior silk gland
SIM	Structured illumination microscopy
STPR	One score and three amino acid peptide repeat
TF	Transcription factor

Abstract

Fibroin modulator binding protein 1 (FMBP-1) is a silkworm transcription factor that has a unique DNA binding domain called the one score and three amino acid peptide repeat (STPR). Here I used fluorescence correlation spectroscopy (FCS) to analyze the diffusion properties of an enhanced green fluorescent protein-tagged FMBP-1 protein (EGFP-FMBP-1) expressed in posterior silk gland (PSG) cells of *Bombyx mori* at the same developmental stage as natural FMBP-1 expression. EGFP-FMBP-1 clearly localized to cell nuclei. From the FCS analyses, I identified an immobile DNA-bound component and three discernible diffusion components. I also used FCS to observe the movements of wild-type and mutant EGFP-FMBP-1 proteins in HeLa cells, a simpler experimental system. Based on previous *in vitro* observation, I also introduced a single amino acid substitution in order to suppress stable FMBP-1-DNA binding; specifically, I replaced the ninth Arg in the third repeat within the STPR domain with Ala. This mutation completely disrupted the slowest diffusion component as well as the immobile component. The diffusion properties of other FMBP-1 mutants (e.g., mutants with N-terminal or C-terminal truncations) were also analyzed. Based on my observations, I suggest that the four identifiable movements might correspond to four distinct FMBP-1 states: 1) diffusion of free protein, 2) and 3) two types of transient interactions between FMBP-1 and chromosomal DNA, and 4) stable binding of FMBP-1 to chromosomal DNA.


Chapter 1

General introduction

Transcription factors (TFs) are proteins which specifically bind to transcriptional regulatory region of DNA. Via the DNA binding, TFs enhance or repress transcription of genes into RNA. It is known that TFs widely play important roles for regulation of biofunctions: promotion of cell fate determination or cell differentiation (e.g. Hox TF family [1] and sex-determining region Y (SRY) TF [2]), response to intracellular signaling (e.g. glucocorticoid receptor [3] and estrogen receptor [4]), response to environmental stimuli (e.g. heat shock factor [5]), cell cycle control (e.g. p53 factor [6] and Myc protein [7]) and so on. Silk production of silkworm *Bombyx mori* is known as one of good example of such ingenious transcriptional regulation. The silk production is also well-known as example of large scale protein expression by organisms. The silk fiber consists of two proteins; fibroin and sericin. In the body of silkworm, silk glands which are the places of silk production are arranged in pairs. Fibroin proteins which are main components of silk fiber are composed of heavy (H) chain, light (L) chain and P25 units; and each unit is expressed in posterior silk gland (PSG) [8-13]. Then, fibroin proteins are stored in middle silk gland (MSG) and coated by sericin proteins which are expressed in there [11, 14-18]. Silk fibril are sent to orifice via anterior silk gland (ASG), and then the fibril and other silk fibril which are produced by another silk gland are combined into a silk fiber [19-21]. In this manner, the expression of silk proteins are proceeded as tissue-specific process. Also, it show developmental stage-specificity. Expression of each silk protein is activated in every feeding stage, but not during any molting stage [9-12, 16,

18, 20, 22]. It is considered that such tissue- and temporal-specific expression of the silk proteins are built on convoluted regulation by combination of TFs. Several fibroin TFs and sericin TFs have identified by previous works [23-30]. Recently, these TFs attract attention as important factors for control of biomaterial production. Genetically manipulated silkworm for industrial production of silk fiber which contains useful protein (e.g. fluorescent protein or spider silk fibroin) have been studied [31, 32]. In those studies, the silk protein gene TFs are used as key-factor in regulation of expression of target gene at appropriate timing and tissue. However, although detail DNA binding properties of several TFs had been reported, whole aspect of the mechanism of transcriptional regulation into silk production is unknown. Especially, detail molecular mechanism of DNA recognition had not been uncovered.

Fibroin modulator binding protein 1 (FMBP-1) is a factor identified by Takiya et al.; and it was reported as binding factor of fibroin H chain gene [20, 29]. The expression timing and tissue of FMBP-1 relatively more overlap with fibroin gene than other TFs [29]. Thus, FMBP-1 is considered as important transcriptional enhancer of fibroin gene. FMBP-1 include a unique structure which consists of four tandem repeats of homologous 23 amino acid residues (sequence identity between each repeat > 80%) (Fig. 1). This tandem repeat region is referred as one score and three amino acid peptide repeat (STPR) domain. And, properties of FMBP-1 and STPR domain were studied by using bioinformatic, biochemical and structural biological methods mainly in our laboratory. The bioinformatic analyses showed that STPR sequences are widely conserved in eukaryote (e.g. *Drosophila*, *Caenorhabditis elegans*, zebra fish, mouse, human and so on) [20]. Although most of silk protein TFs are known as ortholog of *Drosophila*



FMBP-1 (218 a. a.)

MGDLSQADSS WKKMVIAKTN NEWKSSVGED ANAASNTTHF RLSDECIQYD
 NIKKEIEEID EQETLNTVEP VDMIQEMDPL SLLEPKARRR RKGSGPKSET
 SEERAARLAK MSAYAAQRLA NESPEQRATR LKRMSEYAAK RLSSETREQR
 AIRLARMSAY AARRLANETP AQRQARLLRM SAYAAKRQAS KKSLSTVNDS
 LNYSIMPNQS RATNHPFT

Fig. 1. Schema and amino acid sequence of FMBP-1 protein.

Hox TFs [33], ortholog of FMBP-1 which except STPR domain was not found in *Drosophila* [20]. By biochemical experiments, the STPR domain was confirmed that the domain acts as a DNA binding domain of FMBP-1. Even STPR domain alone showed DNA binding ability in EMSA (gel shift assay) experiments [20]. The recognition sequence of FMBP-1 was reported as a 9 bp AT-rich motif (5'-ATNTWTNTA-3') [20]; FMBP-1 binds to the motif located in intronic transcriptional enhancer region "+290 element" of fibroin H-chain gene [29] (Fig. 2). The motif also exists in upstream region "-130 element" of fibroin H-chain gene; and, transcription of the fibroin H-chain gene is also enhanced by binding of FMBP-1 to "-130 element" [29]. In the EMSA experiment using DNA probe which include "+290 element" sequence, the STPR domain bound to the DNA probe with several nM of dissociation constant (K_d) [34, 35]. Previous biochemical studies also reported that the STPR domain binds to the major groove of specific DNA duplex, and the molecular ratio of the STPR domain against DNA duplex in the DNA bound state is 1:1 [34, 36]. From structural biological analyses by using circular dichroism (CD) and nuclear magnetic resonance (NMR), the structure of each repeat of STPR domain was uncovered. It is interesting that N-terminal half of each repeat forms α -helix, while C-terminal half of each repeat of STPR domain is intrinsically disordered; the α -helix of each repeat extends to C-terminal half and become rigid structure in DNA bound state [34, 35]. Those structural analyses also suggested that the extended α -helix structure are stabilized by a salt bridge between first Glu and ninth Arg of each repeat of STPR domain. Actually, the point mutant of STPR domain which was substituted Ala for ninth Arg of third repeat of STPR domain (R9A(rep3) mutant) showed clearly lower DNA binding ability than wild-type STPR domain [34, 35].




Fig. 2. Schema of DNA recognition of FMBP-1.

As described above, various analyses for STPR domain had been performed. However, the number of analyses using full-length FMBP-1 were relatively low because of difficulty of sample preparation and instability in structure of N-terminal half of FMBP-1. Therefore, K_d value of DNA binding of FMBP-1 had not been reported yet. The 3D structure of full-length FMBP-1 is unknown. Also, the properties of DNA interaction of STPR domain and full-length FMBP-1 *in vivo* had not been studied so far. To further consideration of the role of FMBP-1 in native state, detailed observation of movement of FMBP-1 in *in situ* and *in vivo* condition would be crucial.

Not limited to FMBP-1, observation of kinetics of TF in DNA interaction has become big topic. Development of fluorescence observation method enable us to analyze the kinetics of biomolecular with high spatial- and high temporal-resolution under various conditions including *in vitro*, *in vivo* and *in situ*. Recently, many analyses of DNA binding proteins by using fluorescence observation methods with single molecule sensitivity had been performed: e.g. fluorescence recovery after photobleaching (FRAP) analyses of estrogen receptor α (ER α) [37], total internal reflection fluorescence (TIRF) observation of Herpes simplex virus polymerase UL42 and DNA repair protein MutS α [38, 39], fluorescence correlation spectroscopy (FCS) analyses of retinoic acid receptor (RAR) and glucocorticoid receptor (GR) [40, 41] and so on. Results of these observation indicated that the DNA recognition of DNA binding protein are more complicated and dynamic process. Conventional image of DNA recognition process of DNA binding protein was composed of free diffusion state and DNA bound state. However, those results of recent studies reported that more steps are included in the DNA recognition: e.g.

binding of TF with co-transcription factor or transient DNA interactions before stable DNA binding which referred as "facilitated diffusion". The existence of facilitated diffusion movements was predicted and suggested via consideration of the results of biophysical studies since 1980s [42, 43]. The results of modern fluorescence observation studies as above mentioned could identify the movements correspond to facilitated diffusion. Based on the experimental results, computer simulation studies classified some types of facilitated diffusion: "DNA hopping" which is a 3D diffusion movement on DNA chain via transient interaction mainly caused by electrostatic interaction, "DNA sliding" which is a sliding movement along several or several-ten base-pair of DNA phosphate backbone, "intersegmental transfer" which is a jumping from one DNA segment to another DNA segment, and so on (Fig. 3).

In the modern fluorescence observation methods, FCS is a suitable method for analysis of transcription kinetics in live cell. FCS techniques use the fluorescence fluctuation caused by Brownian motion of fluorescently tagged particles in a prolate spheroid microscopic space that is referred to as the confocal volume. Then, an autocorrelation curve is mathematically extracted from measured fluorescence fluctuation; this curve is compared to each of several diffusion models by curve fitting to determine the number and component ratio of diffusion components, and diffusion time of each diffusion component [44-46]. FCS method has several merits: 1) FCS measurement can be used for various condition including live cell, 2) it can analyze the diffusion properties at pinpoint because of very small size of measurement area (the diameter is some hundreds nm), 3) it also can simultaneously determine the detailed diffusion properties of multicomponent, 4) the temporal resolution of FCS




Fig. 3. Representative movements of kinetics of transcription factor (TF) in DNA interaction.

is very higher (it can identify μsec order diffusion movement) than other fluorescence observation method. Utilizing these merits, FCS method had been actually used for many observation of prime proteins in live cells [47, 48]. In recent years, FCS analyses have been applied not only to cultured cells, but also to bio organ or invertebrate animals' cells [49-58]. Particularly, the FCS analyses of Hox TF in *Drosophila* salivary gland which was performed by Vukojević et al. [54] is very interesting as an instance of applicability of FCS method to insect organ. It is known that the silk gland of *B. mori* is homologous to *Drosophila* salivary gland. Successful results of FCS analyses of TF in the salivary gland suggest that I would be able to analyze the diffusion properties of FMBP-1 in silk gland by using FCS method.

Based on these background, in this article, I will mainly report the FCS analyses of FMBP-1 in live cells. First, I tried to express EGFP tagged FMBP-1 (EGFP-FMBP-1) in PSG cells of *B. mori* five-instar larvae at the same developmental stage and place as natural FMBP-1 expression. Then, the kinetics of EGFP-FMBP-1 was analyzed by FCS. Next, the kinetics of R9A(rep3) mutant was compared to wild-type EGFP-FMBP-1. HeLa cells which transiently expressed various EGFP-FMBP-1 mutants were also used for FCS analyses as more simple experimental system than PSG cells. Their diffusion properties were analyzed in detail. Further, FRAP experiments and observation by using super-resolution microscopy were performed to determine the interaction target of FMBP-1. Those results were carefully interpreted based on various supposition. Finally, the process of DNA interaction of FMBP-1 and the role of each region in FMBP-1 were proposed.

References

- 1) Pearson JC, Lemons D & McGinnis W (2005) Modulating Hox gene functions during animal body patterning. *Nat Rev Genet* **6**, 893-904.
- 2) Page DC, Mosher R, Simpson EM, Fisher EMC, Mardon G, Pollack J, McGillivray B, Chapelle A & Brown LG (1987) The sex-determining region of the human Y chromosome encodes a finger protein. *Cell* **51**, 1091-1104.
- 3) Rhen T & Cidlowski JA (2005) Antiinflammatory action of glucocorticoids--new mechanisms for old drugs. *N Engl J Med* **353**, 1711-1723.
- 4) Dahlman-Wright K, Cavailles V, Fuqua SA, Jordan VC, Katzenellenbogen JA, Korach KS, Maggi A, Muramatsu M, Parker MG & Gustafsson JA (2006) International Union of Pharmacology. LXIV. Estrogen receptors. *Pharmacol Rev* **58**, 773-781.
- 5) Sorger PK (1991) Heat shock factor and the heat shock response. *Cell* **65**, 363-366.
- 6) Surget S, Khoury MP & Bourdon JC (2013) Uncovering the role of p53 splice variants in human malignancy: a clinical perspective. *Onco Targets Ther* **7**, 57-68.
- 7) Varmus HE (1984) The molecular genetics of cellular oncogenes. *Ann Rev Genet* **18**, 553-612.
- 8) Suzuki Y & Brown D (1972) Isolation and identification of the messenger RNA for silk fibroin from *Bombyx mori*. *J Mol Biol* **63**, 409-429.
- 9) Suzuki Y & Suzuki E (1974) Quantitative measurements of fibroin messenger RNA synthesis in the posterior silk gland of normal and mutant *Bombyx mori*. *J Mol Biol* **88**, 393-407.
- 10) Maekawa H & Suzuki Y (1980) Repeated turn-off and turn-on of fibroin gene transcription during silk gland development of *Bombyx mori*. *Dev Biol* **78**, 394-406.
- 11) Ishikawa E & Suzuki Y (1985) Tissue- and stage-specific expression of sericin genes in the middle silk gland of *Bombyx mori*. *Dev Growth Differ* **27**, 73-82.
- 12) Couble P, Moine A, Garel A & Prudhomme JC (1983) Developmental variations of a nonfibroin mRNA of *Bombyx mori* silk gland, encoding for a low-molecular-weight silk protein. *Dev Biol* **97**, 398-407.
- 13) Kimura K, Oyama F, Ueda H, Mizuno S & Shimura K (1985) Molecular cloning of the fibroin light chain complementary DNA and its use in the study of the expression of the light chain gene in the posterior silk gland of *Bombyx mori*. *Experientia* **41**, 1167-1171.
- 14) Okamoto H, Ishikawa E & Suzuki Y (1982) Structural analysis of sericin genes. Homologies with fibroin gene in the 50 flanking nucleotide sequences. *J Biol Chem* **257**, 15192-15199.
- 15) Couble P, Michaille JJ, Garel A, Couble ML & Prudhomme JJ (1987) Developmental switches of sericin mRNA splicing in individual cells of *Bombyx mori* silkgland. *Dev Biol* **124**, 431-440.
- 16) Obara T & Suzuki Y (1988) Temporal and spatial control of silk gene transcription analyzed by nuclear run-on assays. *Dev Biol* **127**, 384-391.
- 17) Michaille JJ, Garel A & Couble P (1990) Cloning and characterization of the highly polymorphic Ser2 gene of *Bombyx mori*. *Gene* **86**, 177-184.
- 18) Matsunami K, Kokubo H, Ohno K & Suzuki Y (1998) Expression pattern analysis of SGF-3/POU-M1 in relation to sericin-1 gene expression in the silk gland. *Dev Growth Differ* **40**, 591-597.
- 19) Komatsu K (1975) Kinu no kozo to seishi (The structure of silk and silk reeling). *Proceeding of 28th Summer school of silk reeling*, 1-31.
- 20) Takiya S, Ishikawa T, Ohtsuka K, Nishita Y & Suzuki Y (2005) Fibroin-modulator-binding protein-1 (FMBP-1) contains a novel DNA-binding domain, repeats of the score and three amino acid peptide (STP), conserved from *Caenorhabditis elegans* to humans. *Nucleic Acids Res* **33**, 786-795.
- 21) The International Silkworm Genome Consortium (2008) The genome of a lepidopteran

- model insect, the silkworm *Bombyx mori*. *Insect Biochemistry and Molecular Biology* **38**, 1036-1045.
- 22) Suzuki Y & Giza PE (1976) Accentuated expression of silk fibroin gene *in vivo* and *in vitro*. *J Mol Biol* **107**, 183–206.
 - 23) Durand B, Drevet J & Couble P (1992) P25 gene regulation in *Bombyx mori* silk gland: Two promoter-binding factors have distinct tissue and developmental specificities. *Mol Cell Biol* **12**, 5768-5777.
 - 24) Horard B, Julien E, Nony P, Garel A & Couble P (1997) Differential binding of the *Bombyx* silk gland-specific factor SGFB to its target DNA sequence drives posterior-cell-restricted expression. *Mol Cell Biol* **17**, 1572-1579.
 - 25) Hui CC & Suzuki Y (1989) Enhancement of transcription from the Ad2 major late promoter by upstream elements of the fibroin- and sericin-1-encoding genes in silk gland extracts. *Gene* **85**, 403-411.
 - 26) Hui CC, Matsuno K & Suzuki Y (1990) Fibroin gene promoter contains a cluster of homeodomain binding sites that interact with three silk gland factors. *J Mol Biol* **213**, 651-670.
 - 27) Mach V, Takiya S, Ohno K, Handa H, Imai T & Suzuki Y (1995) Silk gland factor-1 involved in the regulation of *Bombyx* sericin-1 gene contains fork head motif. *J Biol Chem* **270**, 9340-9346.
 - 28) Fukuta M, Matsuno K, Hui CC, Nagata T, Takiya S, Xu PX, Ueno K & Suzuki Y (1993) Molecular cloning of a POU domain-containing factor involved in the regulation of the *Bombyx* sericin-1 gene. *J Biol Chem* **268**, 19471-19475.
 - 29) Takiya S, Kokubo H & Suzuki Y (1997) Transcriptional regulatory elements in the upstream and intron of the fibroin gene bind three specific factors POU-M1, Bm Fkh and FMBP-1. *Biochem J* **321**, 645-653.
 - 30) Zhao XM, Liu C, Li QY, Hu WB, Zhou MT, Nie HY, Zhang YX, Peng ZC, Zhao P & Xia QY (2014) Basic helix-loop-helix transcription factor Bmsage is involved in regulation of fibroin H-chain Gene via interaction with SGF1 in *Bombyx mori*. *PLoS One* **9**, e94091.
 - 31) Kuwana Y, Sezutsu H, Nakajima K, Tamada Y & Kojima K (2014) High-toughness silk produced by a transgenic silkworm expressing spider (*Araneus ventricosus*) dragline silk protein. *PLoS One*, **9**(8), e105325.
 - 32) Iizuka T, Sezutsu H, Tatematsu KI, Kobayashi I, Yonemura N, Uchino K, Nakajima K, Kojima K, Takabayashi C, Machii H, Yamada K, Kurihara H, Asakura T, Nakazawa Y, Miyawaki A, Karasawa S, Kobayashi H, Yamaguchi J, Kuwabara N, Nakamura T, Yoshii K & Tamura T (2013) Colored fluorescent silk made by transgenic silkworms. *Adv Funct Mater* **23**(42):5232-5239.
 - 33) Kimoto M, Tsubota T, Uchino K, Sezutsu H & Takiya S (2014) Hox transcription factor Antp regulates sericin-1 gene expression in the terminal differentiated silk gland of *Bombyx mori*. *Dev Biol* **386**, 64-71.
 - 34) Saito S, Yokoyama T, Aizawa T, Kawaguchi K, Yamaki T, Matsumoto D, Kamijima T, Kamiya M, Kumaki Y, Mizuguchi M, Takiya S, Demura M & Kawano K (2008) Structural properties of the DNA-bound form of a novel tandem repeat DNA binding domain, STPR. *Proteins* **72**, 414-426.
 - 35) Nonaka Y, Muto H, Aizawa T, Okabe E, Myoba S, Yokoyama T, Saito S, Tatami F, Kumaki Y, Kamiya M, Kikukawa T, Mizuguchi M, Takiya S, Kinjo M, Demura M & Kawano K (2010) STPR, a 23-amino acid tandem repeat domain, found in the human function-unknown protein ZNF821. *Biochemistry* **49**, 8367-8375.
 - 36) Takiya S, Saito S, Yokoyama T, Matsumoto D, Aizawa T, Kamiya M, Demura M & Kawano K (2009) DNA-binding property of the novel DNA-binding domain STPR in FMBP-1 of the silkworm *Bombyx mori*. *J Biochem* **146**, 103-111.
 - 37) Jankevics H, Prummer M, Izewska P, Pick H, Leufgen K & Vogel H (2005) Diffusion-time

- distribution analysis reveals characteristic ligand-dependent interaction patterns of nuclear receptors in living cells. *Biochemistry* **44**, 11676-11683.
- 38) Komazin-Meredith G, Mirchev R, Golan DE, van Oijen AM & Coen DM (2008) Hopping of a processivity factor on DNA revealed by single-molecule assays of diffusion. *Proc Natl Acad Sci USA* **105**, 10721-10726.
 - 39) Gorman J, Wang F, Redding S, Plys AJ, Fazio T, Wind S, Alani EE & Greene EC (2012) Single-molecule imaging reveals target-search mechanisms during DNA mismatch repair. *Proc Natl Acad Sci USA* **109**, E3074.
 - 40) Brazda P, Szekeres T, Bravics B, Toth K, Vamosi G & Nagy L (2011) Live-cell fluorescence correlation spectroscopy dissects the role of coregulator exchange and chromatin binding in retinoic acid receptor mobility. *J Cell Sci* **124**, 3631-3642.
 - 41) Mikuni S, Tamura M & Kinjo M (2007) Analysis of intranuclear binding process of glucocorticoid receptor using fluorescence correlation spectroscopy. *FEBS Lett* **581**(3), 389-393.
 - 42) Winter RB, Berg OG & von Hippel OH (1981) Diffusion-driven mechanisms of protein translocation on nucleic acids. 3. The *Escherichia coli* lac repressor-operator interaction: kinetic measurements and conclusions. *Biochemistry* **20**, 6961-6977.
 - 43) Hannon R, Richards EG & Gould HJ (1986) Facilitated diffusion of a DNA binding protein on chromatin. *EMBO J* **5**, 3313-3319.
 - 44) Brock R, Vamosi G, Vereb G & Jovin TM (1999) Rapid characterization of green fluorescent protein fusion proteins on the molecular and cellular level by fluorescence correlation microscopy. *Proc Natl Acad Sci USA* **96**, 10123-10128.
 - 45) Lippincott-Schwartz J, Snapp E & Kenworthy A (2001) Studying protein dynamics in living cells. *Nat Rev Mol Cell Biol* **2**, 444-456.
 - 46) Berland KM (2004) Fluorescence correlation spectroscopy: a new tool for quantification of molecular interactions. *Methods Mol Biol* **261**, 383-398.
 - 47) Giorgio G, Yinghua G, Jagesh VS & Galit L (2013) Activation and control of p53 tetramerization in individual living cells. *Proc Natl Acad Sci USA* **110**, 15497-15501.
 - 48) Tiwari M, Mikuni S, Muto H & Kinjo M (2013) Determination of dissociation constant of the NFκB p50/p65 heterodimer using fluorescence cross-correlation spectroscopy in the living cell. *Biochem Biophys Res Commun* **436**, 430-435.
 - 49) Pan X, Yu H, Shi X, Korzh V & Wohland T (2007) Characterization of flow direction in microchannels and zebrafish blood vessels by scanning fluorescence correlation spectroscopy. *J Biomed Opt* **12**(1), 014034-1-10.
 - 50) Funke SA, Birkmann E, Henke F, Görtz P, Lange-Asschenfeldt C, Riesner D & Willbold D (2007) Single particle detection of Abeta aggregates associated with Alzheimer's disease. *Biochem Biophys Res Commun* **364**(4), 902-907.
 - 51) Berg RH & Beachy RN (2008) Fluorescent protein applications in plants. *Methods Cell Biol* **85**, 153-77.
 - 52) Kappler J, Hegener O, Baader SL, Franken S, Gieselmann V, Häberlein H & Rauch U (2009) Transport of a hyaluronan-binding protein in brain tissue. *Matrix Biol* **28**(7), 396-405.
 - 53) Yu SR, Burkhardt M, Nowak M, Ries J, Petrásek Z, Scholpp S, Schwille P & Brand M (2009) Fgf8 morphogen gradient forms by a source-sink mechanism with freely diffusing molecules. *nature* **461**, 533-536.
 - 54) Vukojević V, Papadopoulos DK, Terenius L, Gehring WJ & Rigler R (2009) Quantitative study of synthetic Hox transcription factor-DNA interactions in live cells. *Proc Natl Acad Sci USA* **107**, 4093-4098.
 - 55) Lee JI, Sato M, Ushida K & Mochida J (2011) Measurement of diffusion in articular cartilage using fluorescence correlation spectroscopy. *BMC Biotechnol* **2**, 11-19.
 - 56) Gehring WJ (2011) How do Hox transcription factors find their target genes in the nucleus of living cells? *Biol Aujourd'hui* **205**(2), 75-85.

- 57) Guldbrand S, Kirejev V, Simonsson C, Goksör M, Smedh M & Ericson MB (2013) Two-photon fluorescence correlation spectroscopy as a tool for measuring molecular diffusion within human skin. *Eur J Pharm Biopharm* **84**(2), 430-436.
- 58) Yaehe K, Tekrony A, Clancy A, Gregoriou Y, Walker J, Dean K, Nguyen T, Doiron A, Rinker K, Jiang XY, Childs S & Cramb D (2013) Nanoparticle accumulation in angiogenic tissues: Towards predictable pharmacokinetics *Small* **9**, 3118-3127.

Chapter 2

In vivo fluorescence-correlation spectroscopy analyses of FMBP-1, a silkworm transcription factor

2-1. Introduction

Silk glands of silkworm *Bombyx mori* larvae are among the most famous protein-producing organs in scientific literature. When cocoon formation begins, the gross weight of these silk glands accounts for nearly 40% of larval weight, and these organs store huge amounts of silk protein. The larval silk gland of *B. mori* is distinctly divided into the anterior silk gland (ASG), the middle silk gland (MSG), and the posterior silk gland (PSG) based on structural and functional criteria. A silk gland comprises about 600 cells, and each cell grows without cell division. Thus, each cell becomes very large; moreover, each nucleus becomes polyploid and develops a dendritic form [1, 2].

Fibroin is the main component of silk protein, and it consists of a heavy (H) chain, light (L) chain, and P25. Throughout larval development, fibroin is expressed in the PSG during every feeding stage, but not during any molting stage [2]. Such tissue- and temporal-specific expression of the fibroin protein is considered to be precisely controlled by a number of transcription factors (TFs). Several fibroin TFs such as BMFA [3], SGFB [3, 4], Fkh/SGF-1 [5-7], SGF-2 [6], POU-M1/SGF-3 [6, 8, 9], Bmsage [10], and FMBP-1 (fibroin modulator binding protein-1) [9, 11] have been identified. However, a comprehensive picture of fibroin gene expression is lacking.

FMBP-1 is a recently identified TF that specifically binds to a 9 bp AT-rich motif (5'-ATNTWTNTA-3') in upstream and intronic promoter elements of the gene encoding the fibroin H chain [9]. FMBP-1 comprises 218 amino acid residues and is divided into several distinctive domains on the basis of amino acid sequence. Notably, the C-terminal half has a unique structure that comprises four tandem repeats of a

23-residue domain, known as the one score and three amino acid peptide repeat (STPR) domain, which acts as a DNA-binding domain in FMBP-1 [11]. Various properties of FMBP-1 have been determined via biochemical or structural biological techniques [12-14]. For example, analyses involving nuclear magnetic resonance, circular dichroism, and limited digestion have shown that the STPR domain adopts a quite rigid, helix-rich structure when bound to DNA, but forms a flexible structure in the absence of DNA [13]. Mutational analysis of the STPR domain demonstrated that each salt bridge between the fourth glutamic acid residue and the ninth arginine residue of each repeat is important to the rigid structure adopted by FMBP-1 in the DNA-bound state. However, the dynamics between FMBP-1 and DNA *in vivo* remain unknown.

Here, I used fluorescence correlation spectroscopy (FCS) techniques to assess these dynamics *in vivo*. FCS is a powerful method used to observe many kinds of molecular dynamics and interactions in aqueous conditions with single-molecule sensitivity [15-17]. FCS measures fluorescence intensity fluctuation caused by Brownian motion of fluorescent molecules in a small volume element (a femtoliter-volume in the shape of a prolate spheroid) that is generated in the sample through the confocal arrangement of optical elements. Then, an autocorrelation curve is mathematically extracted from the measured fluorescence fluctuation; this temporal autocorrelation curve is then compared to autocorrelation functions of several diffusion models by curve fitting. Comparisons between the extracted functions and the models are used to determine 1) the diffusion time, 2) the number of diffusion components, and 3) the relative abundance of each diffusion component. Recently, FCS techniques have been used to study several TFs in cultured cells [18-20]. Furthermore, FCS-based quantitative techniques have been used to analyze homeodomain-DNA interactions in *Drosophila* salivary gland cells [21].

These studies and findings indicate that FCS analysis could be used to study FMBP-1 dynamics in silk gland cells.

In this study, I utilized FCS to analyze the diffusion dynamics of FMBP-1 in the PSGs of *B. mori* fifth instar larvae, which represent the tissue and larval stage positive for endogenous FMBP-1.

2-2. Results

2-2-1. Laser-scanning microscopy (LSM) observations of FMBP-1 in PSG cells

To mimic *in vivo* conditions as closely as possible while observing FMBP-1 mobility, I used transfected cells from PSGs of fifth instar *B. mori* larvae, which transiently express a fusion protein (EGFP-FMBP-1) comprising an EGFP tag and full-length FMBP-1 sequence. As controls, PSG cells that expressed EGFP alone were prepared in parallel. The PSG is a tubular tissue that constitutes the posterior half of the silk gland, and is the site of fibroin protein production (Fig. 1A). A confocal image of a Hoechst-stained EGFP-expressing PSG is shown at low magnification in Fig. 1B. The alternating and consecutive arrangement of the semi-cylindrical cells forms the lumen of the PSG (Fig. 1C). In PSGs that expressed EGFP alone, fluorescence was observed throughout the cell and was not exclusively localized to the cell nuclei (Fig. 1D). In contrast, PSGs that expressed EGFP-FMBP-1 showed clear nuclear localization of the EGFP signal (Fig. 1E).

2-2-2. FCS analyses of EGFP signals in PSG cells

Next, I measured the fluorescence fluctuations of the EGFP-FMBP-1 signal in the nuclei of PSG cells. The fluorescence intensity of EGFP-FMBP-1 showed obvious decay during the first few tens of seconds of each measurement (Fig. 2A). In contrast, the fluorescence intensity of EGFP alone remained almost constant. The photobleaching of EGFP-FMBP-1 likely resulted from the immobilization of a fraction of




Fig. 1. Localization of EGFP-FMBP-1 in the posterior silk gland cells. (A) The structure of a silk gland dissected from a fifth instar *Bombyx mori* larvae. The scale bar is 5 mm. The silk gland is distinctly divided into an anterior silk gland (ASG), a middle silk gland (MSG), and a posterior silk gland (PSG). Isolated PSGs were used for all FCS measurements. (B) Confocal overview image of PSG cells. The scale bar is 500 μ m. Each cell nucleus was stained with

Hoechst33342 (red), and only a few cells expressed the transgene (green). The PSG is a tubular organ consisting of alternately facing cells. Each cell has a dendritic polyploid nucleus. (C) Schematic image of the typical structure of PSG cells. This image was drawn based on the image in (B). (D) Confocal images of a PSG cell transiently expressing EGFP alone. The scale bar is 50µm. (E) Confocal image of a PSG cell transiently expressing wild-type EGFP-FMBP-1. The scale bar is 50µm. These confocal images were taken with Carl Zeiss LSM510 microscope (EGFP fluorescence, Ex. 488 nm, Em. 505-540 nm. Hoechst 33342 [nuclear stain] fluorescence, Ex. 405 nm, Em. 420-480 nm).




Fig. 2. Fluorescence fluctuation from EGFP-FMBP-1 in PSG cells. Sequential record data of fluorescence intensity of (A) wild-type FMBP-1 and (B) R9A(rep3) mutant (red). The intensity of EGFP alone with similar expression level was used as a control (green). Black curves lying on each fluctuation record indicate the fitting curve of photobleaching drawn by the fitting equation (2) described in the *Materials and Methods*.

EGFP-FMBP-1 molecules due to DNA binding, whereas for EGFP alone, free diffusion ensured that the resulting signal did not decay as rapidly.

For correct FCS analysis, the observable photobleaching time period of EGFP-FMBP-1 was removed from the ensemble calculation to extract the autocorrelation function. Then, each of three free diffusion models (one-, two- and three-component) were used to fit the autocorrelation curves (Fig. 3A-C). The residuals showed the systematic deviations for each model. One- and two-component fits had larger χ^2 values than did the three-component fits. Akaike's information criteria (AIC) and *F*-tests were also used to compare the probabilities associated with the respective models; both analyses strongly indicated that the three-component model most closely fit the data (Tables S1 and S2). Furthermore, inclusion of a fourth diffusion component did not improve the fit, nor did it result in reasonable diffusion parameters. In addition, three anomalous-diffusion models (a one-component and two-component models with one anomalous and one free component or with both components anomalous) were also tested. The residuals of the curve fitting with the one-component anomalous-diffusion model were significantly larger than those of the three-component free diffusion model. However, the residuals of the curve fitting with the other two anomalous-diffusion models were slightly smaller than that of three-component free diffusion model. Nevertheless, the preciseness of the fitting parameters was also an important factor in choosing the best-fitting model [22]. However, the diffusion parameters that were determined by both two-component anomalous diffusion models varied greatly (maximum standard deviation of the parameters were larger than 200%); consequently, the interpretation of each diffusion component was very difficult (Table S3). For reference, a three-component anomalous diffusion model was also tested for




Fig. 3. FCS analyses of EGFP-FMBP-1 in the PSG cells. (A) Autocorrelation function of wild-type FMBP-1 FCS (navy blue line) with various fitting models: the one-component free diffusion model (green dot line), the two-component free diffusion model (blue dot line), and the three-component free diffusion model (red line). A graph of the residuals is shown below the images. (B)&(C) Enlarged views of latter half of (A). Fitting curves of the two-component, free diffusion model and three-component, free diffusion model are shown in (B) and (C), respectively. (D) Box plot of diffusion coefficient of EGFP alone (green), three diffusion coefficients of wild-type FMBP-1 (WT, red) and two diffusion coefficients of R9A(rep3) mutant (orange). The whiskers show the minimum and maximum value of each coefficient. These diffusion coefficients were converted from diffusion times, which were measured via FCS, the conversion formula (4) described in the *Materials and Methods* was used for all conversions.

fit (Table S3). The diffusion parameters obtained by the three-component anomalous diffusion model also varied greatly. Therefore, the autocorrelation curves of EGFP-FMBP-1 were accounted for by the three-component free diffusion model. The EGFP-FMBP-1 diffusion parameters measured in the PSG cells are shown in Table 1.

The diffusion times determined from the FCS measurements indicated the transit times of fluorescent particles in a confocal volume (here, approximately 400 nm in diameter and 2000 nm in height). The diffusion coefficient of the 1st component of EGFP-FMBP-1 was similar to that of EGFP alone (Fig. 3D). In contrast, the diffusion coefficients of the 2nd and 3rd components of EGFP-FMBP-1 were significantly smaller than those of EGFP alone. These results indicated that EGFP-FMBP-1 molecules exhibited three different diffusion modes and one immobile (photobleaching) state *in vivo*.

2-2-3. FCS analyses of EGFP-FMBP-1 mutants in PSG cells

Previous mutational analyses had shown that substitution of the ninth arginine with an alanine in repeat 3 of the STPR domain (R9A(rep3)) most clearly disrupted the rigid helix-rich domain conformation and reduced the ability of FMBP-1 to stably bind DNA [13]. To further characterize the four putative components (three diffusion components and one immobile component) of FMBP-1 dynamics, an R9A(rep3) EGFP-FMBP-1 mutant was compared to wild-type EGFP-FMBP-1 in PSG cells.

The R9A(rep3) mutant, like wild-type FMBP-1, localized to the nucleus of each PSG cell (Fig. 4A). However, the mutant did not exhibit photobleaching during FCS measurements (Fig. 2B). In addition, the two-component diffusion model was appropriate for curve fitting of the autocorrelation curve of the R9A(rep3) mutant (Fig. 5);

Table 1. Diffusion parameters of EGFP-FMBP-1 in PSG cells.

	1st component		2nd component		3rd component		Bleach	<i>n</i>
	F_1 (%)	τ_1 (μ s)	F_2 (%)	τ_2 (ms)	F_3 (%)	τ_3 (ms)		
EGFP alone	100	266.7 $\pm$ 53.4					-	13
wild-type	45.1 $\pm$ 14.0	343.2 $\pm$ 59.8	41.4 $\pm$ 7.6	3.28 $\pm$ 1.69	13.5 $\pm$ 9.5	98.2 $\pm$ 88.3	+	29
R9A(rep3)	76.9 $\pm$ 4.0	339.8 $\pm$ 48.7	23.1 $\pm$ 4.0	5.99 $\pm$ 2.99			-	16

Note: These values were derived by curve fitting of the autocorrelation function via the optimum fitting model for each sample. The “ $\pm$ ” in each column indicates standard deviation (SD). The “+” and “-” in the “Bleach” column indicate the presence or absence, respectively, of observable photobleaching.




Fig. 4. Nuclear localization of each EGFP-FMBP-1 mutant. (A) Confocal image of a PSG cell transiently expressing R9A(rep3) mutant. The image was taken with a Carl Zeiss LSM510 microscope (EGFP fluorescence, Ex. 488 nm, Em. 505-540 nm. Hoechst 33342 (nuclear stain) fluorescence, Ex. 405 nm, Em. 420-480 nm). The scale bar is 50 μm. (B)-(J) Confocal images of

HeLa cells expressing (B) EGFP alone, (C) wild-type FMBP-1, (D) R9A(rep3), (E) Δ Ctail, (F) Δ C-rep4, (G) Δ C-STPR, (H) Δ C-HBR, (I) Δ N-HBR, or (J) Δ N-AR mutant were taken with a Carl Zeiss LSM510 microscope (EGFP fluorescence, Ex. 488 nm, Em. 505-540 nm). The scale bar is 10 μ m. (K) Measurement points for FCS analyses in a PSG cell expressing wild-type FMBP-1. The image was taken in the same manner as the images in (A). The scale bar is 20 μ m. (L) Measurement point for FCS analyses in a HeLa cell expressing wild-type FMBP-1. The image was taken in the same manner as image (B)-(J). The scale bar is 5 μ m. In the image (K) and (L), cross marks show the measurement points for FCS, and arrowheads indicate the nucleoli.




Fig. 5. Curve fitting for various mutants of EGFP-FMBP-1. Representative ACFs of various mutants of EGFP-FMBP-1 (navy blue line) with three fitting models: the one-component, free diffusion model (green dot line); the two-component, free diffusion model (blue dot line); and the three-component, free diffusion model (red dot line). A graph of the residuals is shown below each ACF graph. In this study, the number of diffusion components was judged by fitting

residuals and appropriateness of diffusion parameters, which were calculated by the curve fitting. As a result, the three-component model was found to be appropriate for the wild-type FMBP-1 (in both PSG cells and HeLa cells), R9A(rep1), Δ Ctail, Δ N-HBR and Δ N-AR mutants. The two-component model was found to be appropriate for the R9A(rep2), R9A(rep3) (in both PSG cells and HeLa cells), R9A(rep4), Δ C-rep4 and Δ C-STPR mutants. The one-component model was found to be appropriate for EGFP alone (in both PSG cells and HeLa cells) and the Δ C-HBR mutant.

here, inclusion of a third diffusion component did not improve the fit. The results of the fitting analysis showed such a low amplitude that no reliable diffusion time could be determined. Normalized autocorrelation curves derived from all measured data from wild-type FMBP-1 or the R9A mutant were compared. As the R9A(rep3) mutant and wild-type FMBP-1 proteins differed by only one amino acid residue, the two proteins should have similar molecular weights; however, their autocorrelation curves clearly differed (Fig. 6). Thus, this change in the autocorrelation curves should represent changes in the interactions between FMBP-1 and other molecules in the nuclei. The diffusion parameters for the R9A(rep3) mutant measured in the PSG cells are shown in Table 1. Each diffusion time obtained from curve fitting corresponded to the 1st or 2nd component of the wild-type FMBP-1 (Table 1 and Fig. 3D). In other words, the R9A(rep3) mutant simultaneously lost both the 3rd diffusion component and the immobile component. Taken together, this result and previous findings indicated that these two components were dependent on the stable binding of FMBP-1 to DNA.

2-2-4. Observation of wild-type EGFP-FMBP-1 in HeLa cells

To further examine each diffusion component and the role of each domain of FMBP-1 in more detail, it was necessary to observe various FMBP-1 mutants. However, observing such mutants in PSG cells was difficult because sample preparation using the tungsten gun method was complicated. Additionally, measurement success rates were low; in particular, low transformation efficiency with PSG cells caused a decrease of measurement success rates. Furthermore, the strong intrinsic fluorescence of PSG cells and the instability of the fluorescence intensity of EGFP in these cells made it difficult to efficiently measure diffusion properties. Therefore, I analyzed the diffusion of




Fig. 6. Autocorrelation curve of EGFP-FMBP-1. Autocorrelation curves based on measurements from PSG cells (solid lines) or HeLa cells (dotted lines). Each autocorrelation curve represents the average of all measurements taken from each sample; the curves were normalized to have the same amplitude.

EGFP-tagged FMBP-1 mutants in HeLa cells, which represent a simpler experimental system. HeLa cells were also used in a previous analysis of the ZNF821 protein (an ortholog of the STPR domain in humans) that was performed by the authors in our laboratory [23].

First, I determined whether each diffusion movement of wild-type EGFP-FMBP-1 observed in PSG cells could be reproduced in HeLa cells. EGFP-FMBP-1 showed clear nuclear localization in confocal images of HeLa cells, as it had in PSG cells (Fig. 4B, C). The EGFP-FMBP-1 measured in HeLa cell nuclei also showed photobleaching during the early stage of each measurement (Fig. 7A). Both normalized autocorrelation curves of EGFP alone and of EGFP-FMBP-1 showed shapes similar to those seen with PSG cell measurements (Fig. 6). The three-component diffusion model was also appropriate for the curve fitting of data from EGFP-FMBP-1 in HeLa cell nuclei (Fig. 5). The diffusion times of wild-type FMBP-1 are listed in Table 2. The range of diffusion times were very similar to those observed in the PSG cells (Fig. 8). These results confirmed that the diffusion movements of EGFP-FMBP-1 in HeLa cells accurately reproduced those in PSG cells.

2-2-5. FCS analyses of EGFP-FMBP-1 mutants in HeLa cells

Next, I constructed mutant versions of EGFP-FMBP-1 and transfected HeLa cells with each mutant separately (Fig. 9). Substitution mutants similar to the R9A(rep3) mutant were constructed such that Arg9 of each STPR repeat was individually replaced with Ala to generate three additional mutants: R9A(rep1), R9A(rep2), and R9A(rep4). I also generated and analyzed four C-terminal truncation mutants and two N-terminal




Fig. 7. Fluorescence fluctuation of EGFP-FMBP-1 mutants. Sequential record data of fluorescence intensity with (A) wild-type EGFP-FMBP-1 or (B) various FMBP-1 mutants in HeLa cells (red). The initial intensity record data of EGFP alone (green) was compared as the control, as in Figure 2. (C) Comparison between Δ N-AR (red) and wild-type EGFP-FMBP-1 (orange) with regard to photobleaching ratio. The initial intensity levels of both fluctuations were almost identical. Black curves lying on each fluctuation record indicate the fitting curve of photobleaching drawn by the fitting equation (2) described in the *Materials and Methods*.

Table 2. Diffusion parameters of various EGFP-FMBP-1 mutants in HeLa cells.

	1st component		2nd component		3rd component		Bleach	<i>n</i>
	F_1 (%)	τ_1 (μ s)	F_2 (%)	τ_2 (ms)	F_3 (%)	τ_3 (ms)		
EGFP alone	100	278.2 $\pm$ 19.5					-	15
wild-type	33.9 $\pm$ 7.7	335.1 $\pm$ 52.8	52.8 $\pm$ 5.4	2.43 $\pm$ 0.67	13.3 $\pm$ 6.9	112.1 $\pm$ 79.6	+	27
R9A(rep1)	36.9 $\pm$ 4.8	355.8 $\pm$ 42.8	59.3 $\pm$ 4.5	2.51 $\pm$ 0.78	3.8 $\pm$ 1.8	209.1 $\pm$ 203.3	+	8
R9A(rep2)	44.2 $\pm$ 5.2	355.8 $\pm$ 37.0	55.8 $\pm$ 5.2	3.47 $\pm$ 0.51			-	12
R9A(rep3)	40.9 $\pm$ 6.8	354.8 $\pm$ 85.2	59.1 $\pm$ 6.8	3.20 $\pm$ 1.30			-	15
R9A(rep4)	45.0 $\pm$ 7.3	390.6 $\pm$ 85.9	55.0 $\pm$ 7.3	4.21 $\pm$ 0.86			-	7
Δ Ctail	33.0 $\pm$ 4.3	330.9 $\pm$ 45.7	55.2 $\pm$ 6.4	2.67 $\pm$ 0.81	11.8 $\pm$ 4.9	77.3 $\pm$ 44.3	+	18
Δ C-rep4	50.4 $\pm$ 11.5	357.7 $\pm$ 77.0	49.6 $\pm$ 11.5	2.85 $\pm$ 1.19			-	14
Δ C-STPR	80.3 $\pm$ 13.6	326.2 $\pm$ 64.5	19.7 $\pm$ 13.6	1.70 $\pm$ 0.52			-	10
Δ C-HBR	100	345.8 $\pm$ 36.8					-	21
Δ N-HBR	33.5 $\pm$ 4.4	271.7 $\pm$ 36.9	57.4 $\pm$ 3.7	1.65 $\pm$ 0.42	9.1 $\pm$ 4.5	33.2 $\pm$ 14.0	+	7
Δ N-AR	36.9 $\pm$ 3.9	292.7 $\pm$ 41.5	45.5 $\pm$ 5.4	2.29 $\pm$ 0.47	17.6 $\pm$ 5.6	60.6 $\pm$ 33.6	+	10

Note: These values were derived by curve fitting of the autocorrelation function with the optimum fitting model for each sample. The “ $\pm$ ” in each column indicate SD. The “+” and “-” in the “Bleach” column indicate the presence or absence, respectively, of observable photobleaching.




Fig. 8. Diffusion coefficient of wild-type EGFP-FMBP-1 in HeLa cells. Box plot of the diffusion coefficient of EGFP alone (green) and three diffusion coefficients for wild-type EGFP-FMBP-1 (WT, red). The whiskers show the minimum and maximum value of each coefficient. These diffusion coefficients were converted from the diffusion times determined via FCS analyses and calculated via the conversion formula (4) (shown in *Materials and Methods*).




Fig. 9. Representation of mutant forms of EGFP-FMBP-1. Schematic representation of EGFP-FMBP-1 molecules used in this study. The enhanced green fluorescent protein (EGFP) sequences were fused in-frame to the 5' end of the FMBP-1 sequences. Substitution mutants were constructed such that Arg9 of each STPR repeat was separately replaced with Ala to generate the R9A(rep1), R9A(rep2), R9A(rep3), and R9A(rep4) mutants. The C-terminal truncation mutants were as follows: ΔCtail mutant, truncated C-terminal tail; ΔC-rep4 mutant, truncated C-tail and fourth repeat of STPR domain; ΔC-STPR mutant, truncated C-tail and all repeats of STPR domain; and ΔC-HBR mutant, truncated from C-tail to hyper basic region (HBR). Each was used for FCS measurements. The N-terminal truncation mutants were as follows: ΔN-HBR mutant, truncated N-terminal region from N-terminus to HBR and ΔN-AR mutant, truncated from N-terminus to acidic region (AR).

truncation mutants: 1) Δ Ctail, with a truncated C-terminal tail; 2) Δ C-rep4, with a truncated C-tail and fourth repeat of the STPR domain; 3) Δ C-STPR, with a truncated C-tail, and all repeats of the STPR domain; 4) Δ C-HBR, truncated from the C-tail to the hyper basic region (HBR); 5) Δ N-HBR, with a truncated N-terminal region from the N-terminus to HBR; and 6) Δ N-AR, with a truncated from the N-terminus to the acidic region (AR) (Fig. 9).

The mobility of each FMBP-1 R9A mutant was observed in HeLa cells. Each R9A mutant localized in the nuclei, just as the R9A(rep3) mutant had in the PSG cells (Fig. 4D). Of the four R9A mutants, three: R9A(rep2), R9A(rep3), and R9A(rep4), did not show photobleaching (Fig. 7B). The position and shape of the autocorrelation curves of the R9A(rep3) mutant in HeLa cells were similar to those of the R9A(rep3) mutant in PSG cells (Fig. 6). The two-component diffusion model was best for the curve fitting of the autocorrelation curves for these three R9A mutants (Fig. 5). The diffusion movements of the R9A(rep3) mutant in HeLa cells also closely reproduced those of the R9A(rep3) mutant in PSG cells. Additionally, the R9A(rep2) and R9A(rep4) mutants each showed results similar to those of the R9A(rep3) mutant. In contrast, the fluorescence intensity of the R9A(rep1) mutant decayed slightly in the first several seconds of each measurement, and the FCS data were best accounted for by the three-component diffusion model (Fig. 5 and Table S4). Such differences among the R9A mutants were similar to those of previously published *in vitro* findings with respect to their DNA interactions [12]. Thus, these findings suggested that each repeat of the STPR domain does not contribute equally to the DNA-EGFP-FMBP-1 interactions and that the repeats might have differing roles.

The Δ Ctail, Δ C-rep4, Δ N-HBR, and Δ N-AR mutants each showed nuclear

localization (Fig. 4E, F, I, J, respectively). Deletion of the entire STPR domain (Δ C-STPR) resulted in reduced nuclear localization (Fig. 4G), and further deletion of HBR (Δ C-HBR) eliminated nuclear localization completely (Fig. 4H). The Δ C-rep4, Δ C-STPR, and Δ C-HBR mutants did not show photobleaching (Fig. 7B). In contrast, the decay of fluorescence intensity of three mutants: Δ Ctail, Δ N-HBR, and Δ N-AR, was observed at an early stage of each FCS measurement, as was that of wild-type FMBP-1. For these photobleaching truncation mutants, the fluorescence fluctuation data from the observable photobleaching time period were removed from the ensemble average that was used for calculation of the autocorrelation curves. For curve fitting of the autocorrelation curves of the truncation mutants, the three-component diffusion model was appropriate for each of the three mutants (Fig. 5). However, each of the other mutants: Δ C-rep4, Δ C-STPR, and Δ C-HBR, did not show the 3rd component. Furthermore, the Δ C-HBR mutant lost even the 2nd component. These diffusion parameters are shown in Table 2. The diffusion time of the 1st and 2nd components of each truncation mutant were within the same range as those for the wild-type FMBP-1. The component ratios between the immobile component and the three diffusion components of the respective FMBP-1 mutants are summarized in Fig. 10. Notably, the ratio of the 2nd component for the mutant lacking the entire STPR (Δ C-STPR) was decreased. Further deletion of HBR (Δ C-HBR) clearly abolished the 2nd component. On the other hand, the Δ N-HBR mutant, which had only the STPR domain and the C-tail region, showed the 2nd component. These results indicated that the HBR and STPR domains each contributed to a fraction of the 2nd component and that the degree of these contributions differed.




Fig. 10. Component ratios among diffusion movements of EGFP-FMBP-1 mutants. The 100% stacked bar chart of average component ratios for each diffusion movement. This chart includes the immobile component ratio, which represents the decay rate of fluorescence intensity in photobleaching. The other three components (1st, 2nd, and 3rd) were distributed within the remaining component ratio, according to the component distribution F_i ($i = 1, 2, 3$) of diffusion parameters (obtained by FCS analyses, shown in Table 1 and 2).




Fig. 11. Fluorescence recovery after photobleaching (FRAP) of EGFP-FMBP-1 in HeLa cells. (A) Fluorescence recovery of wild-type EGFP-FMBP-1 (red), R9A(rep3) mutant (blue) and EGFP alone (green) in HeLa cell nuclei. Averaged curves for wild-type EGFP-FMBP-1 ($n = 10$), the R9A(rep3) mutant ($n = 5$), and EGFP alone ($n = 5$) are shown. The error bars show standard deviation. (B)&(C) Averaged t_{half} values of fluorescence recovery and fluorescence recovery rates, respectively, of wild-type EGFP-FMBP-1 and mutants that show slow fluorescence recovery ($n \geq 10$).

2-2-6. Fluorescence recovery after photobleaching (FRAP) analyses of wild-type and mutant EGFP-FMBP-1 in HeLa cells

To characterize the immobile component of EGFP-FMBP-1 in more detail, I performed FRAP experiments in HeLa cells. The fluorescence recovery of wild-type EGFP-FMBP-1 was clearly delayed compared to that of EGFP alone (Fig. 11A). The fluorescence recovery curves were fitted and interpreted using an exponential association model. The results of wild-type EGFP-FMBP-1 showed that the fluorescence was completely recovered with $t_{half} \approx 20$ sec (Fig. 11B). For EGFP alone, I could not determine an accurate fluorescence recovery time in this experiment because the recovery was too rapid to analyze the recovery curve. In the nuclei of HeLa cells expressing the R9A(rep1), Δ Ctail, Δ N-HBR, and Δ N-AR mutants, a slow fluorescence recovery ($t_{half} \approx 5\text{--}15$ sec) was also observed (Fig. 11B). In contrast, for HeLa cells expressing other mutants, the fluorescence recovery times could not be determined for the same reason as for EGFP alone. The fluorescence recovery rate of each EGFP-FMBP-1 mutant that showed slow fluorescence recovery also approached 100% (Fig. 11C). Notably, the EGFP-FMBP-1 mutants that showed slow fluorescence recovery in the FRAP experiments completely corresponded to the mutants exhibiting photobleaching during FCS analyses.

2-3. Discussion

My results indicated that STPR and HBR acted cooperatively to function as a nuclear localization signal (Fig. 4). Previous findings on human ZNF821, which contains a STPR motif in the C-terminal tail, also indicated that STPR contributes to nuclear localization [23]. Thus, the nuclear localization function of the STPR motif might be widely conserved among STPR-containing proteins.

Although FCS analyses were carried out under low laser intensity, the EGFP-tagged wild-type FMBP-1 showed observable photobleaching in both PSG and HeLa cells (Fig. 2A and Fig. 7A). The photobleaching was also confirmed and analyzed in detail in HeLa cells using FRAP (Fig. 11). Previous studies suggest that FMBP-1 binds to a specific recognition site consisting of a 9-bp sequence in the vicinity of the gene encoding the fibroin-H chain, where it cooperates with other TFs to promote transcription [11]. However, the length of this recognition sequence is very short in comparison with the genome size of *B. mori* (4.32×10^8 bp, as the DNA content in a PSG cell polyploid nucleus becomes a thousand times larger than the original) and of humans (3.15×10^9 bp). Thus, a rough estimation based on the expected frequency of this consensus sequence (every 8,192 bp in the *B. mori* genome) indicated that multiple FMBP-1 recognition sequences should be present throughout chromosomal DNA in both PSG and HeLa cells. Furthermore, previous electron microscopy findings showed that there are many euchromatic regions that can interact with TFs in the PSG cell nuclei of fifth-instar larvae [2]. Based on these estimations, I proposed that the photobleaching observed in both PSG and HeLa cells nuclei reflected the stable binding of FMBP-1 to its recognition sequence throughout chromosomal DNA (Fig. 2A and Fig.

7A). The disappearance of photobleaching caused by deletion or point mutation of the STPR domain suggested that STPR acted as the DNA binding domain of FMBP-1 *in vivo*, as had been determined using previous *in vitro* assays (Fig. 2B and Fig. 7B). To support the proposed interaction of EGFP-FMBP-1 with chromatin in HeLa cells, colocalization of EGFP-FMBP-1 and mCherry tagged histone H2B protein (H2B-mCherry) was observed using confocal imaging of HeLa cells co-expressing wild-type EGFP-FMBP-1 and H2B-mCherry, wherein the distributions of both proteins overlapped (Fig. 12A). Super-resolution imaging with structured illumination microscopy (SIM) and high-resolution confocal imaging were also performed for detailed observation of the colocalization and to provide evidence of direct interaction, respectively. In the super-resolution imaging of HeLa cells expressing wild-type EGFP-FMBP-1, colocalization of EGFP fluorescence with H2B-mCherry was clearly detected (Fig. 12B). In contrast, the EGFP fluorescence of the R9A(rep3) mutants did not completely co-localize with H2B-mCherry in HeLa cell nuclei. Together, these results confirm the physical and localized interaction of wild-type EGFP-FMBP-1 with chromatin. Thus, the photobleaching observed in the FCS analyses can likely be attributed to the stable binding of FMBP-1 to chromatin. FRAP analysis of the stability of this interaction in HeLa cells (Fig. 11) determined that the t_{half} of the binding of wild-type EGFP-FMBP-1 to chromatin was 20 sec.

In the FCS analyses, FMBP-1 showed three distinct diffusion components in both PSG and HeLa cells based on curve fitting of the autocorrelation functions. Although the photobleaching period (the first 20 sec of each measurement in PSG cells and the first 10 sec in HeLa cells) was removed from the fluorescence fluctuation before curve fitting to determine the number of diffusion components (Fig. 3), the presence of 3rd





Fig. 12. Colocalization of wild-type EGFP-FMBP-1 and chromatin in HeLa cells. (A) Confocal images of HeLa cells in which EGFP, wild-type EGFP-FMBP-1, or the R9A(rep3) mutant were co-expressed with histone H2B-mCherry. The images were taken with a Carl Zeiss LSM510 microscope (EGFP fluorescence, Ex. 488 nm, Em. 505-530 nm; mCherry fluorescence, Ex. 594 nm, Em. LP650 nm). The scale bars represent 5 μ m. The fluorescence intensity profile across the nucleus (shown as a white broken arrow) of each image is also shown for each fluorophore (right). (B) SIM images and high-resolution confocal images of HeLa cells in which wild-type EGFP-FMBP-1, or the R9A(rep3) mutant were co-expressed with histone H2B-mCherry. Those proteins were immunofluorescence stained by AlexaFluor 488 (for EGFP-FMBP-1), or AlexaFluor 594 (for H2B-mCherry). The images were taken with a Nikon N-SIM or Leica TCS SP8 microscope (AlexaFluor 488 fluorescence, Ex. 488 nm, Em. 504-595 nm; AlexaFluor 594 fluorescence, Ex. 595 nm, Em. 603-653 nm). The scale bars represent 5 μ m. Expanded image of yellow frame region in each merge image is shown on the right. Arrow heads in expanded images indicate the colocalization spots of EGFP-FMBP-1 and H2B-mCherry.

component movement was coincident with the presence of photobleaching (Fig. 10). To check the possibility that the 3rd component was a remnant of photobleaching, I compared the autocorrelation curves that were derived from each 10 sec time interval of fluorescence fluctuation in each FCS measurement (consisting of 6 sequential 10 sec of record data) (Fig. 13). The positions and shapes of the autocorrelation curves derived from the first 0–10 sec of fluorescence fluctuation after the measurement start were clearly different from those of the autocorrelation curves derived from other time intervals. On the other hand, the autocorrelation curves derived from the five time intervals that represented later periods during data collection resembled each other. In addition, I determined the diffusion parameters by curve fittings of the autocorrelation curves derived from 20–30 or separately from 50–60 sec of fluorescence fluctuations (Table S5). Generally, photobleaching can be accounted for by an exponential model [24]. Thus, the influence of photobleaching would be expected to be weakest in the last 10-sec time interval. In my results, the component ratios and diffusion times of the 3rd diffusion component derived from 50–60 sec did not decrease from those of the 20–30 sec data, indicating that the 3rd component was not a remnant of photobleaching, but rather an independent component.

Because TFs generally bind to or interact with various molecules, one TF might show multiple distinct diffusion movements. Previous observations of TFs using FCS or FRAP have shown kinetic changes of the same TF via dimerization and ligand binding [18, 25, 26]. Here, I found that the diffusion times of the 1st, 2nd, and 3rd component of each FMBP-1 mutant were similar to those of the wild-type FMBP-1: τ_1 = few hundred microseconds, τ_2 = a few milliseconds, and τ_3 = several tens of milliseconds or more, respectively (Table 1 and 2); nevertheless, the component ratios clearly differed among




Fig. 13. Autocorrelation curves derived from each 10-sec time interval of fluorescence fluctuation in a single FCS measurement. Each autocorrelation curve was calculated from each 10-sec time interval (0-10, 10-20, 20-30, 30-40, 40-50 and 50-60 sec) of fluorescence fluctuation data (sequential 10 sec $\times$ 6 times record) in a single measurement of wild-type FMBP-1 in PSG cell. In the FCS analysis, first two time intervals were removed from the ensemble average of autocorrelation curve for curve fitting to determine the diffusion parameters.

the FMBP-1 variants (Fig. 10). I therefore propose that these three components might reflect FMBP-1 status during three different classes of movements and molecular interactions.

Based on the diffusion times measured via FCS, I assessed the apparent masses of each diffusing component. These calculations were based on the assumption that each diffusion component had a spherical shape, and previous work has stated that these calculations allow estimation of the order of the mass of the respective components [18]. The apparent molecular masses of the diffusion components of FMBP-1 measured in the PSG cells were calculated by comparing their diffusion times with that of EGFP alone (27 kDa). The apparent mass for the 1st component was approximately 58 kDa, nearly that of real monomeric EGFP-FMBP-1, 51.2 kDa. However, the apparent masses for both the 2nd component and the 3rd component were substantially larger than that of actual monomeric FMBP-1 (approximately 50,000 kDa = 50 MDa and 1,300,000,000 kDa = 1.3 TDa, respectively). The ratio between the apparent molecular masses of the 2nd component and the real monomeric wild-type FMBP-1 molecule was approximately 1,000-fold. The apparent mass was also much larger than that of the pre-initiation complex in transcription, which is approximately 3 MDa, or that of a mammalian ribosome, which is the largest intracellular organelle-like complex and has a molecular mass of 4.6 MDa. Furthermore, for the 3rd component, the ratio between the apparent molecular masses and that of the actual monomeric wild-type FMBP-1 molecule was as high as 2.5×10^7 -fold. It is unlikely that such a large diffusing protein complex exists in PSG cells. Thus, it is reasonable to interpret this slow diffusion not as representing the formation of extremely large molecular complexes, but as steric hindrance or transient interaction with another large and immobile structure in the cell

nucleus. Actually, the delay of apparent diffusion times because of interactions with immobile structures has been proposed based on a previous simulation study involving FCS [27]. Here, I did not observe any localization of EGFP-FMBP-1 on nuclear membranes or nucleoli in the confocal images; subsequently, those two nuclear regions were intentionally avoided during all FCS measurements (Fig. 4K, L). I also did not observe any aggregation or speckle formation in the nucleoplasm (Fig. 1E and Fig. 4C). On the other hand, colocalization of the wild-type EGFP-FMBP-1 with chromatin was confirmed by co-expression with H2B-mCherry (Fig. 12). Thus, a probable candidate for such a large immobile structure that could interact with FMBP-1 in the nucleus is chromosomal DNA. In short, these results strongly indicated that the slow diffusion movements (both the 2nd and 3rd components) of FMBP-1 also reflected interactions, albeit transient, between FMBP-1 and chromosomal DNA.

Recent studies of the interaction dynamics between DNA and TFs or DNA repair proteins using single-molecule detection or molecular-dynamics simulation have indicated that very short-term events occur during protein-DNA interactions; such events are called "facilitated diffusion" and include DNA sliding, DNA hopping, and DNA intersegment transfers. Importantly, facilitated diffusion has been suggested to cause proteins to reach a recognition sequence on DNA faster than would free, three-dimensional diffusion [28-36]. Notably, the diffusion coefficient of the 3rd component of wild-type FMBP-1 in the PSG cells in this study, $D = 0.19 \pm 0.20 \mu\text{m}^2/\text{sec}$ (range 0.021–0.812 $\mu\text{m}^2/\text{sec}$) was comparable to that of DNA sliding as determined via single-molecule detection (0.01–0.5 $\mu\text{m}^2/\text{sec}$) [31, 33, 34, 36]. Additionally, the presence of 3rd component movement was coincident with the presence of photobleaching, which was considered to represent stable DNA binding with

chromosomal DNA for all mutants (Fig. 10). Therefore, this result could be considered to support a model wherein the 3rd component movement represents facilitated diffusion that would trigger stable DNA binding. Previous findings have suggested that the DNA binding domains of TFs or DNA repair proteins might enter the groove of a DNA helix during the DNA sliding state; conformational changes in 3D structure are important for such interactions [32, 35]. Findings from previous *in vitro* DNA-binding experiments involving STPR domains indicate that a conformational change into the α -helical-rich structure is important for stable DNA binding and that the R9A mutation abolishes this conformational change [13]. Thus, this conformational change might play an important role during 3rd component movements of the STPR as well as for stable DNA binding.

On the other hand, the diffusion coefficient of the 2nd component ($D = 3.62 \pm 2.07 \mu\text{m}^2/\text{sec}$) was 19-fold larger than that of the reported DNA sliding motion [31, 33, 34, 36]. DNA hopping, which is another type of facilitated diffusion, is also reportedly more rapid than is DNA sliding [34-36]. Here, I focused on the changes in the component ratios caused by mutation. In this study, every mutant that retained the STPR domain or HBR retained the 2nd component (Fig. 10). The isoelectric points calculated based on the amino acid sequences of HBR (pI = 12.5) or STPR (pI = 11.4) are very high, thus these positively charged regions could transiently adsorb onto the negatively charged DNA because of the electrostatic interactions between them. In fact, previous findings indicate that DNA hopping results from such electrostatic interactions and that the diffusion movement becomes faster with increasing salt concentrations [32, 35]. It is likely that the 2nd component movements of FMBP-1 therefore reflected the transient trapping of FMBP-1 on chromosomal DNA (like DNA hopping) via electrostatic

interactions in which both the STPR domain and HBR play important roles.

This study represents the first demonstration of FCS analyses in the silkworm silk gland *in vivo*. From this, I detected three types of interactions between EGFP-FMBP-1 and DNA using *in vivo* FCS and mathematical analyses involving a three-component diffusion model. I note, however, that because of the complicated environment in living cells, the fitting analyses of FCS data often include some indeterminacy. Thus, the interpretation of molecular movements in living cells using only temporal autocorrelation analysis is not straightforward. In this study, the results of systematic mutation analyses support the interpretation suggested by FCS in living cells. Other fluorescence microscopic techniques were also performed to clarify the interaction target. Based on the results, I propose that the 1st, 2nd, 3rd, and immobile (photobleaching) components of the FCS analyses reflected 1) a free diffusion state of monomeric EGFP-FMBP-1 in cell nuclei (Fig. 14A), 2) a transient DNA interaction by electrostatic interaction on chromosomal DNA (Fig. 14B), 3) another transient DNA interaction that is strongly correlated with stable DNA binding (Fig. 14C), and 4) the final stable binding state (Fig. 14D), respectively. These components might correspond to each step that FMBP-1 takes in the recognition of its DNA binding sequence. In particular, as the polyploid nuclei of the PSG cells hold extremely large DNA contents, the two types of transient interactions observed in this study might play important roles, allowing FMBP-1 proteins to reach their final recognition sequence sites on the chromosomal DNA for the efficient expression of the fibroin gene.




Fig. 14. DNA interaction schema of FMBP-1 in vivo suggested by FCS analyses. This schema was drawn based on findings from our FCS analyses. In this schema, four different FMBP-1 movements during progressive interactions between FMBP-1 and DNA are shown.

2-4. Materials and Methods

2-4-1. *Plasmid constructs*

A cDNA encoding FMBP-1 was PCR amplified and then subcloned into the pEGFP-C1 vector (Clontech, Palo Alto, CA) downstream of the EGFP-coding region; *HindIII* and *KpnI* were used to subclone the cDNA. Mutants consisting of R9A(rep1) (substitution of alanine for the ninth arginine of repeat 1 [R107] of the STPR domain of FMBP-1), R9A(rep2) (substitution of alanine for the ninth arginine of repeat 2 [R130]), R9A(rep3) (substitution of alanine for the ninth arginine of repeat 3 [R153]), R9A(rep4) (substitution of alanine for the ninth arginine of repeat 4 [R176]), Δ N-HBR (a deletion extending from M1 to G95 of FMBP-1), Δ N-AR (a deletion extending from M1 to L82), Δ Ctail (a deletion extending from K191 to T218), Δ C-rep4 (a deletion extending from E168 to T218), Δ C-STPR (a deletion extending from E99 to T218), and Δ C-HBR (a deletion extending from K86 to T218) were each created using the QuikChange Site-Directed Mutagenesis Kit (Stratagene, La Jolla, CA) according to the manufacturer's instructions. The integrity of each plasmid was confirmed via DNA sequencing.

2-4-2. *Transient expression in silk gland cells*

The silk glands from fifth instar, day-0 *B. mori* larvae were dissected. Then, the Helios Gene Gun system (Bio-Rad, Hercules, CA) was used as previously described [37] to introduce tungsten particles coated with plasmid DNA into silk gland tissues incubated in Petri dishes. Each silk gland was transplanted into another fifth instar on day 0; the transplants were incubated at 25°C for 17 h. After incubation, the

transplanted silk glands were removed from the host larvae. Then, the PSG tissues were separated from each transplanted silk gland and mounted on a slide glass immediately before LSM imaging and FCS measurement. During these procedures, PSG tissues were maintained in phosphate-buffered saline (PBS). For transient expression in both PSG and HeLa cells, the CMV promoter, which is encoded in the mammalian expression vector pEGFP-C1, was used to drive EGFP-FMBP-1 expression. Expression levels of EGFP-FMBP-1 were high enough to make FCS measurements.

2-4-3. Transient expression in HeLa cells

HeLa cells were plated in 8-well Lab-Tek Chambered coverslips (Nalge Nunc International, Rochester, NY) and incubated in Dulbecco's modified Eagle's medium (DMEM, Sigma-Aldrich, St. Louis, MO) supplemented with 10% fetal bovine serum, 100 units/mL penicillin, and 100 µg/mL streptomycin in a 5% CO₂ humidified atmosphere at 37°C for 24 h. Transfection was conducted using Optifect transfection reagent (Invitrogen, Carlsbad, CA) according to the manufacturer's instructions. Cells were then grown in DMEM supplemented with 10% fetal bovine serum for 24 h in a 5% CO₂ humidified atmosphere. During LSM imaging and FCS measurement, HeLa cells were maintained in Opti-MEM reduced-serum medium (Invitrogen).

2-4-4. LSM imaging and FCS measurement

Both LSM imaging and FCS measurements were performed using an LSM510 inverted confocal laser scanning microscope (Carl Zeiss, Jena, Germany), which comprised a CW Ar⁺ laser, a water immersion objective (C-Apochromat, 40×, 1.2 NA

[Carl Zeiss]), and a ConfoCor 3 (Carl Zeiss). For LSM imaging of PSG cells, the cells were stained with 25 µg/mL Hoechst 33342 (Invitrogen) to visualize the nuclei. The Hoechst dye was excited by a 405-nm diode laser, and the emission signal was detected through a 420–480-nm band-pass filter. EGFP was excited by a 488-nm laser, and the emission signal was detected through a 505–540-nm band-pass filter.

Each FCS measurement was performed in the nucleus of an individual cell, avoiding the nuclear membranes and nucleoli. The output level of the excitation laser was kept at constant intensity during each measurement, at 112.5 µW or 225 µW for PSG or HeLa cells, respectively. The actual values of the laser intensity on the stage of the microscope were approximately 2.5 µW during each measurement for both cell types. The sequential illumination periods during the measurement of fluorescence intensity were 15.0 µsec each. The data were processed in essentially the same manner as described previously [23], except that the measurement time (60 sec, consisting of six consecutive measurements of 10 sec each) differed. For quantitative analyses, the fluorescence autocorrelation function (ACF) $G(\tau)$ was calculated as follows:

$$G(\tau) = \frac{\langle I(t)I(t+\tau) \rangle}{\langle I(t) \rangle^2}, \quad (1)$$

where τ represents the time delay, I is the fluorescence intensity, and the brackets denote the ensemble average. The binning period for extracting the ACF was 0.2 µsec. To calculate the decay rate of fluorescence intensity for samples that showed photobleaching, the fluorescence fluctuations were fitted by the following equation:

$$y = A * \exp(-t * k) + y_0, \quad (2)$$

where A is the height of photobleaching, k is the rate constant of photobleaching, and y_0 is the steady-state level of fluorescence fluctuation. Then, the fluorescence fluctuation

data in the photobleaching time period (first 20 sec or 10 sec for PSG cells or HeLa cells, respectively) were removed from the temporal and ensemble average to extract the ACF.

$G(\tau)$ was fitted with LSM-FCS software (Carl Zeiss) by multicomponent diffusion models that are described as follows:

$$G(\tau) = \frac{1}{N} \left[\sum_{i=1}^M \frac{F_i}{(1+(t/\tau_i)^\alpha) \sqrt{1+1/S^2(t/\tau_i)^\alpha}} \right] + 1, \quad (3)$$

where M is the number of fluorescent components and N is the average number of fluorescent molecules in the detection volume. F_i and τ_i are the component ratio and translational diffusion time of the i -th fluorescent component, respectively. S is the structural parameter of the instrumental set up $S = z/\omega$, where ω and z are the e^{-2} beam radius in the lateral and the axial directions of focus, respectively. The exponent α accounts for the mechanism of diffusion; $\alpha = 1$ for free (Brownian) diffusion and $\alpha < 1$ for obstructed (anomalous) diffusion. The fitting model did not include any expression for the triplet-state relaxation of EGFP, as triplet-state relaxation has a much longer lifetime than that of general fluorescence. Therefore, triplet-state relaxation often causes miscalculation of the diffusion parameters. If the influence of triplet-state relaxation substantially affected the extracted functions, I could use fitting models that included an expression for this; however, when I included an expression for the triplet-state term in the fitting model, it had very little influence on the results (Table S6). In this study, the fitting ranges for ACFs were set from $\tau \approx 50$ μ sec to 3.4 sec in order to avoid large variation in ACFs at early time points. Triplet-state relaxation has been reported to influence the extracted functions mainly at the early ACF (at $\tau \approx$ few μ sec); generally, the influence of triplet-state relaxation decreases exponentially. Thus, the influence of

triplet-state relaxation should be weak in the fitting range ($\tau \approx 50 \mu\text{sec}$ to 3.4 sec) for my FCS analyses. Therefore, I used a fitting model that did not include an expression for the triplet-state term in this study.

In the case of ACFs with correlations that were too small (due to high fluorescence intensity caused by a high concentration of EGFP-tagged protein) to fit a curve normally, the function was removed from the dataset. As a result, the average of the concentration of EGFP-FMBP-1 in this FCS measurement was approximately 400 nM (the range was $\approx 100\text{--}1000$ nM).

The diffusion coefficient D_i ($i = 1, 2, 3$) of a fluorescent species could then be assessed as follows:

$$D_i = \omega^2 / (4\tau_i), \quad (4)$$

where ω at 488-nm excitation was estimated from the diffusion coefficient of rhodamine 6G (R6G) [38].

The apparent molecular masses of EGFP-FMBP-1 monomers or EGFP-FMBP-1-containing complexes (Mw_i), assuming a spherical shape for the diffusing species, were assessed from the diffusion times by using the following mathematical expression based on the Stokes-Einstein equation as follows:

$$Mw_i = 27 \left(\frac{\tau_i}{\tau_{EGFP}} \right)^3. \quad (5)$$

The molecular mass of EGFP is 27 kDa. τ_{EGFP} is the diffusion time of EGFP alone when measured under the same experimental conditions as EGFP-FMBP-1.

The statistical analysis used to compare the averages of the best fit parameters was carried out with unpaired t -tests using Microsoft Excel 2013 software (Microsoft, Redmond, WA). To examine the appropriate number of components for the fitting

model, a comparison of different fitting models was carried out using AIC and with F -tests, also using Microsoft Excel 2013 software, where:

$$AIC = N \cdot \ln\left(\frac{\chi^2}{N}\right) + 2K + \frac{2K(K+1)}{N-K-1} \quad (6)$$

N is the number of data points used for each curve fitting, and K is the number of parameters. More precisely, the formula showed a corrected AIC that was corrected for the influence of sample size:

$$F = \frac{(\chi_1^2 - \chi_2^2)/(DF_1 - DF_2)}{\chi_2^2/DF_2} \quad (7)$$

where χ_1^2 is the χ^2 for the simpler model and χ_2^2 is the χ^2 for the more complicated model. DF is the degree of freedom ($N-K-1$). P -values for each F -ratio were calculated using the "FDIST" formula of Microsoft Excel 2013. The χ^2 values for each fitting were calculated as follows:

$$\chi^2 = \sum_i \left[\frac{y(x_i) - y_i}{\sigma_i} \right]^2. \quad (8)$$

The formula shows the difference between the fitted function $y(x)$ and the experimental data y_i at points x_i weighted by σ_i as a sum over all data points i . Where σ_i is the standard deviation for experimental point i .

For the statistical analysis, I referred to articles [18, 22, 39] and a manual for the online Graphpad Software (<http://www.graphpad.com/quickcalcs/AIC2.cfm>).

2-4-5. FRAP measurement

FRAP measurements were performed in essentially the same manner as described previously [40, 41]. For the measurements, an LSM510 inverted confocal laser scanning microscope (Carl Zeiss) with same settings as for FCS measurements was used. A small

area (approximately 3.0 μm diameter circle) was positioned in a region of the nucleus that did not contain the nucleolus, and bleached using 100% Ar^+ laser power (7.5 mW) with 15 scans. Images were then collected using 2.0% laser power every 2.5 sec for 3.0 min (for the samples showing fast fluorescence recovery) or every 5.0 sec for 6 min (for those with slow fluorescence recovery). The relative fluorescence intensities in the bleached area were normalized using the average intensity of three images measured before bleaching. The normalized intensities were analyzed using a fitting equation for a single exponential association model.

2-4-6. Confocal imaging for observation of colocalization

To observe the colocalization of EGFP-FMBP-1 and chromatin, I utilized a plasmid encoding mCherry-tagged histone H2B protein (H2B-mCherry), which was constructed by substitution of mCherry for GFP in the pBOS-H2BGFP vector [42]. The H2B-mCherry plasmid DNA was transiently co-expressed with EGFP-FMBP-1 in HeLa cells. For confocal imaging, an LSM510 inverted confocal laser scanning microscope (Carl Zeiss) with same settings as those for FCS measurements was used. The mCherry was excited by a 594-nm diode laser, and the emission signal was detected through a 650-nm long-pass filter.

2-4-7. Immunofluorescence staining and super-resolution fluorescence microscopy

Cells were cultivated on glass-based dishes (0.14–0.18 mm) (#3970-035, AGC Techno glass, Shizuoka, Japan), and then fixed with 4% paraformaldehyde buffered

with Hepes-KOH (pH 7.5) at 37°C. The Cells were washed in TBS and permeabilized in the presence of 0.2% (v/v) Triton X-100 (Nacalai tesque, Kyoto, Japan). After blocking non-specific binding activity in a blocking buffer containing 5% normal goat serum (DAKO, Glostrup, Denmark) and 20% Glycerol in PBS, the cells were incubated for 1 hour at room temperature in blocking buffer supplemented with primary antibodies against EGFP (#GF200, Nacalai tesque) and DsRed (#632496, TaKaRa-Clontech). The cells were then incubated with anti-mouse IgG antibody conjugated with AlexaFluor 488 or anti-rabbit IgG antibody conjugated with Alexa Fluor 594 (Thermo Fisher) in blocking buffer for 1 hour at room temperature. Cells stained on cover slips were mounted with ProLong Gold (Thermo Fisher). For super-resolution structured illumination microscopy (SIM), images were captured on an N-SIM system (Nikon, Tokyo, Japan) equipped with an Apo TIRF 100×/1.49 NA oil-immersion objective (Nikon). Alexa Fluor 488 was excited at 488 nm, and Alexa Fluor 594 was excited at 561 nm. Image acquisition and SIM-reconstruction were operated on an NIS-Elements software platform (Nikon). For high-resolution confocal microscopy, images were captured on a laser confocal microscope TCS SP8 (Leica, Wetzlar, Germany) equipped with an HC PL APO CS2 100×/1.40 OIL STED WHITE objective (Leica). AlexaFluor 488 and 594 was excited at 488 nm and 595 nm, respectively. Emitted fluorescence was sequentially detected using a photon-counting HyD detector through a spectral separation module (504-595 nm for AlexaFluor 488 and 603-653 nm for AlexaFluor 594). A pinhole size was set at 0.60 airy units (91 μ m). Acquired images were processed using a deconvolution algorithm on a Huygens software platform (Scientific Volume Imaging, Hilversum, Netherlands).

2-5. References

1. Kondo K, Aoshima Y, Hagiwara T, Ueda H & Mizuno S (1987) Tissue-specific and periodic changes in the nuclease sensitivity of the fibroin gene chromatin in the silkworm *Bombyx mori*. *J Biol Chem* **262**, 5271-5279.
2. Tashiro Y, Morimoto T, Matsuura S & Nagata S (1968) Studies on the posterior silk gland of the silkworm, *Bombyx mori*. 1. Growth of posterior silk gland cells and biosynthesis of fibroin during the fifth larval instar. *J Cell Biol* **38**, 574-588.
3. Durand B, Drevet J & Couble P (1992) P25 gene regulation in *Bombyx mori* silk gland: Two promoter-binding factors have distinct tissue and developmental specificities. *Mol Cell Biol* **12**, 5768-5777.
4. Horard B, Julien E, Nony P, Garel A & Couble P (1997) Differential binding of the *Bombyx* silk gland-specific factor SGFB to its target DNA sequence drives posterior-cell-restricted expression. *Mol Cell Biol* **17**, 1572-1579.
5. Hui CC & Suzuki Y (1989) Enhancement of transcription from the Ad2 major late promoter by upstream elements of the fibroin- and sericin-1-encoding genes in silk gland extracts. *Gene* **85**, 403-411.
6. Hui CC, Matsuno K & Suzuki Y (1990) Fibroin gene promoter contains a cluster of homeodomain binding sites that interact with three silk gland factors. *J Mol Biol* **213**, 651-670.
7. Mach V, Takiya S, Ohno K, Handa H, Imai T & Suzuki Y (1995) Silk gland factor-1 involved in the regulation of *Bombyx* sericin-1 gene contains fork head motif. *J Biol Chem* **270**, 9340-9346.
8. Fukuta M, Matsuno K, Hui CC, Nagata T, Takiya S, Xu PX, Ueno K & Suzuki Y (1993) Molecular cloning of a POU domain-containing factor involved in the regulation of the *Bombyx* sericin-1 gene. *J Biol Chem* **268**, 19471-19475.
9. Takiya S, Kokubo H & Suzuki Y (1997) Transcriptional regulatory elements in the upstream and intron of the fibroin gene bind three specific factors POU-M1, Bm Fkh and FMBP-1. *Biochem J* **321**, 645-653.
10. Zhao XM, Liu C, Li QY, Hu WB, Zhou MT, Nie HY, Zhang YX, Peng ZC, Zhao P & Xia QY (2014) Basic helix-loop-helix transcription factor Bmsage is involved in regulation of fibroin H-chain Gene via interaction with SGF1 in *Bombyx mori*. *PLoS One* **9**, e94091.
11. Takiya S, Ishikawa T, Ohtsuka K, Nishita Y & Suzuki Y (2005) Fibroin-modulator-binding protein-1 (FMBP-1) contains a novel DNA-binding domain, repeats of the score and three amino acid peptide (STP), conserved from *Caenorhabditis elegans* to humans. *Nucleic Acids Res* **33**, 786-795.
12. Saito S, Aizawa T, Kawaguchi K, Yamaki T, Matsumoto D, Kamiya M, Kumaki Y, Mizuguchi M, Takiya S, Demura M & Kawano K (2007) Structural approach to a novel tandem repeat DNA-binding domain, STPR, by CD and NMR. *Biochemistry* **46**, 1703-1713.
13. Saito S, Yokoyama T, Aizawa T, Kawaguchi K, Yamaki T, Matsumoto D, Kamijima T, Kamiya M, Kumaki Y, Mizuguchi M, Takiya S, Demura M & Kawano K (2008) Structural properties of the DNA-bound form of a novel tandem repeat DNA binding domain, STPR. *Proteins* **72**, 414-426.
14. Takiya S, Saito S, Yokoyama T, Matsumoto D, Aizawa T, Kamiya M, Demura M & Kawano K (2009) DNA-binding property of the novel DNA-binding domain STPR in FMBP-1 of the silkworm *Bombyx mori*. *J Biochem* **146**, 103-111.
15. Brock R, Vamosi G, Vereb G & Jovin TM (1999) Rapid characterization of green fluorescent protein fusion proteins on the molecular and cellular level by fluorescence correlation microscopy. *Proc Natl Acad Sci USA* **96**, 10123-10128.
16. Lippincott-Schwartz J, Snapp E & Kenworthy A (2001) Studying protein dynamics in

- living cells. *Nat Rev Mol Cell Biol* **2**, 444-456.
17. Berland KM (2004) Fluorescence correlation spectroscopy: a new tool for quantification of molecular interactions. *Methods Mol Biol* **261**, 383-398.
 18. Brazda P, Szekeres T, Bravics B, Toth K, Vamosi G & Nagy L (2011) Live-cell fluorescence correlation spectroscopy dissects the role of coregulator exchange and chromatin binding in retinoic acid receptor mobility. *J Cell Sci* **124**, 3631-3642.
 19. Giorgio G, Yinghua G, Jagesh VS & Galit L (2013) Activation and control of p53 tetramerization in individual living cells. *Proc Natl Acad Sci USA* **110**, 15497-15501.
 20. Tiwari M, Mikuni S, Muto H & Kinjo M (2013) Determination of dissociation constant of the NF κ B p50/p65 heterodimer using fluorescence cross-correlation spectroscopy in the living cell. *Biochem Biophys Res Commun* **436**, 430-435.
 21. Vukojević V, Papadopoulos DK, Terenius L, Gehring WJ & Rigler R (2009) Quantitative study of synthetic Hox transcription factor-DNA interactions in live cells. *Proc Natl Acad Sci USA* **107**, 4093-4098.
 22. Levin MK & Carson JH (2004) Fluorescence correlation spectroscopy and quantitative cell biology. *Differentiation* **72**, 1-10.
 23. Nonaka Y, Muto H, Aizawa T, Okabe E, Myoba S, Yokoyama T, Saito S, Tatami F, Kumaki Y, Kamiya M, Kikukawa T, Mizuguchi M, Takiya S, Kinjo M, Demura M & Kawano K (2010) STPR, a 23-amino acid tandem repeat domain, found in the human function-unknown protein ZNF821. *Biochemistry* **49**, 8367-8375.
 24. Wachsmuth M, Weidemann T, Müller G, Hoffmann-Rohrer UW, Knoch TA, Waldeck W & Langowski J (2003) Analyzing intracellular binding and diffusion with continuous fluorescence photobleaching. *Biophys J* **84**, 3353-3363.
 25. Hinojos CA, Sharp ZD & Mancini MA (2005) Molecular dynamics and nuclear receptor function. *Trends Endocrinol Metab* **16**, 12-18.
 26. Metivier R, Reid G & Gannon F (2006) Transcription in four dimensions: nuclear receptor-directed initiation of gene expression. *EMBO Rep* **7**, 161-167.
 27. Dix JA, Hom EF & Verkman AS (2006) Fluorescence correlation spectroscopy simulations of photophysical phenomena and molecular interactions: a molecular dynamics/monte carlo approach. *J Phys Chem B* **110**, 1896-1906.
 28. Winter RB, Berg OG & von Hippel OH (1981) Diffusion-driven mechanisms of protein translocation on nucleic acids. 3. The *Escherichia coli* lac repressor-operator interaction: kinetic measurements and conclusions. *Biochemistry* **20**, 6961-6977.
 29. Hannon R, Richards EG & Gould HJ (1986) Facilitated diffusion of a DNA binding protein on chromatin. *EMBO J* **5**, 3313-3319.
 30. Halford SE & Marko JF (2004) How do site-specific DNA-binding proteins find their targets? *Nucleic Acids Res* **32**, 3040-3052.
 31. Komazin-Meredith G, Mirchev R, Golan DE, van Oijen AM & Coen DM (2008) Hopping of a processivity factor on DNA revealed by single-molecule assays of diffusion. *Proc Natl Acad Sci USA* **105**, 10721-10726.
 32. Gorman J & Greene EC (2008) Visualizing one-dimensional diffusion of proteins along DNA. *Nat Struct Mol Biol* **15**, 768-774.
 33. Blainey PC, Luo G, Kou SC, Mangel WF, Verdine GL, Bagchi B & Xie XS (2009) Nonspecifically bound proteins spin while diffusing along DNA. *Nat Struct Mol Biol* **16**, 1224-1229.
 34. Schonhofs JD & Stivers JT (2012) Timing facilitated site transfer of an enzyme on DNA. *Nat Chem Biol* **8**, 205-210.
 35. Terakawa T, Kenzaki H & Takada S (2012) p53 searches on DNA by rotation-uncoupled sliding at C-terminal tails and restricted hopping of core domains. *J Am Chem Soc* **134**, 14555-14562.
 36. Gorman J, Wang F, Redding S, Plys AJ, Fazio T, Wind S, Alani EE & Greene EC (2012)

- Single-molecule imaging reveals target-search mechanisms during DNA mismatch repair. *Proc Natl Acad Sci USA* **109**, E3074.
37. Takiya S, Inoue H & Kimoto M (2011) Novel enhancer and promoter elements indispensable for the tissue-specific expression of the sericin-1 gene of the silkworm *Bombyx mori*. *Insect Biochem Mol Biol* **41**, 592-601.
 38. Petrov EP & Schuille P (2008) State of the art and novel trends in fluorescence correlation spectroscopy. In standardization and quality assurance in fluorescence measurements. *Springer Ser Fluoresc* **6**, 145-197.
 39. Wohland T, Rigler R & Vogel H (2001) The standard deviation in fluorescence correlation spectroscopy. *Biophys J* **80**, 2987-2999.
 40. Kimura H & Cook PR (2001) Kinetics of core histones in living human Cells: little exchange of H3 and H4 and some rapid exchange of H2B. *J Cell Biol* **153**, 1341-1353.
 41. Kitamura A, Kubota H, Pack CG, Matsumoto G, Hirayama S, Takahashi Y, Kimura H, Kinjo M, Morimoto RI & Nagata K (2006) Cytosolic chaperonin prevents polyglutamine toxicity with altering the aggregation state. *Nat Cell Biol* **8**, 1163-1170.
 42. Kanda T, Sullivan KF & Wahl GM (1998) Histone–GFP fusion protein enables sensitive analysis of chromosome dynamics in living mammalian cells. *Curr Biol* **8**, 377-385.

2-7. Supporting Tables

Table S1. AIC test of each fitting model for wild-type EGFP-FMBP-1 data measured in PSG cells

Measurement	Number of data points	1 component, free diffusion model		2 component, free diffusion model		3 component, free diffusion model	
		Chi ²	AIC	Chi ²	AIC	Chi ²	AIC
1	134	97.3	-38.8	13.7	-297.2	13.1	-299.3
2	132	192.6	53.9	18.0	-254.4	15.0	-274.2
3	135	195.2	53.9	25.4	-217.0	8.4	-362.7
4	134	159.2	27.2	23.1	-227.2	10.6	-327.9
5	126	415.0	154.3	37.0	-146.0	32.6	-157.8
6	132	223.9	73.8	32.5	-176.8	20.8	-231.0
7	142	379.9	143.8	45.6	-153.1	35.6	-183.7
8	132	443.4	164.0	35.2	-166.1	32.7	-171.4
9	126	272.2	101.1	54.2	-98.0	51.7	-99.5
10	131	743.4	231.5	39.8	-147.7	27.5	-191.9
11	127	170.3	41.4	17.5	-243.0	11.2	-295.2
12	140	93.4	-52.5	16.0	-295.7	15.5	-295.4
13	130	149.7	22.4	29.7	-183.8	19.8	-232.0
14	134	410.2	154.0	22.8	-228.8	19.4	-246.4
15	133	295.7	110.4	30.4	-188.1	11.6	-311.9
16	129	652.7	213.2	33.2	-166.6	27.8	-185.3
17	119	1331.7	291.5	42.9	-113.1	39.2	-119.3
18	118	164.9	43.6	29.9	-153.5	25.8	-166.7
19	130	213.6	68.6	28.7	-188.3	28.1	-186.3
20	134	183.7	46.3	18.7	-255.4	13.0	-300.0
21	136	108.7	-26.4	16.5	-278.9	12.2	-315.3
22	127	212.9	69.7	32.4	-165.0	30.1	-170.1
23	129	577.3	197.4	29.0	-184.1	28.0	-184.4
24	131	765.7	235.4	58.5	-97.2	53.5	-104.6
25	133	169.6	36.4	17.7	-259.6	10.0	-331.4
26	132	380.4	143.8	29.9	-187.8	13.8	-285.6
27	130	308.0	116.2	19.2	-240.5	15.9	-260.7
28	128	166.2	37.5	19.9	-229.7	13.3	-277.0
29	124	486.5	173.6	23.9	-195.8	20.7	-209.4

The AIC values were calculated from the χ^2 of curve fitting for wild-type FMBP-1 in PSG cells (total 29 fitting). Each AIC value was generated using the formula (6) shown in *Materials and Methods*. AIC represents the appropriateness of the fitting model used in curve fitting. The appropriateness is judged by the value; lower values indicate a more probable fitting model. AIC values in bold type in the three-component, free diffusion model column are the lowest among the three different models.

Table S2. *F*-test of the three-component, free diffusion model fitting against the two-component, free diffusion model for wild-type EGFP-FMBP-1 data in PSG cells

Measurement	<i>F</i> -value	<i>P</i> -value
1	3.15	< 0.05
2	12.59	< 0.005
3	130.50	< 0.005
4	75.61	< 0.005
5	8.19	< 0.005
6	34.92	< 0.005
7	18.87	< 0.005
8	4.74	< 0.05
9	2.80	0.06
10	27.83	< 0.005
11	33.66	< 0.005
12	1.93	0.15
13	30.61	< 0.005
14	11.29	< 0.05
15	102.13	< 0.005
16	11.92	< 0.005
17	5.22	< 0.05
18	8.96	< 0.005
19	1.16	0.32
20	28.01	< 0.005
21	22.54	< 0.005
22	4.65	< 0.05
23	2.22	0.11
24	5.82	< 0.05
25	48.68	< 0.005
26	73.07	< 0.005
27	12.79	< 0.005
28	30.08	< 0.005
29	9.13	< 0.005

Using Microsoft Excel 2013 software, *F*-tests were conducted to compare the χ^2 values of the three-component model with those of the two-component model for each of the 29 measurements from the wild-type EGFP-FMBP-1 measured in PSG cells. *F*-values were calculated by the formula (7) shown in *Materials and Methods*. *F*-values and the corresponding *P*-values or *P*-value upper bounds are listed for each measurement in the Table. Significant values (<0.05) are shown in bold.

Table S3. Diffusion parameters of wild-type EGFP-FMBP-1 in PSG obtained by various diffusion models

(a) Two-component, free diffusion model

1st component		2nd component	
F_1 (%)	τ_1 (μ s)	F_2 (%)	τ_2 (ms)
70.8 ± 7.1	678.7 ± 169.3	29.2 ± 7.1	22.03 ± 12.81

(b) One-component, anomalous-diffusion model

1st component (anomalous)	
τ_1 (μ s)	A
1188.8 ± 462.6	0.63 ± 0.05

(c) Two-component, anomalous-diffusion model (one components is anomalous; the other is free)

1st component (anomalous)			2nd component (free)	
F_1 (%)	τ_1 (μ s)	α	F_2 (%)	τ_2 (ms)
75.1 ± 19.6	3058.0 ± 7535.0	0.76 ± 0.12	24.9 ± 19.6	48.27 ± 89.96

(d) Two-component, anomalous-diffusion model (both components are anomalous)

1st component (anomalous)			2nd component (anomalous)		
F_1 (%)	τ_1 (μ s)	A	F_2 (%)	τ_2 (ms)	α
71.4 ± 28.2	788.4 ± 446.7	0.86 ± 0.43	28.6 ± 28.2	$1.28\text{E}+8 \pm 6.76\text{E}+8$	1.44 ± 1.23

(e) Three-component anomalous-diffusion model (all components are anomalous)

1st component (anomalous)			2nd component (anomalous)			3rd component (anomalous)		
F_1 (%)	τ_1 (μ s)	α	F_2 (%)	τ_2 (ms)	α	F_3 (%)	τ_3 (ms)	α
44.9 ± 35.6	482.3 ± 403.9	33.95 ± 107.57	35.8 ± 30.2	5.82 ± 8.49	1.69 ± 1.15	19.3 ± 19.9	96.8 ± 159.5	5.65 ± 18.25

Note: α values are anomaly parameters; $\alpha = 1$ for free (Brownian) diffusion, $\alpha < 1$ for obstructed (anomalous) diffusion.

(f) Comparison of AIC values between the three-component, free-diffusion model and each anomalous-diffusion model.

	1-component, anomalous		2-component, anomalous (one of the pair is anomalous)		2-component, anomalous (both components are anomalous)		3-component, anomalous (all components are anomalous)		3-component, free
Measurement	Chi ²	AIC	Chi ²	AIC	Chi ²	AIC	Chi ²	AIC	AIC
1	16.5	-274.7	13.1	-301.4	12.8	-301.7	12.4	-301.7	-299.3
2	20.3	-241.1	15.2	-274.7	14.9	-275.2	12.6	-292.7	-274.2
3	10.5	-338.0	7.5	-380.5	7.5	-377.4	7.3	-377.2	-362.7
4	9.9	-343.1	10.1	-335.9	9.8	-337.6	9.0	-344.6	-327.9
5	39.4	-140.2	31.7	-163.2	21.9	-208.0	21.3	-206.8	-157.8
6	22.7	-226.0	19.1	-244.6	19.2	-242.1	17.3	-251.2	-231.0
7	37.7	-182.3	35.7	-185.8	33.6	-192.0	35.2	-180.8	-183.7
8	42.4	-143.6	32.4	-175.0	31.5	-176.7	30.1	-174.5	-171.4
9	54.2	-100.2	51.3	-102.7	48.4	-107.8	47.5	-105.7	-99.5
10	51.3	-116.6	28.2	-190.9	26.6	-196.0	26.2	-193.7	-191.9
11	14.4	-269.9	10.8	-302.9	10.8	-300.3	9.7	-309.5	-295.2
12	20.3	-264.4	18.6	-272.2	18.2	-272.7	17.8	-271.6	-295.4
13	20.9	-231.6	20.1	-232.0	19.7	-232.8	19.6	-228.5	-232.0
14	25.1	-218.5	18.3	-256.5	17.6	-259.6	16.4	-264.6	-246.4
15	25.1	-215.6	8.9	-349.3	8.8	-348.6	8.4	-350.4	-311.9
16	35.3	-161.1	28.6	-183.8	25.1	-198.3	27.3	-182.9	-185.3
17	65.7	-64.5	44.4	-106.9	36.1	-129.1	18.7	-202.7	-119.3
18	31.6	-149.4	26.0	-167.8	26.7	-162.6	23.4	-176.5	-166.7
19	45.7	-129.8	27.6	-190.9	23.5	-209.7	23.1	-207.5	-186.3
20	18.8	-257.1	13.1	-301.5	13.0	-300.0	10.4	-325.4	-300.0
21	10.8	-338.1	12.1	-318.3	11.3	-325.2	11.4	-320.6	-315.3
22	38.3	-146.1	30.2	-172.0	30.9	-166.9	29.0	-170.3	-170.1
23	53.7	-106.8	27.6	-188.6	23.1	-209.4	23.1	-204.8	-184.4
24	48.4	-124.1	44.2	-131.8	48.4	-117.8	48.3	-113.6	-104.6
25	9.9	-338.9	10.0	-333.8	9.7	-335.8	9.4	-335.4	-331.4
26	16.6	-267.8	14.6	-280.1	14.8	-276.6	13.1	-287.4	-285.6
27	18.7	-246.1	16.6	-257.4	14.6	-271.2	11.8	-295.0	-260.7
28	12.8	-288.3	11.4	-299.0	11.7	-293.8	10.4	-303.5	-277.0
29	19.0	-226.6	19.7	-217.3	19.0	-219.7	14.9	-257.8	-209.4

The AIC values calculated from the fitting residuals of wild-type FMBP-1 in PSG cells (total 29 fitting). This comparison of AIC values was processed in the same manner as described for Table S1. In more than half of the fitting samples, the one-component, anomalous-diffusion model was judged to be improbable compared with the three-component, free diffusion model. The other two anomalous-diffusion models were judged to be relatively probable. However, the differences in AIC values between the two anomalous-diffusion models and those of the three-component free diffusion model were not large for most fitting samples. For reference, the χ^2 and AIC values for the three-component, anomalous diffusion model are also displayed.

Table S4. Examination of the appropriateness of the interpretation with the three-component model for the R9A(rep1) mutant in HeLa cells

(a) Diffusion parameters of the R9A(rep1) mutant determined by using the two-component, free diffusion model

1st component		2nd component	
F_1 (%)	τ_1 (μ s)	F_2 (%)	τ_2 (ms)
63.7 ± 12.2	617.9 ± 174.4	36.3 ± 12.2	6.62 ± 3.44

In comparison with the diffusion parameters obtained by the three-component free diffusion model (shown in Table 2), the component ratios and diffusion times of each component were clearly different. Also, dispersion of each parameter was increased.

(b) Comparison of fitting residuals of the R9A(rep1) mutant by the three-component model and the two-component model

Measurement	3-component model		2-component model	
	Chi ²	AIC	Chi ²	AIC
1	10.0	-352.8	22.2	-246.6
2	5.6	-395.9	12.1	-300.7
3	21.9	-207.6	24.8	-196.5
4	8.6	-358.6	12.0	-318.3
5	21.2	-223.4	29.6	-183.9
6	16.7	-253.7	27.0	-195.8
7	29.0	-200.1	36.6	-172.6
8	10.2	-314.9	11.6	-302.5

The AIC values calculated from the fitting residuals of R9A(rep1) mutants in HeLa cells (total 8 fitting). This comparison of AIC values was processed in the same manner as described in Table S1. In all eight fittings, the three-component free diffusion model was judged to be more probable than the two-component model.

Table S5. Influence of photobleaching for diffusion parameters

	1st component		2nd component		3rd component		<i>n</i>
	F_1 (%)	τ_1 (μ s)	F_2 (%)	τ_2 (ms)	F_3 (%)	τ_3 (ms)	
wild-type FMBP-1 (20-30 sec)	41.5 ± 11.7	299.1 ± 77.4	43.5 ± 8.0	2.78 ± 1.61	14.9 ± 11.1	114.6 ± 75.7	16
wild-type FMBP-1 (50-60 sec)	37.5 ± 17.1	330.7 ± 63.5	43.6 ± 14.8	2.86 ± 1.96	18.9 ± 9.9	73.3 ± 51.9	19

These values are derived by curve fitting of the autocorrelation function (ACF) calculated from the third (20-30 sec) or last (50-60 sec) 10 seconds of continuous fluorescence fluctuation data ($10 \text{ sec} \times 6 \text{ times}$) of wild-type FMBP-1 in PSG cells with the three-component, free diffusion model. Each of 16 or 19 samples of all 29 measured samples could be fit. However, other samples could not be fit well, resulting in very few component ratios and large diffusion times for each component. It is impossible to consider the ratios and times are real values. Such difficulty of curve fitting would be caused by large variation of the ACF, calculated from shorter fluorescence fluctuation than normal curve fitting of ACF in this study (which was calculated from continuous $10 \times 4 \text{ times}$ measurement data).

Table S6. Influence of triplet-state relaxation for diffusion parameters

	1st component		2nd component		3rd component		
	F_1 (%)	τ_1 (μ s)	F_2 (%)	τ_2 (ms)	F_3 (%)	τ_3 (ms)	n
wild-type FMBP-1	45.6 ± 13.7	347.4 ± 56.9	41.1 ± 7.1	3.37 ± 1.70	13.3 ± 9.6	102.1 ± 88.9	28

These values are derived by curve fitting of the autocorrelation function of wild-type FMBP-1 in PSG cells with the three-component diffusion model incorporating triplet-state relaxation as follows:

$$G(\tau) = \frac{1 + \frac{T}{1-T} e^{-\tau/\tau_T}}{N} \left[\sum_{i=1}^M \frac{F_i}{(1 + \tau/\tau_i) \sqrt{1 + \tau/(S^2 \tau_i)}} \right] + 1.$$

The triplet-state term was inserted in each fitting model that was used for the present study (shown in *Materials and Methods*). T and τ_T are the fractional population and decay time of the triplet state, respectively. I tested the fitting model for the same data samples of wild-type EGFP-FMBP-1 measured in PSG cells. Each of the 28 samples could be fit well with this fitting model. However, one sample (the 29th measurement sample in Table S1 and S2) could not be fit well, resulting in very few component ratios of the 3rd component, which is impossible to consider real value. Thus, I excluded the one datum from the calculation of average diffusion parameters as shown in the table above.

Acknowledgements

I gratefully acknowledge the invaluable suggestions and support of Associate Prof. T. Aizawa and Prof. M. Kinjo. I would like to express my gratitude to Emeritus Prof. K. Kawano, Prof. M. Demura, Associate Prof. S. Takiya, Associate Prof. T. Kikukawa, Assistant Prof. A. Kitamura, Assistant Prof. M. Kamiya and Dr. H. Muto. They have been a tremendous mentor for me. I would like to thank you for encouraging my research and for allowing me to grow as a research scientist. I also appreciate a lot of advices by Prof. H. Haga as deputy chief examiner of my doctoral defense.

I also deeply appreciate the efforts of Dr. M. Kimoto and Mr. S. Myoba in collaborating in this study.

I would also like to thank the Nikon Imaging Center at Hokkaido University for allowing us to use N-SIM on super-resolution microscopy. I also thank Mr. T. Igarashi and Ms. R. Hayashi (Leica microsystems) for high-resolution fluorescence microscopy.

Last but not the least, I would like to thank my parents, my brother, my friends and my junior in our laboratory for supporting me spiritually throughout writing this thesis and my life in general.