

HOKKAIDO UNIVERSITY

Title	Mercury concentrations in primary feathers reflect pollutant exposure in discrete non-breeding grounds used by Short-tailed Shearwaters
Author(s)	Watanuki, Yutaka; Yamamoto, Takashi; Yamashita, Ai et al.
Citation	Journal of ornithology, 156(3), 847-850 https://doi.org/10.1007/s10336-015-1205-6
Issue Date	2015-07
Doc URL	https://hdl.handle.net/2115/62327
Rights	The final publication is available at Springer via http://dx.doi.org/10.1007/s10336-015-1205-6
Type	journal article
File Information	watanuki.pdf



Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

2015.2.14

Short Note to J Ornithology: 2nd revised version

Mercury concentrations in primary feathers reflect pollutant exposure in discrete non-breeding grounds used by Short-tailed Shearwaters

Yutaka Watanuki[1], Takashi Yamamoto[1, 2], Ai Yamashita[1], Chihiro Ishii[3], Yoshinori Ikenaka[3], Shouta M.M. Nakayama[3], Mayumi Ishizuka[3],Yuya Suzuki[1], Yasuaki Niizuma[4], C.E. Meathrel[5], R.A. Phillips[6]

[1]Graduate School of Fisheries Sciences, Hokkaido University, Minato-cho 3-1-1, Hakodate 041-8611, Japan ywata@fish.hokudai.ac.jp, [2]Arctic Environment Research Center, National Institute of Polar Research, Midori-cho 10-3, Tachikawa, 190-8518, Japan, [3]Graduate School of Veterinary Medicine, Hokkaido University, Kita-18 Nishi-9, Sapporo 060-0818, Japan, [4]Faculty of Agriculture, Meijyo University, Shiogamaguchi 1-501, Tenpaku-ku, Nagoya 468-8502, [5]LaTrobe University, Wodonga, Victoria 3086, Australia, [6]British Antarctic Survey, Natural Environment Research Council, High Cross, Madingley Road, Cambridge CB3 0ET, UK

Abstract We measured mercury concentrations ([Hg]) and nitrogen stable isotope ratios ($\delta^{15}N$) in the primary feathers of Short-tailed Shearwaters (*Puffinus tenuirostris*) that were tracked year-round. [Hg] were highest in 14 birds that used the Okhotsk and northern Japan Seas during the non-breeding period ($2.5 \pm 1.4\ \mu$ g/g), lowest in 9 birds that used the eastern Bering Sea ($0.8 \pm 0.2\ \mu$ g/g), and intermediate in 5 birds that used both regions ($1.0 \pm 0.5\ \mu$ g/g), with no effects of $\delta^{15}N$. The results illustrate that samples from seabirds can provide a useful means of monitoring pollution at large spatial scale.

Key word: geolocator; trophic level; northern North Pacific; monitoring

INTRODUCTION

Mercury (Hg) is diffused globally through atmospheric and oceanic transport (Laurier et al. 2004), methylated and biomagnified in the marine ecosystem (Jaeger et al. 2009) and has adverse physiological effects on consumers. Although reliable approaches to monitoring spatial pattern of marine Hg pollution are needed, intensive and repeated sampling of seawater in offshore regions is financially and logistically challenging. As an alternative, seabird feathers may offer a viable method for monitoring Hg pollution including over large spatial scales (Bearhop et al. 2000).

Studies of seabirds tracked using geolocators show that individuals from the same breeding colony may use various discrete regions in pelagic waters during the non-breeding period (Phillips et al. 2008). Recent study showed that individuals of Great Skuas (*Catharacta skua*) carried rather different levels of contaminants depending on their wintering area (Leat et al. 2013). If molt patterns are known or inferred, Hg concentrations of feathers grown at a specific time of the year may therefore provide valuable information on environmental exposure within particular regions (Ramos et al. 2009).

Short-tailed Shearwaters (*Puffinus tenuirostris*) are trans-equatorial migrants that breed in southern Australia and spend the non-breeding period (May–Sep) in the northern North Pacific. To assess Hg exposure in these non-breeding grounds, we tracked Short-tailed Shearwaters using geolocators, and sampled outer most primary feathers (P10) when the birds returned to the colony for later analysis of

56 Hg concentrations ([Hg]) and $\delta^{15}N$ (a proxy of trophic level; Jaeger et al. 2009).

57

58 ## STUDY AREA AND METHODS

59 **Field work** The study was conducted at Great Dog, Flinders Islands, Tasmania
60 (40°15´S, 148°15´E). Geolocators (Mk15; British Antarctic Survey, Cambridge)
61 were attached to 50 and 46 incubating birds in early December in 2009 and 2010,
62 respectively. Geolocators weighed 2.5g (< 1% of mean body mass of study birds),
63 and were attached to aluminum leg bands with plastic ties (Carey et al. 2014).
64 Fifteen and 27 birds in the burrows were recaptured in early December 2010 and
65 2011, respectively (including 3 birds recaptured 2 years after deployment), and
66 one recovered from a beached bird in 2010. Tracking data were obtained from 40
67 birds (3 loggers could not be downloaded). At recapture, 1 cm from the tip of P10
68 was collected and stored at -20 °C. Based on body measurements (Carey 2011),
69 the sample of tracked birds was male-biased (32 males, 4 females, and 7
70 unknown) since males usually take the first incubation spell in early December
71 when we conducted the fieldwork.

72

73 **Track analysis** The geolocators measured light intensity, and immersion and
74 temperature in seawater. We estimated sunset and sunrise times from light curves,
75 then derived latitudes on the basis of day length, and longitudes from the time of
76 local midday and midnight. Day and night locations were averaged to give a
77 single location per day. During the period around the equinoxes, when latitude
78 cannot be estimated from the day length, we used the water-temperature data and
79 the light-based longitudes to estimate the daily latitude from maps of

remotely-sensed sea surface temperature (8-day composite, resolution 9 km, measured by Aqua-MODIS). Location data that were unreliable because of obvious interruptions around sunset and sunrise, or unrealistic flight speeds (>70 km/h) were replaced with those estimated by linear interpolation.

Chemical analysis Each feather was split into two at the rachis for Hg and stable isotope analyses. For Hg analysis, the feathers were washed using 99.5% acetone and Milli-Q water and dried in an oven at 50°C for 24 hrs. We measured [Hg] using CV-AAS (Cold Vapor-Atomic Absorption Spectroscopy) and a Mercury Analyzer MA-3000 (Nippon Instruments Corporation, Japan). Hg recoveries were between 90% and 105% for the laboratory standards (fish; DORM-3 and DOLT-4), and the detection limit was 0.2 ng/g (dry weight). For $\delta^{15}N$ analysis, feathers were cleaned using 0.25M sodium hydroxide, rinsed in Milli-Q water and dried at 60°C for 24 hrs. Dried samples were ground in an auto-mill after freezing using liquid nitrogen. The nitrogen stable isotope ratio (in ‰) was measured using a gas-source isotope ratio mass spectrometer (ANCA-GSL and Hydra 20-20, Sercon Ltd, UK), and is presented as deviations from atmospheric N_2, where $\delta^{15}N = [(^{15}N/^{14}N \text{ sample} / ^{15}N/^{14}N \text{ standard}) - 1] \times 1{,}000$. All samples were measured in triplicate and average values used in all statistical tests. If the coefficient of variation on triplicate measurements was over 0.3, the value with the largest deviation was excluded from calculation of the mean.

Statistical analyses The effect of year and non-breeding grounds (fixed effects) and $\delta^{15}N$ (covariate) on [Hg], and the effects of year and non-breeding grounds

(fixed) on $\delta^{15}N$ were examined by GLMs using SPSS statistics ver. 22. Sex was not included as a factor because of the male bias. No interaction terms were included. Means are presented ± SD.

RESULTS

Non-breeding grounds Feather samples of Short-tailed Shearwaters were not collected in every case, or too small for analysis, so [Hg] and $\delta^{15}N$ data were available for 28 tracked birds in this study. During the non-breeding period, 14 birds stayed in the southern Okhotsk and northern Japan Seas (WEST), 9 birds (including three tracked for two years) migrated to the eastern Bering Sea and around the Aleutian Islands (EAST), and 5 birds initially used the western region but later moved to the eastern region (MIX) (Fig. 1).

Mercury and $\delta^{15}N$ There was no significant effect of year ($F_{(1,23)}$=1.863, P=0.185) or $\delta^{15}N$ ($F_{(1,23)}$=0.053, P=0.820) on [Hg], but the effect of non-breeding grounds was significant ($F_{(2,23)}$=6.789, P=0.002). Mean [Hg] was higher in WEST than EAST birds (Fig. 2; P<0.05, Bonferroni post-hoc test), and did not differ significantly between MIX and WEST or EAST birds (P>0.05). The effect of non-breeding grounds on $\delta^{15}N$ was marginally significant ($F_{(2,24)}$=3.336, P=0.053) with $\delta^{15}N$ tended to be higher in EAST (15.2±2.9‰) than WEST (13.4±0.8‰) and MIX birds (13.0±1.7‰). Year effect was not significant ($F_{(1,24)}$=0.229, P=0.637).

DISCUSSION

128 Primary molt of Short-tailed Shearwaters has been recorded during late June and
129 July (non-breeding period) and is completed before return to the breeding site
130 (Marchant and Higgins 1990). Based on stable isotope and other data, the Sooty
131 Shearwaters *Puffinus griseus* also start to molt primaries on arrival at the
132 non-breeding grounds in the north-western Atlantic (Hedd et al. 2012). Although
133 there was substantial variation in molting patterns among populations and
134 individuals in Yelkouan Shearwaters *Puffinus yelkouan*, they rarely molted wing
135 feathers during breeding (Bourgeois and Dromzee 2014). Thus, the Hg
136 concentrations in P10 of the short-tailed shearwaters in our study presumably
137 reflect exposure to pollutants in the non-breeding grounds. Further information on
138 molt sequence in this species, including birds of different status and from different
139 colonies, would be valuable, however, as chemical analysis of different feather
140 types may provide details on pollutant exposure during the non-breeding period at
141 a finer temporal and spatial scale (González-Solís et al. 2011).
142
143 [Hg] in P10 of our Short-tailed Shearwaters (0.8 – 2.5 μg/g, Fig. 2) were lower
144 than those in the flight feathers that were replaced during the non-breeding period
145 by other species (1.2 – 3.9 μg/g in Cory's Shearwater *Calonectris diomedea*, Ramos
146 et al. 2009; 4.3 – 6.0 μg/g in Great Skua, Bearhop et al. 2000), and lower than the
147 levels sometimes associated with impaired reproduction (>5.0μg/g, Burger and
148 Gochfeld 1997). We further found that feather [Hg] were higher for the
149 Short-tailed Shearwaters that spent the non-breeding period in the southern
150 Okhotsk and northern Japan Seas (WEST), than those in the eastern Bering Sea
151 and around the Aleutian Islands (EAST). The influence of $\delta^{15}N$ on [Hg] in

feathers was not significant in our study birds, in comparison with results from other seabird species (Bond 2010). In addition, the $\delta^{15}N$ of WEST birds tended to be lower than that of EAST birds, and so the differences in trophic level would not explain the higher [Hg] in the WEST birds unless we postulate a substantial disparity in isotopic baselines in the two regions.

The spatial pattern of Hg pollution in offshore waters of the northern North Pacific Ocean has been examined by sampling seabird tissues and seawater during research cruises. [Hg] in the liver of Glaucous-winged Gulls (*Larus glaucescens*) increased towards the west along the Aleutian Island chain (Ricca et al. 2008). [Hg] in seawater was higher in the western North Pacific shelf area off Japan than in the basin and central North Pacific (Laurier et al. 2004). These findings provide general support for the spatial pattern in Hg exposure found in the Short-tailed Shearwaters tracked in our study.

Although the method we applied provided information on pollutant levels only in the areas visited by the tracked birds, our study nevertheless demonstrates the utility of this technique for monitoring the spatial pattern of Hg pollution in large offshore regions, which are beyond the ranges of breeding birds and where ship-based sampling is expensive and logistically challenging.

ACKNOWLEDGEMENTS

We thank R. Bruce, B. Nishizawa, and A. Ito for field assistance, L. Hardy for recovering geolocator from a beached bird. Research was carried out under La

Trobe University Animal Ethics Project AEC 10/32 (W) and Tasmanian Department of Primary Industries Permits to Take Wildlife for Scientific Purpose FA10161 and FA11212 held by CE Meathrel. The study was supported by JSPS (#20241001, #22405007) and GRENE project.

REFERENCES

Bearhop S, Phillips RA, Thompson DR, Waldron S, Furness RW (2000) Variability in mercury concentrations of great skuas Catharacta skua: the influence of colony, diet and trophic status inferred from stable isotope signatures. Mar Ecol Prog Ser 195:261-268.

Bond AL (2010) Relationships between stable isotopes and metal contaminants in feathers are spurious and biologically uninformative. Environ Pollut 158:1182–1184

Bourgeois K, Dromzee D (2014) Moulting strategies of the Yelkouan Shearwater *Puffinus yelkouan* during the breeding season. J Ornithol 155:265–271

Burger J, Gochfeld M (1997) Risk, mercury levels, and birds: relating adverse laboratory effects to field biomonitoring. Environ Res 75:160-172

Carey MJ (2011) Sexual size dimorphism, within-pair comparisons and assortative mating in the short-tailed shearwater (*Puffinus tenuirostris*). Notornis 58: 8-16

Carey MJ, Phillips RA, Silk JRD, Shaffer SA (2014) Trans-equatorial migration of Short-tailed Shearwaters revealed by geolocators. Emu 114:352–359

González-Solís J, Smyrli M, Militão T, Gremillet D, Tveraa T, Phillips RA, Boulinier T (2011) Combining stable isotope analyses and geolocation to reveal kittiwake migration. Mar Ecol Prog Ser 435, 251-261

200 Hedd A, Montevecchi WA, Otley H, Phillips RA, Fifield DA (2012)
201 Trans-equatorial migration and habitat use by sooty shearwaters *Puffinus*
202 *griseus* from the South Atlantic during the nonbreeding season. Mar Ecol
203 Prog Ser 449:277–290
204 Jaeger I, Hop H, Gabrielsen GW (2009) Biomagnification of mercury in selected
205 species from an Arctic marine food web in Svalbard. Sci Total Environ
206 407:4744–4751
207 Laurier FJG, Mason RP, Whalin GAG (2004) Mercury distribution in the North
208 Pacific ocean–20 years of observations. Mar Chem 90:3–19
209 Leat EHK, Bourgeon S, Magnusdottir E, Gabrielsen GW, Grecian WJ, Hanssen SA,
210 Olafsdottir K, Petersen A, Phillips RA, Strøm H, Ellis S, Fisk AT, Bustnes JO,
211 Furness RW, Borgå K (2013) Influence of wintering area on persistent organic
212 pollutants in a breeding migratory seabird. Mar Ecol Prog Ser 491, 277-293.
213 Marchant S, Higgins PJ (1990) Handbook of Australian, New Zealand and
214 Antarctic Birds. Volume 1: Ratites to ducks: Part A: Ratites to petrels.
215 Melbourne, Oxford University Press, pp 632–644
216 Phillips RA, Croxall JP, Silk JRD, Briggs DR (2008) Foraging ecology of
217 albatrosses and petrels from South Georgia: two decades of insights from
218 tracking technologies. Aquat Conserv Mar Freshw Ecosyst 17:6–21
219 Ramos R, González-Solís J, Forero MG, Moreno R, Gomez-Diaz E, Ruiz X,
220 Hobson KA (2009) The influence of breeding colony and sex on mercury,
221 selenium and lead levels and carbon and nitrogen stable isotope signatures in
222 summer and winter feathers of *Calonectris* shearwaters. Oecologia
223 159:345–354

224 Ricca MK, Miles AK, Anthony RG (2008) Sources of organochlorine
225 contaminants and mercury in seabirds from the Aleutian archipelago of
226 Alaska: inferences from spatial and trophic variation. Sci Total Environ
227 406:308–323
228

FIGURE CAPTIONS

Fig. 1 Kernel density map of Short-tailed Shearwaters with different distributions during the non-breeding period: (a) WEST (14 birds), (b) MIX (5 birds), and (c) EAST (9 birds). The kernel density contours represent the proportions of the overall kernel density surface from the highly utilized, core area (black: 25%) to the periphery of the winter distribution (light grey: 95%).

Fig. 2 Mercury (Hg) concentration (in μg/g dry weight) of the outermost primary feather of Short-tailed Shearwaters that spent the non-breeding period in the southern Okhotsk Sea and northern Japan Sea (WEST, 14 birds), in the eastern Bering Sea and around the Aleutian Islands (EAST, 9 birds), or that moved from WEST to EAST (MIX, 5 birds).



(a) WEST
(b) MIX
(c) EAST
120°E
140°E
160°E
180°
160°W
140°W
120°W
70°N
50°N
30°N
0
3,000
km
Kernel density contour (%)
0
25
50
75
95

Fig. 1




Fig. 2