
HOKKAIDO UNIVERSITY

Title	Increased Bacterial Load and Expression of Antimicrobial Peptides in Skin of Barrier-Deficient Mice with Reduced Cancer Susceptibility
Author(s)	Natsuga, Ken; Cipolat, Sara; Watt, Fiona M.
Citation	Journal of investigative dermatology, 136(1), 99-106 https://doi.org/10.1038/JID.2015.383
Issue Date	2016-01
Doc URL	https://hdl.handle.net/2115/62619
Rights(URL)	https://creativecommons.org/licenses/by/4.0/
Type	journal article
File Information	mmc1.pdf, Supplementary Figures


Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

Supplementary Figures




Supplementary Figure 1. Quantification of Ly6G+ dermal cells. Number of Ly6G+ cells per mm^2 dermis in WT, EPI-/-, antibiotics-treated EPI-/- and flora-deficient EPI-/- skin. Data are means ± SEM from at least 4 mice per group. No statistically differences were found between groups.

Supplementary Figure 2



Supplementary Figure 2. AMP expression in EPI-/- epidermis. (A, B) Immunofluorescence labelling of Camp (**A**) and β3 defensin, encoded by Defb3, (**B**) in WT, EPI-/- and flora deficient EPI-/- skin (green fluorescence, with red propidium iodide nuclear counter-stain). Scale bars: 20 μm.

Supplementary Figure 3



Supplementary Figure 3. Serpina1b expression in WT epidermis treated with antibiotics. qRT-PCR of Serpina1b in WT and antibiotic-treated WT epidermis. Data are means ± SEM from 4 mice per group. No statistically significant differences were found between groups.