



HOKKAIDO UNIVERSITY

Title	Molecular weight-dependent degradation of D-lactate-containing polyesters by polyhydroxyalkanoate depolymerases from <i>Variovorax</i> sp C34 and <i>Alcaligenes faecalis</i> T1
Author(s)	Sun, Jian; Matsumoto, Ken'ichiro; Tabata, Yuta et al.
Citation	Applied microbiology and biotechnology, 99(22), 9555-9563 https://doi.org/10.1007/s00253-015-6756-1
Issue Date	2015-11
Doc URL	https://hdl.handle.net/2115/63400
Rights	The final publication is available at link.springer.com
Type	journal article
File Information	Matsumo-AMB(99)suppl. file.pdf



Supporting information

Molecular weight-dependent degradation of D-lactate-containing polyesters by polyhydroxyalkanoate depolymerases from *Variovorax* sp. C34 and *Alcaligenes faecalis* T1

Jian Sun¹, Ken'ichiro Matsumoto¹, Yuta Tabata^{2,3}, Ryosuke Kadoya¹, Toshihiko Ooi^{1,4}, Hideki Abe^{2,3} and Seiichi Taguchi^{1,4,*}

¹Division of Biotechnology and Macromolecular Chemistry, Graduate School of Engineering, Hokkaido University, N13-W8, Kita-ku, Sapporo 060-8628, Japan,

²Department of Innovative and Engineering Materials, Tokyo Institute of Technology, 4259 Nagatsuta, Midori-ku, Yokohama, Kanagawa 226-8502, Japan

³Bioplastic Research Team, Biomass Engineering Program Cooperation Division, RIKEN Center for Sustainable Resource Science, 2-1, Hirosawa, Wako-shi, Saitama 351-0198, Japan

⁴CREST, JST, 4-1-8 Honcho, Kawaguchi, Saitama, 332-0012, Japan

*Corresponding author

staguchi@eng.hokudai.ac.jp

TEL/FAX +81-117066610

Figure S1 ^1H NMR analysis of chemically synthesized trimers. (a) 3HB-LA-LA-methyl, (b) LA-3HB-LA-methyl, (c) LA-LA-3HB-methyl.

