



HOKKAIDO UNIVERSITY

Title	Catalytic Depolymerization of Chitin with Retention of N-Acetyl Group
Author(s)	Yabushita, Mizuho; Kobayashi, Hirokazu; Kuroki, Kyoichi et al.
Citation	Chemsuschem, 8(22), 3760-3763 https://doi.org/10.1002/cssc.201501224
Issue Date	2015-11-23
Doc URL	https://hdl.handle.net/2115/63650
Rights	This is the accepted version of the following article: http://onlinelibrary.wiley.com/doi/10.1002/cssc.201501224/abstract , which has been published in final form at Chemsuschem, Vol.8(22), November 23, 2015. This article may be used for non-commercial purposes in accordance with the Wiley Self-Archiving Policy [olabout.wiley.com/WileyCDA/Section/id-820227.html].
Type	journal article
File Information	CSSC_manuscript_150909hk.pdf



Catalytic Depolymerization of Chitin with Retention of *N*-Acetyl Group**

Mizuho Yabushita, Hirokazu Kobayashi,* Kyoichi Kuroki, Shogo Ito, and Atsushi Fukuoka*

Abstract: Chitin, a polymer of *N*-acetylglucosamine units with β -1,4-glycosidic linkages, is the most abundant marine biomass. Chitin monomers containing *N*-acetyl groups are useful precursors to various fine chemicals and medicines. However, the selective conversion of robust chitin to *N*-acetylated monomers currently requires a large excess of acid or a long reaction time, which has limited its application. We demonstrate a fast catalytic transformation of chitin to monomers with retention of *N*-acetyl group by combining mechanochemistry and homogeneous catalysis. Mechanical-force-assisted depolymerization of chitin with a catalytic amount of H_2SO_4 gave soluble short-chain oligomers. Subsequent hydrolysis of the ball-milled sample provided *N*-acetylglucosamine in 53% yield, and methanolysis afforded 1-*O*-methyl-*N*-acetylglucosamine in yields of up to 70%. Our process can greatly reduce the use of acid compared to the conventional process.

The transformation of biomass to value-added chemicals with particular functional groups is an attractive alternative to multistep functionalization of hydrocarbons from fossil fuels.^[1] The most abundant terrestrial and marine biomasses are cellulose and chitin, respectively. Notably, chitin is a characteristic biopolymer consisting of nitrogen-containing monomer units, *N*-acetylglucosamine (GlcNAc), with β -1,4-glycosidic linkages (Scheme 1). It is therefore an attractive renewable feedstock for organic nitrogen compounds.^[2] However, unlike cellulose conversion, in which thermal and mechanocatalytic depolymerization can both yield monomers,^[3] chitin requires selective depolymerization of glycosidic bond to prevent the loss of *N*-acetyl group for the wide application.

Hydrolysis of the glycosidic bonds of chitin gives GlcNAc, which is useful as a biologically active agent, an ingredient for cosmetics, and a raw material for *N*-containing organic compounds.^[4] Meanwhile, methanolysis of chitin can afford 1-*O*-methyl-*N*-acetylglucosamine (denoted as MeGlcNAc), which inhibits hemagglutination, and thus can be used for suppression of influenza and cancer.^[5] MeGlcNAc is also applicable to the

synthesis of biodegradable polyesters and polyamides,^[6] organocatalysts,^[7] ligands,^[8] and gelators.^[9] Additionally, MeGlcNAc is a useful molecule for further transformations in glycoscience, since the hemiacetal group is protected by a methyl group.^[10]

Despite a great deal of effort in various fields, efficient conversion of chitin to *N*-acetylated monomers remains a challenge. In biochemistry, chitin is depolymerized by enzymes to form GlcNAc in yields of 64–77% under ambient conditions; however, it takes 10 days for the reaction to be completed due to low catalytic activity.^[11] Conventional chemical/mechanical pretreatments slightly improved the reactivity of chitin by the amorphization, but a harsh pretreatment (supercritical water with converge milling) was necessary to accelerate the reaction.^[12] In synthetic organic chemistry, HCl provides GlcNAc in ca. 65% yield, but only when HCl is used in large excess (i.e., the molar ratio of HCl to GlcNAc units in chitin = 100) and at high concentrations of 15–36%.^[4a, 12a] Therefore, the serious corrosiveness of HCl becomes an issue, and a huge amount of acidic waste is produced. Dilution of HCl in the depolymerization reaction of chitin results in low selectivity for GlcNAc.^[13] Moreover, the use of acid results in deacetylation of chitin,^[13] but preservation of the *N*-acetyl group is essential for high physiological activity and wide application of the resulting chitin monomers.^[14] The synthesis of MeGlcNAc has not been studied to the same extent as that of GlcNAc; in fact, its production from chitin has not yet been reported. Fischer glycosidation of GlcNAc has been necessary to obtain MeGlcNAc.^[15] Herein, we report catalytic depolymerization of chitin to GlcNAc and MeGlcNAc (Scheme 1) in a process which reduces the use of acid by 99.8% and is completed within one day (including all handling).

The first key step in our reaction is mechanocatalytic conversion of chitin to soluble oligomers. It is known that mechanical force enhances chemical reactions under mild conditions.^[16] Chitin was impregnated with a small amount of H_2SO_4 [substrate to catalyst ratio (S/C) = 8.1], and the resulting mixed solid was ball-milled at 500 rpm for 6 h to achieve mechanocatalytic hydrolysis (Scheme 1, product name: Chitin- H_2SO_4 -BM). The water used for hydrolysis should be physisorbed water (1.5 wt%) in chitin. The process quantitatively converted chitin to soluble compounds such as GlcNAc (4.7% yield), (GlcNAc)₂ (7.8% yield), (GlcNAc)₃ (11% yield), (GlcNAc)₄ (9.7% yield), and (GlcNAc)₅ (8.6% yield), where (GlcNAc)_{*n*} denotes an oligomer consisting of *n* GlcNAc units (Figure 1, Chitin- H_2SO_4 -BM). Other products were longer and branched oligomers, as determined by liquid chromatography/mass spectroscopy (LC/MS) and nuclear magnetic resonance (NMR) spectroscopy (Figures S1 and S2). It is known that side chains improve the solubility of oligomers.^[16] Chitin oligomers are useful as antitumor, immunoenhancing, and antimicrobial agents^[17] as well as useful precursors of monomers (vide infra).


[*] Dr. M. Yabushita, Dr. H. Kobayashi, Prof. Dr. A. Fukuoka
Catalysis Research Center, Hokkaido University
Kita 21 Nishi 10, Kita-ku, Sapporo, Hokkaido 001-0021 (Japan)
E-mail: kobayashi.hi@cat.hokudai.ac.jp (H.K.),
fukuoka@cat.hokudai.ac.jp (A.F.)

K. Kuroki, S. Ito
Graduate School of Chemical Sciences and Engineering, Hokkaido University

Kita 13 Nishi 8, Kita-ku, Sapporo, Hokkaido 060-8628 (Japan)

[**] We would like to thank Dr. Y. Yamakoshi (Hokkaido Research Organization) for the LC/MS analysis and Dr. A. Shrotri for useful discussion. This work was supported by a Grant-in-Aid for Young Scientists (A) (KAKENHI, 26709060) from the Japan Society for the Promotion of Science (JSPS) and a Grant-in-Aid for the JSPS Fellows.

Supporting information for this article is given via a link at the end of the document.



Scheme 1. Two-step depolymerization of chitin to *N*-acetylated monomers, catalyzed by H_2SO_4 . The first step is mechanocatalytic depolymerization of chitin to water-soluble oligomers, and the second is solvolysis of the oligomers to *N*-acetylated monomers.





Figure 1. Mechanocatalytic depolymerization of chitin. Dissolution conditions: chitin weight 406 mg; distilled water 40 mL; 298 K.

In controlled experiments (Figure 1), pristine chitin contained no soluble compounds, while chitin that had been ball-milled in the absence of H_2SO_4 (Chitin-BM) afforded soluble compounds in 5.2% yield, with the chitin becoming amorphous (Figure S3), and H_2SO_4 -impregnated chitin that had been aged at 298 K for 6 h (Chitin- H_2SO_4) had a soluble fraction of only 3.7%. Hence, both ball-milling and H_2SO_4 are necessary for depolymerization of chitin. To examine the synergy between mechanical force and H_2SO_4 , chitin was ball-milled, impregnated with H_2SO_4 , and aged at 298 K for 6 h (Chitin-BM- H_2SO_4); this sample had a soluble fraction of only 7.8%. Therefore, mechanical force was only effective for the depolymerization of chitin in the presence of H_2SO_4 .

Notably, acetyl groups in the GlcNAc units were preserved after the mechanocatalytic reaction, since Chitin- H_2SO_4 -BM consisted of *N*-acetylated oligomers and did not contain acetic acid. This result is specific to the mechanocatalysis, as the conventional hydrolysis using a small amount of acid results in significant removal of *N*-acetyl group.^[13] In the mechanocatalytic reaction (Scheme 2), we propose that amorphization of chitin

first occurs by the ball-milling, which enhances the accessibility of acid to chitin due to lower steric hindrance.^[18] Then, mechanical force assists the cleavage of the connecting points of polymers, in which polymers are subjected to macro-scale power as tensile stress.^[19] Thus, the use of mechanical force resulted in selective activation of glycosidic bonds connecting the GlcNAc units rather than amide bonds in the acetamide



Scheme 2. Proposed schematic of mechanochemical depolymerization. The mechanical force first amorphizes chitin, which increases the accessibility of acid to the substrate. The main chain is subjected to strong mechanical force, which leads to selective dissociation of the connecting points of the chain (glycosidic bonds), while the side chains (acetamide groups) do not dissociate.

Table 1. Solvolysis of chitin samples to *N*-acetylated monomers.^[a]

Entry	Sample	S/C	Reaction temperature [K]	Solvent	Product ^[b] [%]		
					GlcNAc	MeGlcNAc	Other soluble products
1	Chitin-H ₂ SO ₄ -BM	8.1	463	Water	32	–	68
2	Chitin-H ₂ SO ₄ -BM	2.0	443	Water	53	–	47
3	Chitin	– ^[c]	463	Methanol	–	0.0	0.0
4	Chitin-H ₂ SO ₄	8.1	463	Methanol	–	4.0	1.7
5	Chitin-BM	– ^[c]	463	Methanol	–	0.0	2.2
6	Chitin-BM-H ₂ SO ₄	8.1	463	Methanol	–	16	4.6
7	Chitin-H ₂ SO ₄ -BM	8.1	463	Methanol	–	68	27
8	Chitin-H ₂ SO ₄ -BM	4.1	463	Methanol	–	70	28

[a] Common reaction conditions: chitin weight 406 mg; solvent 40 mL; rapid heating-cooling conditions (time course is shown in Figure S4). [b] Based on moles of GlcNAc units. [c] No H₂SO₄ was impregnated on chitin.

groups hanging from the units. This is the remarkable merit of mechanocatalysis in the selective depolymerization of chitin. The merit is different from that in the conversion of simple sugar polymers like cellulose that have only one functional group to be hydrolyzed (glycosidic bond).^[16]

To produce GlcNAc, Chitin-H₂SO₄-BM was further hydrolyzed in water under rapid heating-cooling conditions (Table 1). The catalyst for hydrolysis was H₂SO₄ contained in the sample (S/C = 8.1), which gave a 32% yield of GlcNAc at 463 K in 1 h (entry 1). The yield of GlcNAc was improved to 53% when the amount of H₂SO₄ was increased to S/C = 2.0 and the reaction temperature was lowered to 443 K (entry 2). However, the formation of by-products was still observed in these reactions because of the degradation of GlcNAc, which is thermally unstable in water; in a stability test of GlcNAc, only 17% of GlcNAc was recovered after being subjected to the same heating conditions (Table S1, entry S1). Since the generation of acetic acid was limited (<1%), the degradation must have resulted from the conversion of hemiacetal groups in our rapid reaction.

The second key point in our reaction is to overcome the stability issue by using methanolysis instead of hydrolysis, as the product MeGlcNAc does not contain a hemiacetal group. Indeed, 94% of the MeGlcNAc was recovered after heat treatment at 463 K (Table S1, entry S2). It is notable that the advantage of MeGlcNAc is not only the good stability but also the wide applications in various fields as described above.^[5–10] Recently, Yan et al. demonstrated that alcoholysis of chitin was possible in the presence of H₂SO₄ (S/C = 0.9), although the deacetylated product [i.e., 1-*O*-(2-hydroxyethyl)-glucosamine] was obtained predominantly (24% yield) due to the harsh conditions required for direct depolymerization of the robust chitin.^[20] Therefore, we can expect selective synthesis of

MeGlcNAc from the reactive chitin oligomers contained in Chitin-H₂SO₄-BM.

Chitin samples were subjected to methanolysis at 463 K to demonstrate selective synthesis of MeGlcNAc. The reaction of Chitin-H₂SO₄-BM provided MeGlcNAc in 68% overall yield, even when only a small amount of H₂SO₄ was used (S/C = 8.1) (entry 7). The ratio of α - to β -anomers in MeGlcNAc was 5.0, and the high stereoselectivity is favorable for further applications (the formation mechanism of the α -anomer is discussed in Figure S5). The turnover number (TON) of H₂SO₄ for MeGlcNAc production was 5.6, showing that the acid acted as a catalyst. This result was in sharp contrast to those obtained from conventional systems, which require a large excess of HCl (TON < 0.01) to obtain the *N*-acetylated monomer.^[4a] With regard to by-products, only a trace amount of methyl acetate (0.90%) was detected by gas chromatography. Almost no deacetylation occurred during methanolysis under our conditions. The yield of methanol-insoluble humins was 4.7 wt% based on the weight of chitin used. The yield of MeGlcNAc was increased to 70% under optimum conditions (S/C = 4.1) (entry 8). In contrast, methanolysis of other samples—pristine chitin, Chitin-H₂SO₄, Chitin-BM, and Chitin-BM-H₂SO₄—gave low yields of MeGlcNAc ($\leq 16\%$; entries 3–6). These results clearly show that mechanocatalytic depolymerization of chitin prior to methanolysis is essential for high-yield synthesis of MeGlcNAc. The reactivity of chitin is dramatically different from that of cellulose; the methanolysis of cellulose gives a good yield of methylglucose with no pretreatment.^[21]

In summary, we demonstrated the catalytic conversion of chitin to *N*-acetylated monomers, notably MeGlcNAc. The process consists of mechanocatalytic depolymerization of chitin to soluble oligomers, followed by solvolysis to monomers. Mechanical force was found to be essential to achieving selective catalytic depolymerization with retention of *N*-acetyl

group, demonstrating the remarkable merit of mechanocatalysis. The second thermocatalytic methanolysis converted oligomers to the *N*-acetylated monomer MeGlcNAc in 68–70% yield, owing to the improved reactivity of the substrate and the good stability of the product. In the overall process, the amount of acid used for the reaction was drastically reduced (99.8%) compared with the conventional process. Our method overcomes critical issues in conventional systems for chitin depolymerization, and opens the door to various applications of chitin-based compounds.


Keywords: biomass • catalysis • chitin • hydrolysis • mechanochemistry

- [1] a) A. Corma, S. Iborra, A. Velty, *Chem. Rev.* **2007**, *107*, 2411–2502; b) F. M. Kerton, Y. Liu, K. W. Omari, K. Hawboldt, *Green Chem.* **2013**, *15*, 860–871; c) H. Kobayashi, A. Fukuoka, *Green Chem.* **2013**, *15*, 1740–1763.
- [2] X. Chen, N. Yan, *Catal. Surv. Asia* **2014**, *18*, 164–176.
- [3] a) J. S. Luterbacher, D. M. Alonso, J. A. Dumesic, *Green Chem.* **2014**, *16*, 4816–4838; b) A. Fukuoka, P. L. Dhepe, *Angew. Chem. Int. Ed.* **2006**, *45*, 5161–5163; *Angew. Chem.* **2006**, *118*, 5285–5287; c) Y. Su, H. M. Brown, X. Huang, X. Zhou, J. E. Amonette, Z. C. Zhang, *Appl. Catal. A: Gen.* **2009**, *361*, 117–122.
- [4] a) J.-K. Chen, C.-R. Shen, C.-L. Liu, *Mar. Drugs* **2010**, *8*, 2493–2516; b) K. W. Omari, L. Dodot, F. M. Kerton, *ChemSusChem* **2012**, *5*, 1767–1772; c) Y. Wang, C. M. Pedersen, T. Deng, Y. Qiao, X. Hou, *Bioresour. Technol.* **2013**, *143*, 384–390; d) M. Osada, K. Kikuta, K. Yoshida, K. Totani, M. Ogata, T. Usui, *Green Chem.* **2013**, *15*, 2960–2966; e) Y. Ohmi, S. Nishimura, K. Ebitani, *ChemSusChem* **2013**, *6*, 2259–2262; f) X. Chen, S. L. Chew, F. M. Kerton, N. Yan, *Green Chem.* **2014**, *16*, 2204–2212.
- [5] a) F. Tabary, J. Font, R. Bourrillon, *Arch. Biochem. Biophys.* **1987**, *259*, 79–88; b) N. Kochibe, K. L. Matta, *J. Biol. Chem.* **1989**, *264*, 173–177.
- [6] S. Fujii, Y. Kondo, M. Matsui, K. Ichihashi, JP Patent 5 178 904, **1993**.
- [7] J. Agarwal, R. K. Peddinti, *Eur. J. Org. Chem.* **2012**, 6390–6406.
- [8] T. V. RajanBabu, T. A. Ayers, G. A. Halliday, K. K. You, J. C. Calabrese, *J. Org. Chem.* **1997**, *62*, 6012–6028.
- [9] N. Goyal, S. Cheuk, G. J. Wang, *Tetrahedron* **2010**, *66*, 5962–5971.
- [10] K. M. Koeller, C.-H. Wong, *Nat. Biotechnol.* **2000**, *18*, 835–841.
- [11] H. Sashiwa, S. Fujishima, N. Yamano, N. Kawasaki, A. Nakayama, E. Muraki, K. Hiraga, K. Oda, S. Aiba, *Carbohydr. Res.* **2002**, *337*, 761–763.
- [12] a) J. A. Bohlmann, D. O. Schisler, K.-O. Hwang, J. P. Henning, J. R. Trinkle, T. B. Anderson, J. D. Steinke, A. Vanderhoff, US Patent 6 693 188, **2004**; b) I. Karube, T. Morita, US Patent 5 262 310, **1993**; c) M. Osada, C. Miura, Y. S. Nakagawa, M. Kaihara, M. Nikaido, K. Totani, *Carbohydr. Polym.* **2013**, *92*, 1573–1578.
- [13] a) A. Einbu, K. M. Vårum, *Biomacromolecules* **2007**, *8*, 309–314; b) N. Gandhi, J. K. Laidler, US Patent 6 486 307, **2002**.
- [14] a) A. R. Shikhman, K. Kuhn, N. Alaaeddine, M. Lotz, *J. Immunol.* **2001**, *166*, 5155–5160; b) L. I. Álvarez-Añorve, M. L. Calcagno, J. Plumbridge, *J. Bacteriol.* **2005**, *187*, 2974–2982.
- [15] J. Yoshimura, H. Ando, Y. Takahashi, H. Ono, T. Sato, *Nippon kagaku zasshi* **1964**, *85*, 142–145.
- [16] a) S. L. James, C. J. Adams, C. Bolm, D. Braga, P. Collier, T. Friščić, F. Grepioni, K. D. M. Harris, G. Hyett, W. Jones, A. Krebs, J. Mack, L. Maini, A. G. Orpen, I. P. Parkin, W. C. Shearouse, J. W. Steedk, D. C. Waddell, *Chem. Soc. Rev.* **2012**, *41*, 413–447; b) Q. Zhang, F. Jérôme, *ChemSusChem* **2013**, *6*, 2042–2044; c) R. Rinaldi, *Angew. Chem. Int. Ed.* **2014**, *53*, 8559–8560; *Angew. Chem.* **2014**, *126*, 8699–8701; d) G.-W. Wang, *Chem. Soc. Rev.* **2013**, *42*, 7668–7700; e) R. G. Blair in *Production of Biofuels and Chemicals with Ultrasound* (Eds.: Z. Fang, R. L. Smith, Jr., X. Qi), Springer, Berlin, **2014**, pp. 269–288; f) S. M. Hick, C. Griebel, D. T. Restrepo, J. H. Truitt, E. J. Buker, C. Bylda, R. G. Blair, *Green Chem.* **2010**, *12*, 468–474; g) N. Meine, R. Rinaldi, F. Schüth, *ChemSusChem* **2012**, *5*, 1449–1454; h) R. Carrasquillo-Flores, M. Kåldström, F. Schüth, J. A. Dumesic, R. Rinaldi, *ACS Catal.* **2013**, *3*, 993–997; i) A. Shrotri, L. K. Lambert, A. Tanksale, J. Beltrami, *Green Chem.* **2013**, *15*, 2761–2768; j) S. Liu, Y. Okuyama, M. Tamura, Y. Nakagawa, A. Imai, K. Tomishige, *ChemSusChem* **2015**, *8*, 628–635.
- [17] F. Shahidi, J. K. V. Arachchi, Y.-J. Jeon, *Trends Food Sci. Tech.* **1999**, *10*, 37–51.
- [18] T. Hama, H. Ueda, A. Kouchi, N. Watanabe, H. Tachikawa, *J. Phys. Chem. Lett.* **2014**, *5*, 3843–3848.
- [19] a) M. K. Beyer, H. Clausen-Schaumann, *Chem. Rev.* **2005**, *105*, 2921–2948; b) D. A. Davis, A. Hamilton, J. Yang, L. D. Cremer, D. V. Gough, S. L. Potisek, M. T. Ong, P. V. Braun, T. J. Martinez, S. R. White, J. S. Moore, N. R. Sottos, *Nature* **2009**, *459*, 68–72.
- [20] Y. Pierson, X. Chen, F. D. Bobbink, J. Zhang, N. Yan, *ACS Sustainable Chem. Eng.* **2014**, *2*, 2081–2089.
- [21] W. Deng, M. Liu, Q. Zhang, X. Tan, Y. Wang, *Chem. Commun.* **2010**, *46*, 2668–2670.

Entry for the Table of Contents

COMMUNICATION

Preserving *N*-acetyl groups: Chitin is depolymerized to *N*-acetylated monomers in good yields by combining mechanochemistry and homogeneous catalysis. The first mechanocatalytic hydrolysis of chitin provides water-soluble oligomers, which are more reactive than chitin. The subsequent solvolysis of the oligomers easily produces monomers without deacetylation.



Mizuho Yabushita, Hirokazu Kobayashi,* Kyoichi Kuroki, Shogo Ito, and Atsushi Fukuoka*

Page No. – Page No.

Catalytic Depolymerization of Chitin with Retention of *N*-Acetyl Group