HOKKAIDO UNIVERSITY

Title	天蛾科の二新種
Author(s)	松村, 松年
Citation	札幌博物学会会報, 11(3), 119-120
Issue Date	1930-11-30
Doc URL	https://hdl.handle.net/2115/63870
Type	journal article
File Information	Vol.11No.3_001.pdf

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

TWO NEW SPECIES OF SPHINGID-MOTHS

BY

SHÔNEN MATSUMURA

(With 2 Text Figures)

天蛾科の二新種

松村松年

1. *Oxyambulyx takamukui* n. sp. (Fig. 1)

♂. Primaries yellowish brown, with a light shade of purple; at the base with 2 and at the costa with 3 velvety fuscous spots; the medial line double, interrupted, fuscous, geniculated at the medial vein; the discocellulars black, geniculated, having a paler line in the middle; the postmedial line broad, olivaceous, defined by a black band internally; the space from this line to the bow-like marking at the margin pale yellowish, scattering fuscous spots and lines within it; at the inner side of the tornus with a large velvety black patch. Secondaries yellowish, with about 4 black irregular bands, those of the outer 2 being much narrower, zigzac, the discoidal spot reddish yellow; the posterior margin broadly black, internally defined by a velvety black



Fig. 1.

Oxyambulyx takamukui n. sp.

[Transact. Sapporo Nat. Hist. Soc., Vol. XI, Pt. 3, 1930]

band. Body above dark brown, abdomen at the sides of the 3rd and 4th segments and at the middle of the 4th with each a fuscous spot; below yellowish.

Expanse—115 mm.

Hab.—Formosa; one male specimen was collected by the late T. TAKAMUKU at Horisha.

The nearest ally: *O. panopus* WK.

2. *Macroglossum palauensis* n. sp. (Fig. 2)

♂. Primaries dark grey, at the base with about 4 obsolete fuscous bands; in the middle with 2 fuscous bands, that of the inner being somewhat incurved at the hind angle of cell; the apical mark broadly fuscous, intersected by the longitudinal veins. Secondaries velvety black, with an orange-yellow band somewhat constricted in the middle, not extending to the apex.



Fig. 2.

Macroglossum palauensis n. sp.

This resembles somewhat *M. iwasakii* MATS., but may easily be distinguished in having a narrower yellowish band to the secondaries.

Expanse—52 mm.

Hab.—Palau; one male specimen was collected by Mr. S. UCHIYAMA.

摘　　要

今回二種の天蛾を發見したれば爰に發表せん。その和名は下の如し。

1. *Oxyambulyx takamukui* MATS. (n. sp.)　タカムクホソバスズメ
2. *Macroglossum palauensis* MATS. (n. sp.)　パラウホウジヤク

前者は印度産のパノープス (*O. panopus* WK.) に、後者は沖繩産のイハサキホウジヤク (*M. iwasakii* MATS.) に似る。