



Title	センブリ属の一新種
Author(s)	本田, 正次; 館脇, 操
Citation	札幌博物学会会報, 14(3), 192-192
Issue Date	1936-06-30
Doc URL	https://hdl.handle.net/2115/64181
Type	journal article
File Information	Vol.14No.3_006.pdf



BUCHI, VIII. 28, 1890; KINGO TOGASHI, VIII. 6, 1935—type in Herb. Fac. Agr. Hokkaido Imp. Univ.).

Remarks. The present species is closely related to *Calanthe reflexa* MAXIM., from which it differs by the shape of the labellum and sepals, and also by the appressedly hairy character of the under surface of the leaves.

A NEW SPECIES OF SWERTIA

BY

MASAJI HONDA AND MISAO TATEWAKI

(本田正次・館脇操)

Swertia crysantha HONDA ET TATEWAKI, sp. nov.

Radix simplex exilis. Caulis simplex vel subsimplex erectus glaber angulato-alatus. Folia ovato-lanceolata sessilia vel brevissime petiolata utrinque glabra vulgo 3-5 nervia apice obtusiuscula basi rotundata vel leviter cordata 2-2.5 cm longa 0.8-1.5 cm lata; inferiora minora oblongo-spathulata vel ovato-lanceolata. Flores minusculi omnes quadrifidi. Pedunculi axillares et terminales umbellato-fastigiati. Calyx fere usque ad basin 4-partitus, lobis lineari-lanceolatis 4-5 mm longis 0.5-1.5 mm latis. Corolla rotata 4-partita expansa 0.5-1 cm in diametro decidua. Corollae segmenta lutea vel flava vix basin cohaerentia patentia elliptica 4-5.5 mm longa 1.5-2 mm lata, intus infra medio foveola et sub fovea papillosa; fovea solitaria quadrata, marginibus pilis tuberculosus pauci-ciliata.

NOM. JAP. *Yakeishi-senburi*, n. n. (H. IWABUCHI).

HAB. *Honshu*: Mt. Yakeishi, Prov. Rikuchu (H. IWABUCHI, VIII. 9, 1935).

摘 要

センブリ属の新種

本種はヤケイシセンブリと稱し、岩淵初郎氏が陸中焼石岳に於て發見新和名を提稱せる新種にして、チシマセンブリに類似せるも、花冠裂片の黄色なると腺体の位置形狀を異にす。