HOKKAIDO UNIVERSITY

Title	Caspase-dependent cleavage regulates protein levels of Epstein-Barr virus-derived latent membrane protein 1
Author(s)	Togi, Sumihito; Hatano, Yosuke; Muromoto, Ryuta et al.
Citation	FEBS Letters, 590(6), 808-818 https://doi.org/10.1002/1873-3468.12119
Issue Date	2016-03
Doc URL	https://hdl.handle.net/2115/64895
Rights	This is the peer reviewed version of the following article: Caspase-dependent cleavage regulates protein levels of Epstein-Barr virus-derived latent membrane protein 1, FEBS Letters590(6), pp808-818, 2016, which has been published in final form at 10.1002/1873-3468.12119. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.
Type	journal article
File Information	suppl.pdf, Summplementary information

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

Summplementary information

Supplementary Figure S1. **Effect of MG132 on TNF-α-induced IκBα degradation in HeLa cells.** HeLa cells in a 24 well plate were treated with DMSO or MG132 (10 μM) for 12h, followed by TNF-α (10 ng/ml) stimulation for the indicated periods. Cells were then lysed, and an aliquot of total cell lysate (TCL) (1%) was blotted with anti-IκBα (Santa Cruz Biotechnology, Santa Cruz, CA) or anti-actin antibody.

Supplemental Figure 1