



HOKKAIDO UNIVERSITY

Title	Occurrence of the Kuril Harbour Seal (<i>Phoca vitulina</i>) at a small fixed fishing net in Akkeshi Bay, Hokkaido, Japan
Author(s)	Kobayashi, Yumi; Kobayashi, Mari; Sakurai, Yasunori et al.
Citation	International Journal of Agricultural Policy and Research, 5(1), 18-25 https://doi.org/10.15739/IJAPR.17.003
Issue Date	2017-01-21
Doc URL	https://hdl.handle.net/2115/64923
Rights(URL)	https://creativecommons.org/licenses/by/4.0/
Type	journal article
File Information	Occurance seal.pdf





Original Research Article

Occurrence of the Kuril Harbour Seal (*Phoca vitulina*) at a small fixed fishing net in Akkeshi Bay, Hokkaido, Japan

Received 11 November, 2016

Revised 15 January, 2017

Accepted 18 January, 2017

Published 21 January, 2017

**Yumi Kobayashi^{1,2 *},
Mari Kobayashi³,
Yasunori Sakurai¹,
and
Kiyoharu Takada⁴**

¹Graduate School of Fisheries Sciences, Hokkaido University, 3-1-1 Minato-cho, Hakodate, Hokkaido, 041-8611, Japan.

²Current Address: Laboratory of Animal Ecology, Research Faculty of Agriculture, Hokkaido University, Kita-9, Nishi-9, Kita-ku, Sapporo, Hokkaido, 060-8589, Japan.

³Department of Aqua Bioscience and Industry, Tokyo University of Agriculture, Abashiri, Hokkaido, 099-2493, Japan

⁴Akkeshi town Fisherman, Akkeshi, 088-0875, Japan.

*Corresponding Author Email: kobayumi_kotomasu@hotmail.com

Tel: +81-138-40-8863

Humans frequently come into conflict with marine mammals that compete for the same resources, such as seals exploiting the same areas as industrial fisheries. To develop efficient interventions to minimize the negative impact of seals on fisheries, it is essential to understand the behaviour of seals in the fishery area, such as where and when they occur. The present study characterizes an instance of human-seal conflict, using acoustic telemetry to examine patterns in the occurrence of three Kuril harbour seals (*Phoca vitulina*) around a small fixed set-net in Akkeshi Bay, eastern Hokkaido, Japan, in late spring in 2009. To examine the environmental variables affecting the occurrence of Kuril seals around the net, 3 Kuril seals were captured and tracked using an acoustic monitoring system. Two of the tracked individuals (a male and a female) usually came to the net in the evening, and at high tide, but the seals were never present at times when fishing occurred (04:00–07:00 h), which indicates that they actively avoided human activity in the coastal fishery. Seals did not appear in the net after the fishery season (April), a behavioural characteristic that suggests that these seals were adults that used a nearby haul-out site during the breeding season (May–June). A third individual, an adult male, never occurred near the net. These observations demonstrate the extent to which individual seals vary in their interactions with human activities. Given the current selective removal of seals in the fishery area by controlled killing or driving away, these findings have the potential to inform targeted management intervention to minimize conflict between economic and conservation interests.

Key words: human-seal conflicts, harbour seals, small set-net, Akkeshi Bay -Japan, acoustic telemetry system

INTRODUCTION

The production of seafood for human consumption has led to long-running and widespread conflicts with many wild marine mammals around the world. One type of global human-marine mammal conflict that has recently increased in intensity is that between coastal fisheries and seals, which damage both the catch and the fishing gear (*e.g.*, northern Sweden; Lunneryd et al, 2003). In Japan, the Kuril harbour seal (*Phoca vitulina*, henceforth “Kuril seal”) is resident in eastern Hokkaido all year round. Itoo and Shukunobe (1986) estimated that during the 1940s, the population of Kuril harbour seals that lived along the southeastern coast

of Hokkaido was 1500–4800, based on the observations of Inukai (1942a,b). The population was estimated at only a few hundred in 1970s (*e.g.*, Hayama 1988), and the causes of that decline may have been commercial harvesting of the seals and human activities including coastal fisheries (*e.g.*, Itoo and Wada 1982). In particular, the annual hunt was estimated to be at least 250–300 individual seals (Itoo and Syukunobe 1986). Moreover, bycatch from salmon set-net fisheries has also been reported (*e.g.*, Wada et al., 1991). The commercial harvest ended in the late 1980s, and the harbour seal population has thus been increasing for




Figure 1: The study area: Akkeshi Bay, eastern Hokkaido, Japan, showing 10 m isobath diagram. ●: Kuril seal haul-out sites; ①② seal trapping location, ★: receiver position

almost 30 years (Kobayashi et al, 2014). The second-largest Kuril seal haul-out site in Japan is on Daikoku Island in the Akkeshi area of Hokkaido, where recent observations have identified over 300 seals. In concert with this increasing population, in the 1980s, Itoo and Wada (1982) reported damage caused by seals in this area to fish caught in salmon set-nets and gillnets during the autumn. More recently, damage to fish has also been reported from gill nets and small fixed nets during the spring (March-May) in Akkeshi Bay (Kobayashi et al, 2010).

To develop efficient interventions to reduce the interactions between seals and fisheries, it is essential to understand the behaviour of seals in the fishery area, such as where and when they occur. Direct observation of seal behaviour is not straightforward, with the observation of nocturnal activity at sea being a particular challenge, but biologging methods can be used effectively to overcome these constraints.

In the present study, we use acoustic telemetry to track and characterize the behavioural patterns of Kuril seals in the fishery area in Akkeshi Bay.

MATERIALS AND METHODS

Study area

This study was conducted in the spring (April-June) of 2009 in the Akkeshi area, eastern Hokkaido, Japan (Figure 1). The town of Akkeshi has a population of about 11,000, of which approximately 1,500 people over the age of 15 years were employed in the fishing industry as of 2010 (Akkeshi town 2015). Akkeshi Bay supports many coastal fisheries, including spring/autumn salmon set-nets, small fixed nets and dragnets, as well as production of other marine resources such as Kombu seaweed and shellfish. Usually, a

small fixed net can catch several tens of fish species, but only several dominant fish species (Sweke et al, 2006), and damage to seals can also occur. The fishes caught in this area include sailfin sandfish (*Arctoscopus japonicus*), righteye flounder (Pleuronectidae) Pacific herring (*Clupea pallasii*), icefish (*Salangichthys microdon*), saffron cod (*Eleginus gracilis*) and fluvial sculpin (family Cottidae) (Kobayashi et al, 2013). The spring small set-net used in the present study is generally worked by fishermen once a day, around 04:00–07:00 h, except on Sundays and public holidays. In 2009, the fishing season for small set-nets in Akkeshi Bay was March 21st–May 15th (56 days in total).


Four Kuril seal haul-out sites exist in the Akkeshi area: Daikoku Island and Akkeshi A, B, and C (Figure 1). Daikoku Island is an uninhabited island, with a coastline of 6 km, located near the mouth of Akkeshi Bay. This is the second-largest haul-out site (>300) in Hokkaido, Japan.

Acoustic telemetry monitoring system

Kuril seal behaviour was tracked using an acoustic monitoring system. The transmitter was a V16P-5H unit (VEMCO Ltd, Halifax, Canada), and the receiver a VR2W unit (VEMCO Ltd, Halifax, Canada). During a scan, the receiver listened regularly on each frequency for pinger pulses (69 kHz), and the time interval between detected pulses was recorded (Voegeli et al, 1998).

The transmitter used in this study measured 105×15×15 mm and weighed 52.0 g. Pinger pulses had a frequency of one every 10–30 s (mean 20 s), giving a battery lifespan of approximately 178 days. The receiver had a battery life of approximately 15 months and an average coverage of 300 m.

The receiver was installed at a small set-net for icefish (*Salangichthys microdon*) (net no. 44 in the Tomata area, Akkeshi bay; N43°02' 03", E144°45' 34") (Figure 1)



Number of readings registered by the receiver
at each hour of measurement = day + time




Figure 3: The time interval between each seal's occurrences at the net, presented as a percentage of total records.

(every 60th minute) + tide (cm) + mean wind speed (m/s) + interactions between these variables. The importance of each factor was evaluated by selecting the most parsimonious model based on Akaike's Information Criterion (AIC). All statistical analyses were performed using R, version 3.2.3 (R Development Core Team, 2015).

RESULTS AND DISCUSSION

Seal occurrence around the net

In total, 1,368 pulses were received from individual no. 1 (adult female) over 7 days (April 7th–22nd), and pulses from individual no. 2 (adult male) were detected 870 times over 17 days (April 6th–May 8th). No pulses were recorded from individual no. 3 (adult male), and no records were received from any seal after the fishing season ended (May 15th) (Figure 2). Individuals no. 1 and no. 2 were never recorded at the net at the same time. These two individuals occurred on 28.0% (7 occurrence days/25 days of receiver activity) and 52.0% (13/25 days) of recording days in April, respectively, and on 0.0% (0/26 days) and 15.4% (4/26 days) in May. The total duration of occurrence near the net was 607.2 min (mean stay time per day 86.5 ± 77.4 min) for individual no. 1 and 343.2 min (mean stay time per day 20.1 ± 43.5 min) for individual no. 2, with individual no. 1 having a significantly longer total recording duration (Wilcoxon

signed-rank test, $p < 0.05$). Stay time per visit, however, did not vary between these seals (no. 1: 4.3 ± 0.7 min, $N = 142$ visits; no. 2: 4.2 ± 0.7 min, $N = 81$ visits; KS test, $p > 0.05$). There was no daily variation in stay time for either seal (KS test, $p > 0.05$). The interval between occurrences of the seals around the net was mostly 5 min or less, but occasionally over 3 days passed between occurrences (Figure 3).

In general, foraging habitat selection by harbour seals depends on small-scale habitat differences, but habitat use also differs between individual seals (Tollit et al, 1998).

Individual no. 1 was not recorded after late April, but was only observed at a haul-out site. Although this female seal lost her tag halfway through the study period, it is notable that she showed signs of pregnancy when she was initially caught. In the pre-breeding period, adult female harbour seals make regular trips between the haul-out site and foraging areas, but around the time of breeding, these trips stop as females spend much longer at the haul-out sites (Thompson et al, 1994).

Moreover, haul-out patterns vary between geographical areas and seasons, probably in accordance with regional and seasonal dietary variation as well as differing demands across different phases of the seals' life cycle (Niizuma 1986). Although female no. 1 was only recorded in the fishery area during early April, she spent more time there than individual no. 2.

No records were obtained from individual no. 3, an adult male caught in a different area (Monshizu; individuals no. 1




Figure 4. The time of each seal's occurrence at the net, presented as a percentage of total records

Table 1. Results for the generalized linear mixed model (GLM) explaining the occurrence of seals at the net in relation to fishery catch area and environmental variables.

	Estimate	Std. Error	Z value	Pr (> Z)
(Intercept)	-2.150e+06	2.115e+05	-10.165	< 2e-16 ***
Date	1.070e-01	1.053e-02	10.165	< 2e-16 ***
Time	8.246e+04	9.593e+03	8.595	< 2e-16 ***
Tide	1.283e+04	1.242e+03	10.330	< 2e-16 ***
Wind	2.248e-01	3.874e-02	5.802	6.55e-09***
Tide×Wind	-2.224e-03	4.488e-04	-4.955	7.25e-07***
Date×Time	-4.104e-03	4.757e-04	-8.595	< 2e-16 ***
Date×Tide	-6.835e-04	6.181e-05	-10.330	< 2e-16 ***

and no. 2 were caught in the Tomata area, near where the receiver was deployed). Collectively, this suggests that individual no. 3 was either not feeding around the Tomata fishery area or feeding in a different area. Harbour seals have a lek-like terrestrial mating system, and from the pre-breeding season to the mating season, intense male-male competition has been reported (Hayes et al, 2004), which demands that much be spent at the haul-out site (Niizuma 1986). To further elucidate these various explanations of seal behaviour, a challenge for future studies is to simultaneously deploy many receivers in this area.

Environmental variables affecting occurrence around the net

The best-fit model describing the occurrence time of seals around the net showed that most records were received at high tide levels and during the evening and night (Table 1; Figure 4). Throughout the tracking period, records were never received between 04:00 and 07:00 h, which is the time of highest activity in the fishery, suggesting that the

seals avoided human activity. However, there was no relationship between seal activity and breaks in fishery activity resulting holidays. Some fishermen use the fishery area to repair their nets during holidays; therefore, the holidays might have not actually affected seal behaviour near the nets.

The times at which the two recorded seals occurred most frequently at the net differed (KS test, $p < 0.001$), but seals occurred in the evening at 16:00–18:00. Seals show a strong preference to haul out at low tide, and a diel rhythm in haul-out behaviour has been described during months with day–night cycling and midnight sun (Niizuma 1986). Previous work has reported a preferential haul-out period of 03:00–08:59, with the night period (21:00–03:00) spent at sea (Frost et al, 2001).

Another explanation for when and how often seals occur around set-nets might ultimately involve prey availability or feeding behaviour, as wind speeds had an effect in the model (Table 1).

A study of human–seal conflict in Scotland has shown that only a small number of individuals appeared to be using

rivers and thus interfering with fisheries, which allows conflict management to focus on the selective removal of so-called 'problem' or 'rogue' animals (Graham et al., 2011). In this study, tracked individuals (captured near the set-net area) frequently and carefully visited the fishery area. The results of the present study provide further insights into individual seal behaviour around economically important fishing gear, and will thus help to inform effective management of human-seal conflict in this area.

Acknowledgements

This study was funded by the Japanese Ministry of the Environment through the Environmental Research and Technology Development Fund, Research No. 4-1301 and JSPS KAKENHI Grant Number JP16K20883 Dr. Y Watanuki are acknowledged for their constructive suggestions on drafts of the manuscript. Furthermore, we are also grateful to Akkeshi city hall, Fisheries Division, Fisheries Cooperative Associations (FCAs) staff, Hokkaido Akkeshi Forestry Center, Akkeshi Memorial Hall for Marine, and Akkeshi Marine Station, Aquatic Research Station Field Science Center for Northern Biosphere of Hokkaido University for their support during data collection. We would like to thank the reviewers for their helpful comments and suggestions. We also extend our gratitude to fisherman in both Akkeshi for their time and patience during surveys. Without their assistance, this work could not have been possible.

Conflict of interests

The authors declare that they have no conflicting interests

REFERENCES

- Akkeshi town (2015). Akkeshi town statistical documentation. In Akkeshi town, (Eds.). Akkeshi, Japan, 177pp (in Japanese)
- Bowen WD, Tully D, Boness DJ, Bulheier BM, Marshall GJ (2002). Prey-dependent foraging tactics and prey profitability in a marine mammal. *Mar. Ecol. Prog. Ser.*, 244: 235-245.
- Frost KJ, Michael AS, Lowry LF (2001). Diving behavior of subadult and adult harbor seals in Prince William Sound, Alaska. *Mar. Mam. Sci.*, 17: 813-834.
- Fujii K, Ishikawa T, Watanabe Y, Saito S, Nakagawa E, Kobayashi Y (2005). Tagging and measuring of harbor seals (*Phoca vitulina stejnegeri*) in Erimo Cape, Japan. *Bull Erimo Town Mus.*, 2: 1-8 (in Japanese with English abstract)
- Graham IM, Harris RN, Matejusková IS, Middlemas J (2011). Do 'rogue' seals exist? Implications for seal conservation in the UK. *Anim. Cons.*, 14: 587-598.
- Hayama S (1988). Kuril seal -Present status in Japan-. *AMBIO*, 17: 75-78.
- Hayes AS, Daniel PC, James TH, Le Boeuf BJ (2004). Aquatic mating strategies of the male Pacific Harbor seal (*Phoca vitulina richardii*): are males defending the hotspot? *Mar. Mam. Sci.*, 20: 639-656.
- Hokkaido University Rules and Regulations (2007) Guidelines for Animal Experiments and Animal welfare. http://www.hokudai.ac.jp/jimuk/reiki/reiki_honbun/u01ORG00000658.html (accessed 10 December 2008)
- Inukai T (1942a). True Seals in northern Ocean (1). *Plants and Animals* 10(10): 927-932 (in Japanese)
- Inukai T (1942b). True Seals in northern Ocean (2). *Plants and Animals* 10(11): 1025-1030 (in Japanese)
- Ito T, Shukunobe T (1986). Number and present of Kuril seal In Wada K, Ito T, Niizuma A, Hayama S, Suzuki M (Eds.). *Ecology and Protection of Kuril seal* (pp. 18-58). Tokyo, Japan: Tokai University Press (in Japanese with English abstract).
- Ito T, Wada K (1982). A Preliminary Reports of the Relation between the Kuril Seal and Fisheries on the Pacific Coastal Waters of Eastern Hokkaido, Japan. *Honyurui Kagaku (Mam. Sci.)*, 22: 39-57 (In Japanese)
- Japan Meteorological Agency (2011). http://www.data.jma.go.jp/obd/stats/etrn/index.php?pre_c_no=19&block_no=47418 &year=&month=&day=&view (accessed 10 December 2011)
- Kobayashi Y, Furotani H, Ishikawa K, Sakurai Y (2013). Fishery damage by seals around Akkeshi Bay, eastern Hokkaido, Japan- A questionnaire survey for the local fishermen during 2004-2007-. *Wild. Cons. Japan*, 14(1/2): 53-60 (in Japanese with English abstract)
- Kobayashi Y, Ishikawa K, Harimura I, Goto M, Sakurai Y (2010). Collection of seal-damaged fishes in Akkeshi bay, eastern Hokkaido, Japan. *Nemuro City Mus. His. & Nat.*, 22:29-36 (in Japanese with English abstract)
- Kobayashi Y, Kariya T, Chishima J, Fujii K, Wada K, Baba S, Ito T, Nakaoka T, Kawashima M, Saito S, Aoki N, Hayama SI, Osa Y, Osada H, Niizuma A, Suzuki M, Uekane Y, Hayashi K, Kobayashi M, Ohtaishi N, Sakurai Y (2014). Population trends of the Kuril harbour seal *Phoca vitulina stejnegeri* from 1974 to 2010 in southeastern Hokkaido, Japan. *End. Spe. Res.*, 24: 61-72.
- Kobayashi Y, Kobayashi M, Takada K, Kuraya S, Ogawa I, Horiuchi S, Baba H, Watanabe Y, Sakurai Y (2011). Box and bag traps for capturing Kuril harbour seals *Phoca vitulina stejnegeri*. *Honyurui Kagaku (Mam. Sci.)*, 51: 47-52 (in Japanese with English abstract).
- Lunneryd S G, Fjälling A, Westerberg H (2003). A large-mesh salmon trap: a way of mitigating seal impact on a coastal fishery. *ICES. J. Mar. Sci.*, 60: 1194-1199.
- Niizuma A (1986). Socio-ecology and reproductive strategy of the Kuril seal In Wada K, Ito T, Niizuma A, Hayama S, Suzuki M (Eds.). *Ecology and protection of Kuril seal* (pp. 59-102). Tokyo, Japan: Tokai University Press (in Japanese with English abstract).
- R Development Core Team (2015). R: a language and

- environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. www.r-project.org (accessed 10 December 2015)
- Sweke EA, Kobayashi Y, M Makino, Y Sakurai (2016). Comparative job satisfaction of fishers in northeast Hokkaido, Japan for coastal fisheries management and aquaculture development. *Oce. & Coas. Manag.*, 120 : 170-179.
- The Mammal Society of Japan (2009). Guidelines for the Procedure of Obtaining Mammal Specimens as Approved by the Mammal Society of Japan, The Committee of Reviewing Taxon Names and Specimen Collections, Mammal Society of Japan. <http://www.mammalogy.jp/en/guideline.pdf> (accessed 10 December 2008)
- Thompson PM, Miller D, Cooper R, Hammond PS (1994). Changes in the distribution and activity of female harbour seals during the breeding season: implications for their lactation strategy and mating patterns. *J. Anim. Ecol.*, 63: 24-30.
- Tollit DJ, Black AD, Thompson PM, Mackay A, Corpe HM, Wilson B, Van Parijs SM, Grellier K, Parlane, S (1998). Variations in harbour seal *Phoca vitulina* diet and dive-depths in relation to foraging habitat. *J. Zool.*, 244: 209-222.
- Voegeli FA, Lacroix GL, Anderson JM (1998). Development of miniature pingers for tracking Atlantic salmon smolts at sea. *Hydrobiologia.*, 371/372: 35-46.
- Wada K, Hayama S, Nakaoka T, Uno H (1991) Interactions between Kuril seals and salmon trap net fishery in the coastal waters of southeastern Hokkaido. *Mar. Mam. Sci.*, 7: 75-84.

Appendix 1. Overview of the sampled Kuril seals

Number	Capture location (lat., long.) / trap type	Capture date	Release date	S e x	Age (yr)	Weight (kg)	Total length (cm)	Body length (cm)	Chest girth (cm)	Note
1	① (N43.00363, E144.44379) /	4/1/2009	4/1/2009	F	>4	60.0	144.0	128.0	97.0	pregnant
2	Box trap	4/4/2009	4/4/2009	M	>7	135.0	186.0	166.0	136.0	
3	② (N43.02.251, E144.46737)/ Bag trap	4/6/2009	4/7/2009	M	>7	105.0	170.0	155.5	118.5	