



HOKKAIDO UNIVERSITY

Title	Phylogeography of <i>Littorina brevicula</i> suggests postglacial colonization from south to north along the Japanese Archipelago
Author(s)	Azuma, Noriko; Chiba, Susumu
Citation	The Journal of molluscan studies, 82, 259-267 https://doi.org/10.1093/mollus/eyv052
Issue Date	2016-05
Doc URL	https://hdl.handle.net/2115/65189
Rights	This is a pre-copyedited, author-produced PDF of an article accepted for publication in Journal of Molluscan Studies following peer review. The version of record J. Mollus. Stud. (2016) 82: 259-267 is available online at: http://mollus.oxfordjournals.org/content/82/2/259
Type	journal article
File Information	Azuma.pdf



1 Phylogeography of *Littorina brevicula* suggests post-glacial colonization from south to north along the
2 Japanese Archipelago

3

4 Noriko Azuma¹, Susumu Chiba²

5

6 ¹Graduate School of Fisheries Sciences, Hokkaido University, 3-1-1, Minato-cho, Hakodate 041-8611, Japan

7 ²Department of Aquatic Bioscience, Tokyo University of Agriculture, 196, Yasaka, ABS, Hokkaido, 099-2493

8

9 Running head: Phylogeography of *Littorina brevicula*

10

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31

ABSTRACT

Littorina brevicula is one of the most common gastropods in the intertidal-supralittoral zone around Japan. The northernmost population of this species is around Hokkaido, and the determinant of this northern limit is likely seawater and air temperature. To reconstruct an evolutionary history of this species, we investigated genetic differentiation among 12 populations (three from Hokkaido, six from Honshu and three from Kyushu) using a mitochondrial DNA marker (partial sequence of the NADH dehydrogenase 6 gene). The haplotype network showed shallow genetic divergence within the species, suggesting a bottleneck followed by population expansion. One major haplotype that occurred in 70.5% of all individuals examined was the most frequent in every population sampled. A second major haplotype was abundant around Kyushu but not found in Hokkaido. This skewed haplotype distribution resulted in significant genetic differentiation along the north-south axis of Japan. The importance of the southern clade, which included the second major haplotype, was supported by population analyses of datasets that excluded either the southern clade or the northern clades. The north-south differentiation remained when datasets were analyzed that excluded the northern clades, but disappeared when datasets were analyzed that excluded the southern clade. The combined evidence of shallow divergence and the north-south population structure suggests that the *L. brevicula* population around Japan once declined and then expanded and colonized northward. Although the time of population reduction and re-colonization could not be precisely estimated, the observation that this species is absent further north in Japan suggests that it would have been unable to survive in northern Japan during the last glacial maximum (LGM) and therefore re-colonization likely occurred after the LGM, probably from south to north.

INTRODUCTION

Many species in the northern hemisphere probably underwent changes to their distribution as a result of strong influence from climate oscillation during glacial periods, particularly in the last glacial maximum (LGM) that lasted until around 20,000 years ago and in the subsequent rapid warming. Such distributional changes can be detected by phylogeographic investigations using suitable molecular markers (Hewitt, 2000). The coastal ecosystem was considerably influenced by the LGM and thus various species exhibit genetic evidence of rapid expansion and/or colonization, probably occurring in the post-glacial period. Examples include the near-shore fish *Syngnathus leptorhynchus* (Wilson, 2006) and *Xiphister mucosus* (Marko *et al.*, 2010), seaweed *Pelvetia canaliculata* (Neiva *et al.*, 2014), gastropods *Nassarius nitidus* (Albaina *et al.*, 2012), *Littorina sitkana* and *L. scutulata* (Marko *et al.*, 2010), and starfish *Pisaster ochraceus* and *Evasterias troschelii* (Marko *et al.*, 2010). Around Japan, glacial effects on intertidal molluscs have been suggested in *Batillaria cumingi* (Kojima *et al.*, 2004) and *Cellana nigrolineata* (Nakano, Sasaki & Kase, 2010). However, the more stable temperature of the ocean compared to that of the land allowed some coastal species to maintain their genetic diversity during glacial maxima. Furthermore, the glacial effects sometimes varied within a single species, resulting in a complex population genetic structure, as in *Littorina saxatilis*, in which some populations seemed to have colonized following climate change, while others maintained diversity during a long continuous history (Panova *et al.*, 2011). Such a spatially variable effect within a single species has also been suggested for *Nucella ostrina* and *N. lamellosa* (Marko *et al.*, 2010). The variable genetic consequences of the LGM between or within species are probably due to specific ecological characteristics of each species, i.e. distribution range, habitat, life cycle and abundance, and also to area-specific physical effects of climatic oscillation such as temperature and sea-level change.

In addition to the effects of climate oscillations over paleontological time scales, genetic population structure within a species is affected by contemporary gene flow. Populations of marine species are often weakly structured (i.e. show a low level of differentiation among local populations and/or weak correlation between geography and genetic differentiation) over the species' range (Palumbi, 1994), due to a high level of gene flow caused by active and passive dispersal. In invertebrates that generally showed poor active mobility,

59 gene flow has been thought to be largely affected by the developmental mode and the length of the planktonic
60 larval period, and colonization and contemporary gene flow in marine species are generally mediated by larval
61 dispersal along ocean currents (Palumbi, 1994; Tsang *et al.*, 2008). Recently, larval dispersal has been
62 revealed to be less crucial for determining the genetic structure of marine species than previously supposed
63 (Marko, 2004; Weersing & Toonen, 2009), although it remains an important factor. A comparative study of
64 two planktonic- and two direct-developing *Littorina* species suggested that planktonic larval dispersal reduced
65 genetic differentiation between local populations (Kyle & Boulding, 2000). In their analyses,
66 planktonic-developing species (*Littorina scutulata* and *L. plena*) showed very low levels of genetic
67 differentiation among local populations, while the direct-developing *L. subrotundata* showed higher
68 structuring with larger differentiation between localities. Subsequent research revealed that direct-developing
69 *Littorina* species showed spatial rather than temporal differentiation, and that planktonic-developing species
70 showed the reverse, due to sweepstakes-like reproductive success (Lee & Boulding, 2009). These studies
71 show that *Littorina* is a good model in which to investigate the relationships between genetic population
72 structure, larval dispersal, and historical factors in marine coastal species.

73 *Littorina brevicula* occurs in the littoral fringe of the temperate coast of the northwestern Pacific. Its
74 range extends from Hong Kong to Peter the Great Bay along the continental coast of Asia and from Okinawa
75 to Hokkaido along the Japanese Archipelago (Reid, 1996; Okutani, 2000). On the central to northern Japanese
76 coast, *L. brevicula* is one of the most common snails, occurring in dense aggregations on rocky and boulder
77 shores and on artificial constructions such as breakwaters, tetrapods and slipways. However, it is scarce along
78 the southeastern coast of Hokkaido, which is under the influence of a cold current (Ohgaki, 1983). The
79 spawning season is January to April around Japan and Korea (Kojima, 1957; Son & Hong, 1998). The egg
80 capsule is pelagic and the veligers hatch about 7 days after spawning (Son & Hong, 1998). The total length of
81 the planktonic egg and larval period is several weeks (Golikov, 1976). Such a long planktotrophic period is
82 expected to lower the genetic differentiation between local populations of *L. brevicula* (Reid, 1996).
83 Nevertheless, a spatial genetic structure could potentially be found if samples from a wide geographic range
84 were investigated using highly variable molecular markers (Palumbi, 1994). The fine details of genetic
85 structure across a wide area may provide knowledge about the effects of environmental change on coastal

species on a paleontological scale.

Several studies of population genetics in *L. brevicula* using allozyme variation appeared in the 1990s (e.g. Zaslavskaya, Sergievsky & Tatarenkov, 1992; Tatarenkov, 1995; Park *et al.*, 1999; Zaslavskaya & Takada, 1998), and some of these studies reported genetic differentiation among samples. This was variously attributed to reduced gene flow between distant populations (the east and west coasts of the Sea of Japan; Zaslavskaya & Takada, 1998), the effect of heavy metal pollution (Park *et al.*, 1999), and possible local selection on specific genes (Tatarenkov, 1995). Since these studies did not aim to clarify the total phylogeography of *L. brevicula* throughout the distribution range, they provided little information about hierarchical genetic structure among populations. Nevertheless, Zaslavskaya & Takada (1998) showed genetic differentiation between distant populations, which suggested the possibility of some structuring throughout the species' range.

More recently, DNA markers were used to investigate population structure in *L. brevicula* around Korea (Kim *et al.*, 2003b; Kim, Rodriguez-Lanetty & Song, 2003a). Kim *et al.* (2003b) failed to detect genetic differences among populations, even for populations 1,000 km distant from each other along the coastline. Kim *et al.* (2003a) found significant population differentiation which they attributed to a loss of genetic diversity caused by heavy metal pollution of some populations.

In this study, we investigated the genetic structure among populations of *L. brevicula* in its the main distribution area around Japan, from Hokkaido to Kyushu (Fig. 1). These populations span approximately 2,500 km, and experience considerably different climatic conditions, therefore we expected to observe genetic structure. We chose the mitochondrial NADH dehydrogenase 6 gene (ND6) as a genetic marker, because this region has previously been shown to be more polymorphic than the commonly-used cytochrome-*b* in *L. brevicula* (Kim *et al.*, 2003a). We found population genetic structure that revealed the evolutionary history of *L. brevicula* around Japan, and suggested patterns of population decline, expansion, and migration during recent glacial and post-glacial periods.

110 MATERIAL AND METHODS

111

112 *Sampling, DNA sequencing, and analysis of haplotype genealogy*

113 We examined 540 individuals collected from 12 populations in three regions: three populations from
114 Hokkaido (SSB, ABS, USJ), six from Honshu (FKU, HCH, OBM, CHO, URG, OSK) and three from Kyushu
115 (SIK, AMK, MKR) (Table 1, Fig. 1). Genomic DNA was extracted from a piece of muscle (approximately 1
116 mm³) of ethanol-fixed specimens using the Pure Gene Kit (Qiagen) according to the manufacturer's protocol
117 and resuspended in 100 µl of Tris-EDTA buffer.

118 A partial sequence of ND6 was amplified by PCR in a 30-µl reaction mixture containing template DNA
119 (approximately 200 pg), dNTPs, a pair of primers [Lbnd-F (5' -AGG TAC ATA TTC CTG CGC TCT GAA
120 A-3') and Lbnd-R (5' -GTG TGC GCA TGA AAT GTA T-3')] (Kim *et al.*, 2003) and ExTaq (Takara Bio
121 Inc.), according to the manufacturer's instructions. The thermal cycling profile included precycling
122 denaturation at 95°C for 1 min, followed by 35 cycles of denaturation at 95°C for 30 s, annealing at 50°C for
123 20 s, and extension at 72°C for 45 s. The PCR products were then examined by electrophoresis on a 2%
124 agarose gel, purified with magnetic beads (AMPure, Agencourt), cycle-sequenced using the above-mentioned
125 forward primer and the BigDye® Terminator v. 3.1 Cycle Sequencing Kit (Applied Biosystems) and loaded
126 onto an automated sequencer, ABI PRISMTM 3130 (Applied Biosystems). The obtained sequences were
127 aligned and edited to 450 bp using DNASIS-Mac v. 3.5 (Hitachi) and ClustalX v. 1.81 software (Thompson *et*
128 *al.*, 1997) for defining haplotypes and deposited in the DDBJ/Genbank database with accession nos.
129 AB968435–AB968512.

130 The haplotype genealogy was resolved with a parsimony network using the TCS Network Program
131 (Clement, Posada & Crandall, 2000) under a 95% connection limit. The age of lineage divergence was
132 estimated using the divergence rate of 2.46% per million years (Myr). The rate was based on a calibration for
133 mtDNA including the ND6 region in *L. saxatilis* (Quesada *et al.*, 2007), which assumed a sequence
134 divergence rate of $1.83 \pm 0.21\%$ per Myr. Since Quesada *et al.* applied this rate for the 1.8-kbp sequence
135 including a not very variable region, the calibration may underestimate the evolutionary rate of our ND6 data.
136 To correct the rate, we examined the GenBank data of *L. saxatilis* (accession nos. AM500948–500955,

137 deposited by Quesada *et al.*) and compared sequence divergence within 1.8 kbp of the full length of these data
138 and 0.45 kbp of the ND6 region, orthologous to our sequence data. In the 1.8-kbp sequence, the number of
139 different nucleotides per site was 0.145, while it was 0.195 in the ND6 region in the *L. saxatilis* data. We
140 therefore employed the corrected divergence rate of 2.46% ($1.83 \times 0.195 / 0.145$) per Myr.

141

142 *Population genetic analyses*

143 Arlequin v. 3.1 (Excoffier, Laval & Schneider, 2005) was used to estimate haplotype (h) and nucleotide
144 diversity (π) in each population and to calculate pairwise F_{ST} (Weir & Cockerham, 1984). Genetic distance
145 between populations based on the pairwise F_{ST} was visualized on a two-dimensional surface by nonmetric
146 multidimensional scaling (nMDS) plotting using statistical software R v. 2.9.0 (R Development Core Team,
147 2005). To assess geographic-genetic correlation, the isolation-by-distance (IBD) model (Wright, 1943) was
148 tested using Arlequin 3.1. For the IBD test, the matrix of geographic distances along the coastline between
149 sampling sites was compared with the F_{ST} matrix, and the significance of the correlations was evaluated using
150 the Mantel test with 10,000 permutations.

151 To test the significance of the hierarchical population structure, analysis of molecular variance (AMOVA;
152 Excoffier, Smouse & Quattro, 1992) was conducted with Arlequin 3.1, comparing the three categories of
153 clustering suggested by geography and/or nMDS plotting based on F_{ST} analysis: (1) three regional group
154 clusterings—Hokkaido, Honshu and Kyushu, i.e. [SSB, ABS, USJ], [FKU, HCH, OBM, CHO, URG, OSK]
155 and [SIK, AMK, MKR], respectively; (2) three groups suggested by nMDS plotting based on pairwise F_{ST}
156 analysis—Hokkaido+HCH, Honshu (except HCH), and Kyushu, i.e. [SSB, ABS, USJ, HCH], [FKU, OBM,
157 CHO, URG, OSK] and [SIK, AMK, MKR], respectively; and (3) two groups of clustering by coast—the Sea
158 of Japan and the Pacific Ocean (including the Yellow Sea), i.e. [SSB, ABS, FKU, OBM] and [USJ, HCH,
159 CHO, URG, OSK, SIK, AMK, MKR], respectively. Since Category (3) took into account ocean currents, ABS
160 on the coast of the Sea of Okhotsk, where a branch of the Tsushima current flows in from the Sea of Japan
161 (Fig. 1), was included in the Sea of Japan cluster.

162 When we detected significant genetic structure, we evaluated the effect of each clade (haplotype group
163 derived from the most abundant haplotype shown in Figure 3) on population structure by preparing virtual

164 datasets for population analyses, performing an IBD test, estimating pairwise F_{ST} , and conducting AMOVA
165 using these datasets. Datasets were individually modified by excluding an area-restricted clade (haplotype
166 groups connected to each other in a haplotype network; Fig. 3) and then used to test whether or not the
167 structure appeared. If we found that the structure was lost when a certain clade was excluded from the
168 modified dataset, we would know that the excluded clade was essential for the structure.

169 To test recent population expansions, neutrality tests and mismatch distribution (MMD) analysis with
170 Arlequin 3.1 were added. When our MMD analysis detected population and range expansion, the expansion
171 age was estimated based on the τ value, assuming a sequence-divergence rate of 2.46% per Myr. Neutrality
172 tests of Fu's F_S and Tajima's D were conducted to detect the possibility of population expansion.

173 Based on the genealogy and distribution of haplotypes, we detected the population structure and inferred
174 the shift of the distribution area following a series of steps in the flow chart in Figure 2. The results of several
175 analyses were employed to test the determinants in Figure 2: pairwise F_{ST} estimation for step 1; test of
176 isolation by distance (IBD), nMDS plotting of populations based on pairwise F_{ST} , and AMOVA, for steps 2
177 and 4; construction of the haplotype network and survey of the genealogy for step 3; and estimation of indices
178 of genetic diversity in each population and each area for step 5. The first step, checking heterogeneity between
179 populations, is also informative to evaluate the level of contemporary gene flow, which causes panmictic
180 populations all through the distribution area in extreme cases. The second step, checking the
181 genetic-geographic association, can reveal genetic drift and is useful for detecting population structures
182 skewed by human impact as in *L. brevicula* (Kim *et al.*, 2003b) and *Neptunea arthritica* around Japan (Azuma
183 *et al.*, 2015).

184

185 RESULTS

186

187 *Genealogy and distribution of haplotypes*

188 A total of 78 haplotypes were detected. *LbN6AB03* was the most abundant haplotype, occurring in 381
189 individuals (70.5% of all individuals examined in the present study). This haplotype was also the most
190 frequent in all populations, and was located at the centre of the haplotype network (Fig. 3). The second most

191 frequent haplotype, *LbND6OT02*, occurred in 44 individuals (8.1% of all individuals examined) from Honshu
192 and Kyushu, and the third, *LbND6AO10*, occurred in 10 individuals from Honshu and Hokkaido. Other
193 haplotypes occurred in less than 9 individuals, and 65 haplotypes were private, i.e. found exclusively in a
194 single population (see Supplementary Data). Given that nucleotide divergence between *LbN6AB03* and
195 *LbN6OT02* was 0.22% per site, the time of divergence was estimated as 89 thousand years ago (kya). The
196 largest number of nucleotide substitutions observed was six, 1.33% per site.

197 Several clades appeared in the haplotype network. Six of the clades were found only in Hokkaido and
198 Honshu (northern clades), and one only in Honshu and Kyushu (a southern clade). The rest of the clades
199 appeared in Hokkaido, Honshu and Kyushu, or only in Honshu (middle clades). Some of the following
200 analyses employed modified datasets that excluded the northern clades or the southern clade, to evaluate the
201 importance of these clades in creating the significant genetic structure that we detected.

202

203 *Genetic population structure*

204 As shown in Table 1, h in each population ranged from 0.199 for USJ to 0.677 for MKR, and π ranged from
205 0.0004 for USJ to 0.0025 for URG. The mean diversities of each population were 0.304 (h) and 0.0009 (π) in
206 Hokkaido, 0.546 (h) and 0.0018 (π) in Honshu, and 0.503 (h) and 0.0013 (π) in Kyushu. Although these
207 diversity indices seemed highest in Honshu followed by Kyushu and Hokkaido, no significant differentiation
208 between these three areas appeared in the Kruskal-Wallis test for both h ($\chi^2 = 4.5256$, $df = 2$, $P = 0.104$) and π
209 ($\chi^2 = 3.2488$, $df = 2$, $P = 0.197$).

210 The pairwise F_{ST} values between Hokkaido and Kyushu populations was always higher than 0.07 and
211 always was significantly different from 0, whereas none of the pairwise values from population comparisons
212 within Hokkaido and Kyushu were significant ($\alpha = 0.01$; Table 2). These results indicated a north-south
213 genetic differentiation in this species. The HCH population in Honshu, which did not include haplotypes from
214 the southern clade, showed a significant difference from populations in Kyushu and the URG population. The
215 nMDS plot based on F_{ST} indicated that genetic distance was correlated with a spatial distribution at the region
216 (island) level (Fig. 4 (A)). On the two-dimensional plot, Hokkaido populations were on the left, Kyushu on the
217 right, and Honshu in the middle. However, the relationship among populations within each region did not

218 always follow this genetic-geographic trend.

219 The DNA sequences of Honshu populations did not follow the spatial pattern. FKU, the northernmost
220 among Honshu populations, appeared closest to the Kyushu populations, while HCH, the second northernmost
221 population, was genetically closer to the Hokkaido populations than to the other Honshu populations. The
222 OSK population, the southernmost in Honshu, was genetically close to populations in Hokkaido and not to
223 those in Kyushu. The reason for this variance from an IBD model will be discussed later.

224 The Mantel test showed a significant correlation between genetic (F_{ST}) and geographic distance ($P < 0.01$),
225 indicating that the set of study populations of *L. brevicula* followed an IBD model.

226 As shown in Table 3, AMOVA supported hierarchical population structure clustering (1) comprising
227 Hokkaido, Honshu and Kyushu, and (2) comprising Hokkaido+HCH, Honshu (except HCH) and Kyushu, (P
228 = 0.001 and 0.000, respectively). However, the AMOVA did not support the structure of clustering (3) that
229 comprised two groups—the Sea of Japan and the Pacific Ocean ($P = 0.762$)—indicating that the west and east
230 sides of the Japanese Archipelago could not be separated by their genetic profiles. In each clustering, the
231 variance component was the highest in the category of “within a population” and much higher than “among
232 groups” or “among populations within a group” because of the relatively high genetic diversity in each
233 population and the genetic similarity between populations due to the high frequency of *LbN6AB03* in every
234 population.

235 When the significance of the northern clades and the southern clade was tested using modified datasets, it
236 appeared that the southern clade was essential for population structure. While the unmodified original dataset,
237 the six modified datasets that each excluded one of the six northern clades, and a dataset that excluded all
238 northern clades all showed significant IBD correlations ($P < 0.01$), the genetic-geographic correlation was not
239 significant ($P = 0.840$) when a modified dataset that excluded the southern clade was used. Further the nMDS
240 plot based on pairwise F_{ST} values showed no genetic structure for the modified dataset that excluded the
241 southern clade (Fig. 4 (B)). The significance of the hierarchical structure of [Hokkaido+HCH], [Honshu
242 except HCH] and [Kyushu] was supported by the lowest P -value (0.000) in AMOVA with the unmodified
243 dataset. Table 3 shows the results of analyses on modified datasets testing this structure where all six northern
244 clades were excluded ((2a), significant with $P = 0.000$) and where only the southern clade was excluded ((2b),

not significant with $P = 0.679$), indicating the importance of the southern clade as the determinant of the north-south structure of *L. brevicula* populations around Japan.

Demographic history

The MMD analysis with pooled populations showed that the observed frequencies of haplotype difference did not significantly deviate from the sudden-expansion model when either the sum of squared deviation (SSD; $P = 0.995$) or Harpending's raggedness index ($P = 0.724$) was used. Additionally, the deviation from spatial expansion assuming a constant deme (local population) size model was not significant using either the SSD ($P = 0.990$) or Harpending's raggedness index ($P = 0.730$). The τ values corresponding to the sudden expansion of a population and spatial expansion were estimated to be 0.72 and 0.70, respectively. Given $\tau = 2ut$ ($2u$ = divergence rate, t = time from expansion) and $2u = 2.46\%$ per Myr, the population and spatial expansion dates were estimated to be 65.0 kya and 63.2 kya, respectively (i.e. Upper Pleistocene).

Both Fu's F_S (-29.709, $P=0.0000$) and Tajima's D (-2.653, $P = 0.0000$) indicated that the observed haplotype frequency deviated from the one expected in neutrality, suggesting that the population expanded.

DISCUSSION

Characterization of population genetic structure of *Littorina brevicula* around Japan

The haplotype distribution of *Littorina brevicula* showed a population genetic structure with north-south differentiation. As expected from the planktotrophic developmental mode of *L. brevicula* (Reid, 1996), genetic differentiation between geographically close populations is not significant, and significant genetic differentiation observed between distant populations agreed with the results of Zaslavskaya & Takada (1998). The correlation of geographic and genetic distance was statistically significant in the isolation-by-distance (IBD) test, and the north-south structure was supported by both AMOVA and F_{ST} analyses. The deviation from IBD within the Honshu group (Figure 4A), was probably caused by a stochastic balance of private haplotypes resulting from genetic drift and contemporary gene flow, similar to the lack of genetic structure in the Korean populations (Kim *et al.*, 2003a). In the case of OSK, however, it was possible to attribute the

unreasonable situation to recent human disturbance. OSK showed an unexpected deviation from IBD (Fig. 4 A) and the lowest observed values of both haplotype and nucleotide diversity among all the Honshu populations that could be the result of high genetic drift. Its small effective population size could be attributed to the effects of human impacts such as pollution, as has been argued for the Korean populations (Kim *et al.*, 2003b). OSK is in the Seto Inland Sea, where pollution likely accumulates more than in the open sea. Overall, genetic-geographic correlation and a fair level of genetic diversity within each population indicated that examined populations had not been as affected by recent human activity as it was in the case of *Neptunea arthritica* (Azuma *et al.*, 2015). The higher resistance of *L. brevicula* to anthropogenic effects compared to that of *N. arthritica* is attributable to much larger population sizes and higher gene flow in *L. brevicula* than in *N. arthritica*.

Population genetic structure can be greatly affected by ocean currents. *Littorina brevicula* distributed around Japan currently experiences warm currents (the Kuroshio and Tsushima currents). Comparison of Figures 1 and 5 show that the proportion of the southern clade in each population declines as one travels along the direction of both these currents, until no haplotypes of this clade occur in Hokkaido and HCH. This correlation implies corresponding dispersal pathways for the *L. brevicula* veligers. The HCH population, which is on Honshu where it is less affected by these warm currents, did not include the haplotypes characteristic of the southern clade. We argue below that gene flow by passive dispersal along ocean currents was an important factor for the colonization of *L. brevicula* on a paleontological time scale. Surprisingly, contemporary gene flow mediated by currents has not been strong enough to establish uniform or panmictic populations throughout the distribution of *L. brevicula*; the southern clade has not reached Hokkaido from Kyushu even though it has been 20 ky since the LGM.

AMOVA failed to find east-west genetic differentiation, a result that may be attributable to the remarkable north-south differentiation; the high variation within the east and west groups, which corresponds to a north-south difference, hindered any statistical significance for east-west differentiation in AMOVA, based on the analysis of variance. Additionally, the frequency of the southern clade within each population was similar across Honshu Island, as shown in Figure 5, indicating that east-west differentiation was fundamentally small. In fact, pairwise F_{ST} was 0.000 between CHO and OBM, which are at a similar latitude

on the east and west coasts of Honshu Island respectively.

A glacial effect on Littorina brevicula populations inferred from population genetics

We inferred the glacial effect following the flow chart in Figure 2. First, the presence of a possible glacial effect was evaluated in steps 1 to 3, and then, if a glacial effect was suggested, the position of glacial refugia and post-glacial colonization was identified in steps 4 and 5. The determinants and resolutions in this chart are based on empirical and theoretical knowledge accumulated in phylogeography as reviewed in Avise (2000), Hewitt (2000) and Freeland (2005). The determinant keys of steps 1 and 2 correspond to gene flow and genetic drift. The IBD model is a powerful tool for detecting continuous genetic-geographic structure based on the balance of gene flow and genetic drift. The key of step 3 corresponds to coalescent theory and sequence mismatch distribution (MMD) analysis detailed in Avise (2000). A comparison of genetic diversity in each population can be a good way to detect the direction of colonization, as the sequential founder effect likely caused lower diversity in newly established populations. The southern refugia is plausible for species in the temperate zone of the northern hemisphere (Hewitt, 2000), and the southern refugia often caused post-glacial colonization to trend northward and likely resulted in higher diversity in southern than in northern populations.

Testing the determinants in the flow chart in Figure 2, the presence of step 1 (Genetic difference between local populations) and step 2 (Populations with genetic-geographic correlation) were supported by pairwise F_{ST} , IBD and AMOVA, and we assumed that the genetic-geographic pattern of *L. brevicula* was basically shaped by natural evolution and distribution during the paleontological time scale. Since step 3 (Deep genealogy of haplotypes) was absent, we assumed that recent population expansion probably occurred after the LGM. Significant deviation from neutral evolution indicated by Fu's F and Tajima's D as well as MMD analysis confirmed recent population expansion. Step 4 (North-south differentiation) was present, and step 5 (Higher diversity in the south) could not be proved by statistical analyses, although the mean of diversity indices are lowest in Hokkaido compared to Honshu and Kyushu.

Though the glacial effect was highly possible, northward post-glacial colonization could not be directly supported by genetic evidence. The skewed frequency of the southern clade, however, which appeared as a

key factor in the north-south structure, suggested northward colonization. The fact that the highest proportion of the southern clade was found at MKR (which faces the point of east-west separation) and that the proportion of the southern clade decreased in a similar pattern along both the east and west sides of the Japanese Archipelago (Fig. 5) indicated a gradual colonization of the southern clade to the north in a stepwise manner. Since the northern population of *L. brevicula* likely declined with a decline in air and seawater temperatures, it is possible to assume that a shift in the distribution of *L. brevicula* occurred during glacial and post-glacial periods. Following the decline in temperatures, the northern range of this species trended south, and it seems that a bottleneck effect left only a single haplotype, *LbN6AB03*, around Honshu. An additional haplotype, *LbN6OT02*, survived in the southern area of the Yellow Sea, where temperatures were higher than around Hokkaido or Honshu. In the present study, genetic diversities in Hokkaido populations tend to be lower than those in Honshu populations, though the differences cannot be statistically significant due to the low number of examined Hokkaido populations. This tendency, plus the direction of the ocean currents and the ecological character of *L. brevicula* as described below, implies that this species was absent around Hokkaido during the glacial period and recovered during the post-glacial colonization.

Time estimation corresponding to population expansion

Population and range expansion was estimated to have occurred at 65.0 kya and 63.2 kya, long before the end of the LGM at 20 kya. The estimated date might be earlier than the actual expansion time, however, as the divergence rate of 2.46% per Myr, a rate modified in this study from the molecular clock calibration by Quesada *et al* (2007), might overestimate the divergence time within *L. brevicula*. The calibration by Quesada *et al* (2007) may not be adequate for the time scale of the present study, as the evolutionary rate of $1.83 \pm 0.21\%$ per Myr was based on the split of the *Hydrobia/Peringia* 5.33 million years ago, far earlier than the age in the scope of our study, and such calibration based on speciation is known to overestimate divergence time at the population level within a species (Ho *et al.*, 2005). Thus, for precise dating, molecular clock calibration based on a more recent event should be established for *L. brevicula* or closely-related *Littorina* species.

Ecological and paleontological perspectives supporting the hypothesis of post-glacial colonization

353 The hypothesis of glacial effects and post-glacial colonization inferred from phylogeography is also supported
354 by ecological and paleontological perspectives of *L. brevicula* around Japan. Fossil records suggest that the
355 speciation event of *L. brevicula* occurred no later than the upper Pliocene, and molecular phylogeny suggests
356 that the divergence between *L. brevicula* and *L. mandshurica* occurred by the lower Pleistocene at the latest
357 (Reid, Dyal & Williams, 2012). The species should therefore have a long history, but the genetic genealogy of
358 the extant population at present was shallow, indicating that the population declined in the upper Pleistocene.
359 Probably only two mitochondrial lineages (*LbN6AB03* and *LbN6OT02*) remained after the latest bottleneck
360 event, which was likely associated with glacial cooling. Generally, temperature is arguably the most critical
361 factor determining the geographical distribution of species (e.g., Raffaelli & Hawkins, 1996). Considering the
362 temperature range in which extant populations are currently found, the low temperatures during glacial
363 periods can be expected to have been severe for the survival and reproduction of *L. brevicula* around northern
364 Japan. Though the lower temperature limit for the reproduction of *L. brevicula* has not been reported, it is
365 probably 3 - 5 °C given that this species is rare along areas of the Pacific coast under the influence of the cold
366 Kuril Current but abundant on the Sea of Okhotsk coast of Hokkaido (Ohgaki, 1983; Azuma & Chiba personal
367 observation). Since low seawater temperature likely determines the present northern limit of *L. brevicula* in
368 Hokkaido, it is safe to assume that there were no *L. brevicula* around Hokkaido during glacial periods and that
369 post-glacial colonization formed the extant Hokkaido populations.

370 We could not find higher genetic diversity in Kyushu, the southern area, as mean diversity indices in each
371 population were estimated to be higher in Honshu than Kyushu (Table 1). This might be attributable to
372 population decline around Kyushu in recent years. Although both haplotype and nucleotide diversity at the
373 southernmost sampling point (MKR) was the highest among all examined populations, *L. brevicula* is not
374 common there at present, suggesting that conditions are no longer optimal and might possibly be too warm for
375 this species. The upper limit for spawning was reported to be 13 °C (Son & Hong, 1998), and high seawater
376 temperatures around Kyushu, ordinarily greater than 15 °C even in January, may also limit the reproduction of
377 *L. brevicula* and may have caused population decline and a reduction in genetic diversity in this area after
378 post-glacial colonization. Since genetic diversity in Kyushu was probably higher in the past (long after the
379 LGM) compared to the present, we assume a southern refugia existed around Kyushu or further south during

380 LGM and that subsequent northward post-glacial colonization resulted in higher genetic diversity in the south
381 than in the north.

382 A glacial effect was also suggested in *Batillaria cumingi* (Kojima *et al.*, 2004) and *Cellana nigrolineata*
383 (Nakano *et al.*, 2010) around Japan, although no evidence of post-glacial colonization was provided in those
384 studies. The present study represents the first report of genetic evidence for such colonization in an intertidal
385 mollusc species along the Japanese Archipelago.

386

387

388 ACKNOWLEDGEMENTS

389

390 The authors are grateful to editor David Reid for the English language review and to K. Imai, Z. Ishizuka, T.
391 Kokita, J. Shoji, H. Takahara and T. Sonoda for their kind assistance in collecting samples. The present study
392 was partly supported by Grant-in-Aid for Advanced Study Projects from the Tokyo University of Agriculture.

393

394

395 REFERENCES

396 ALBAINA, N., OLSEN, J.L., COUCEIRO, L., RUIZ, J.M. & BARREIRO, R. 2012. Recent history of the
397 European *Nassarius nitidus* (Gastropoda): phylogeographic evidence of glacial refugia and colonization
398 pathways. *Marine Biology*, **159**: 1871–1884.

399 AVISE, J.C. 2000. *Phylogeography: the history and formation of species*. Harvard University Press,
400 Cambridge.

401 AZUMA, N., MIRANDA R.M., GOSHIMA, S. & ABE, S. 2015. Phylogeography of Neptune whelk
402 (*Neptunea arthritica*) suggests sex-biased impact of tributyltin pollution and overfishing around northern
403 Japan. *Journal of Molluscan Studies*, **81**: 131–138.

404 CLEMENT, M., POSADA, D. & CRANDALL, K.A. 2000. TCS: a computer program to estimate gene
405 genealogies. *Molecular Ecology*, **9**: 1657–1659.

406 EXCOFFIER, L., LAVAL, G. & SCHNEIDER, S. 2005. Arlequin v. 3.0: an integrated software package for

407 population genetics data analysis. *Evolutionary Bioinformatics Online*, **1**: 47–50.

408 EXCOFFIER, L., SMOUSE, P.E. & QUATTRO, J.M. 1992. Analysis of molecular variance inferred from
 409 metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data.
 410 *Genetics*, **131**: 479–491.

411 FREELAND, J.R. 2005. *Molecular ecology*. JohnWiley, Chichester.

412 GOLIKOV, A. N. 1976. Gastropoda. *The animals and plants of Peter the Great Bay*. Nauka, Leningrad: 79–92
 413 (in Russian).

414 HEWITT, G.M. 2000. The genetic legacy of the Quaternary ice ages. *Nature*, **405**: 907–913.

415 HO, S. Y. W., PHILLIPS, M. J., COOPER, A. & DRUMMOND, A. J. Time Dependency of Molecular Rate
 416 Estimates and Systematic Overestimation of Recent Divergence Times. *Molecular Biology and Evolution*,
 417 **22**: 1561–1568.

418 KIM, S.J., RODRIGUEZ-LANETTY, M. & SONG, J.I. 2003a. Genetic population structure of *Littorina*
 419 *brevicula* around Korean waters. *Hydrobiologia*, **505**: 41–48.

420 KIM, S.J., RODRIGUEZ-LANETTY, M., SUH, J.H. & SONG, J.I. 2003b. Emergent effects of heavy metal
 421 pollution at a population level: *Littorina brevicula* a study case. *Marine Pollution Bulletin*, **46**: 74–80.

422 KOJIMA, S., HAYASHI, I, KIM, D., IIJIMA, A. & FUROTA, T. 2004. Phylogeography of an intertidal
 423 direct-developing gastropod *Batillaria cumingi* around the Japanese Islands. *Marine Ecology Progress*
 424 *Series*, **276**: 161–172.

425 KOJIMA, Y. 1957. On the breeding of periwinkle, *Littorina brevicula* (Philippi). *Bulletin of the Biological*
 426 *Station of Asamushi, Tohoku University*, **8**: 59 – 62.

427 KYLE, C.J. & BOULDING, E.G. 2000. Comparative population genetic structure of marine gastropods
 428 (*Littorina* spp.) with and without pelagic larval dispersal. *Marine Biology*, **137**: 835–845.

429 LEE, H.J. & BOULDING, E.G. 2009. Spatial and temporal population genetic structure of four northeastern
 430 Pacific littorinid gastropods: the effect of mode of larval development on variation at one mitochondrial
 431 and two nuclear DNA markers. *Molecular Ecology*, **18**: 2165–2184.

432 MARKO, P.B. 2004. ‘What’s larvae got to do with it?’ Disparate patterns of post-glacial population structure
 433 in two benthic marine gastropods with identical dispersal potential. *Molecular Ecology*, **13**: 597–611.

MARKO, P.B., HOFFMAN, J.M., EMME, S.A., MCGOVERN, T.M., KEEVER, C. & COX, L.M. 2010. The expansion-contraction model of Pleistocene demography: rocky shores suffer a sea change? *Molecular Ecology*, **19**:146–169.

NAKANO, T., SASAKI, T. & KASE, T. 2010. Color polymorphism and historical biogeography in the Japanese patellogastropod limpet *Cellana nigrolineata* (Reeve) (Patellogastropoda: Nacellidae). *Zoological Science*, **27**: 811–820.

NEIVA, J., ASSIS, J., FERNANDES, F., PEARSON, G.A. & SERRÃO, E.A. 2014. Species models and mitochondrial DNA phylogeography suggest an extensive biogeographical shift in the high-intertidal seaweed *Pelvetia canaliculata*. *Journal of Biogeography*, **41**: 1137–1148.

OHGAKI, S. 1983. Distribution of the family Littorinidae (Gastropoda) in Hokkaido, with special emphasis on the distribution in Akkeshi Bay. *Nankiseibutu*, **25**: 173–180. (in Japanese)

OKUTANI, T. 2000. *Marine Mollusks in Japan*. Tokai University Press, Kanagawa.

PALUMBI, S.R. 1994. Genetic divergence, reproductive isolation, and marine speciation. *Annual Review of Ecology and Systematics*, **25**: 547–572.

PANOVA, M., BLAKESLEE, A.M.H., MILLER, A.W., MÄKINEN, T., RUIZ, G.M., JOHANNESSEN, K. & ANDRE, C. 2011. Glacial history of the North Atlantic Marine snail, *Littorina saxatilis*, inferred from distribution of mitochondrial DNA lineages. *PLoS ONE*, **6**: e17511.

PARK, K.S., SONG, J.I., CHOE, B.L. & KIM, S.J. 1999. Amylase polymorphism of *Littorina brevicula* from polluted and unpolluted sites, Korea. *Bulletin of Environmental Contamination and Toxicology*, **63**: 633–638.

QUESADA, H., POSADA, D., CABALLERO, A., MORÁN, P. & ROLÁN-ALVAREZ, E. 2007. Phylogenetic evidence for multiple sympatric ecological diversification in a marine snail. *Evolution*, **61**: 1600–1612.

R DEVELOPMENT CORE TEAM. 2005. *R: a language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL <http://www.R-project.org>.

RAFFAELLI, D. & HAWKINS, S. 1996. *Intertidal ecology*. Chapman & Hall, London.

461 REID, D.G. 1996. *Systematics and evolution of Littorina*. Ray Society, London.

462 REID, D.G., DYAL, P. & WILLIAMS, S.T. 2012. A global molecular phylogeny of 147 periwinkle species
 463 (Gastropoda, Littorininae). *Zoologica Scripta*, **41**: 125–136.

464 SON, M.H. & HONG, S.U. 1998. Reproduction of *Littorina brevicula* in Korean waters. *Marine Ecology*
 465 *Progress Series*, **172**: 215–223.

466 TATARENKOV, A.N. 1995. Genetic heterogeneity in populations of *Littorina brevicula* (Philippi) (Mollusca:
 467 Gastropoda) in the northern part of Peter the Great Bay (Sea of Japan). *Veliger*, **38**: 85–91.

468 THOMPSON, J.D., GIBSON, T.J., PLEWNIK, F., JEANMOUGIN, F. & HIGGINS, D.G. 1997. The
 469 ClustalX windows interface: flexible strategies for multiple sequence alignment aided by quality analysis
 470 tools. *Nucleic Acids Research*, **24**: 4876–4882.

471 WEERSING, K. & TOONEN, R. J. 2009. Population genetics, larval dispersal, and connectivity in marine
 472 systems. *Marine Ecology Progress Series*, **393**: 1–12.

473 WEIR, B.S. & COCKERHAM, C.C. 1984. Estimating *F*-statistics for the analysis of population structure.
 474 *Evolution*, **38**: 1358–1370.

475 WILSON, A.B. 2006. Genetic signature of recent glaciation on populations of a near-shore marine fish species
 476 (*Syngnathus leptorhynchus*). *Molecular Ecology*, **15**: 1857–1871.

477 WRIGHT, S. 1943. Isolation by distance. *Genetics*, **28**: 114–138.

478 YOKOYAMA, M. 1960. Tertiary fossils from various localities in Japan. Part IV. *Palaeontological Society of*
 479 *Japan Special Papers*, **6**: 1–89.

480 YOKOYAMA, M. 1925. Tertiary Mollusca from Shinano and Echigo. *Journal of Faculty of Science. Imperial*
 481 *University of Tokyo. Section 2*, **1**: 1–23.

482 ZASLAVSKAYA, N.I. & TAKADA, Y. 1998. Allozyme variation and behavioural dimorphism among
 483 populations of *Littorina brevicula* (Philippi) from Japan. *Hydrobiologia*, **378**: 53–57.

484 ZASLAVSKAYA N.I., SERGIEVSKY S.O. & TATARENKOV A.N. 1992. Genetic biochemical comparison
 485 of Atlantic and Pacific species of *Littorina* (Gastropoda: Littorinidae). *Genetika*, **28**: 89–98. (Russian)
 486

487 **FIGURE CAPTIONS**

488 **Figure 1.** Ocean currents around Japan, and sampling locations of *Littorina brevicula* in the present study:
489 three in Hokkaido, six in Honshu, and three in Kyushu.

490 **Figure 2.** Flow chart to evaluate glacial effects on population structure and post-glacial colonization inferred
491 from genetic data. The path of inference in the present study follows the circled determinations (present or
492 absent).

493 **Figure 3.** Parsimony network of the mtDNA ND6 haplotypes of *Littorina brevicula*. A solid line between
494 circles indicates a single nucleotide substitution. Circle size reflects haplotype abundance (number of
495 individuals that had the haplotype), and the colors in the circles indicate the proportions of Hokkaido, Honshu,
496 and Kyushu individuals. Dotted lines indicate the criteria of each clade (a monophyletic group of multiple
497 haplotypes). Colors of dotted lines indicate clade categories (northern, southern, middle).

498 **Figure 4.** nMDS plot of *Littorina brevicula* with pairwise F_{ST} values based on a 450-bp sequence of partial
499 mtDNA ND6 for unmodified dataset (A) and a dataset excluding the southern clade (B). While population
500 structure along the Japanese Archipelago from north to south (from left to right in this plot) appeared in (A),
501 no such structure was observed in (B), indicating the north-south structure was attributable to the existence of
502 the southern clade.

503 **Figure 5.** Distribution of mtDNA ND6 haplotype groups in each sampling locality of *Littorina brevicula*.
504 *LbND6AB03* was the most abundant in all populations, while the southern clade was abundant in southern but
505 not in northern populations.

Table 1. Geographical and genetic information for *Littorina brevicula* populations analyzed in the present study. Sampling locality, Collection date, Sample size (number of individuals), number of haplotype and diversity indices, haplotype richness (*HR*, number of haplotypes standardized by the smallest sample size), haplotype diversity (*h*) and nucleotide diversity (π) estimated in partial ND6 sequence. *Average for all samples. **Calculated pooling all samples as a single population.

Region	code. Sample name	latitude; longitude	year / month	Sample size	No. of haplotype	<i>HR</i>	<i>h</i>	π
Hokkaido	1. SSB	44°32'N; 141°45'E	2010/July	48	10	5.5	0.343	0.0011
	2. ABS	44°03'N; 144°15'E	2010/June	39	8	5.6	0.371	0.0012
	3. USJ	41°56'N; 140°56'E	2013/June	48	6	3.5	0.199	0.0004
	mean of Hokkaido					4.8	0.304	0.0009
Honshu	4. FKU	40°46'N; 140°03'E	2011/October	48	11	7.6	0.647	0.0018
	5. HCH	40°32'N; 141°33'E	2011/November	48	11	6.8	0.469	0.0015
	6. OBM	35°32'N; 135°42'E	2011/October	48	12	7.6	0.578	0.0018
	7. CHO	35°42'N; 140°52'E	2012/January	47	17	9.8	0.646	0.0021
	8. URG	35°15'N; 139°44'E	2013/August	48	14	8.4	0.629	0.0025
	9. OSK	34°17'N; 132°54'E	2011/November	47	7	4.4	0.311	0.0009
	mean of Honshu					7.4	0.546	0.0018
Kyushu	10. SIK	32°48'N; 131°53'E	2011/September	46	5	3.6	0.423	0.0011
	11. AMK	32°36'N; 130°29'E	2011/September	24	3	3	0.409	0.0009
	12. MKR	31°15'N; 130°18'E	2013/May	49	14	7.9	0.677	0.0020
	mean of Kyushu					4.8	0.503	0.0013
Total				540	78	6.0*	0.497**	0.0015**

Table 2. Left lower: Pairwise F_{ST} between *Littorina brevicula* populations based on partial sequence of ND6. Bold font indicates the P value < 0.05. Right upper: Significance level of F_{ST} . + indicates a P value < 0.01 after sequential Bonferroni correction.

		1.	2.	3.	4.	5.	6.	7.	8.	9.	10.	11.	12.
Hokkaido	1. SSB		-	-	+	-	-	-	-	-	+	+	+
	2. ABS	0.010		-	-	-	-	-	+	-	+	+	+
	3. USJ	0.000	0.010		-	-	+	-	-	+	+	+	+
Honshu	4.FKU	0.033	0.016	0.034		-	-	-	-	-	-	-	-
	5. HCH	0.009	0.000	0.010	0.009		-	-	+	-	+	-	+
	6. OBM	0.014	0.017	0.021	0.005	0.014		-	-	-	-	-	-
	7. CHO	0.007	0.002	0.005	0.000	0.001	0.000		-	-	+	-	-
	8. URG	0.010	0.021	0.017	0.016	0.020	0.011	0.002		-	-	-	-
	9. OSK	0.006	0.008	0.009	0.011	0.006	0.006	0.000	0.010		-	-	-
Kyushu	10. SEK	0.089	0.093	0.128	0.017	0.078	0.027	0.030	0.027	0.041		-	-
	11. AMK	0.073	0.077	0.138	0.003	0.060	0.013	0.013	0.011	0.030	0.000		-
	12. MKR	0.080	0.079	0.098	0.019	0.071	0.032	0.034	0.030	0.042	0.000	0.000	

509

510

511

512

Table 3. Results of five separate AMOVA using K2P distance. (1): 3 groups = [Hokkaido] [Honshu] [Kyushu]. (2): 3 groups = [Hokkaido+HCH] [Honshu (except HCH)] [Kyushu]. (3): 2 groups = [Pacific Ocean + Yellow Sea] [Sea of Japan + Sea of Okhotsk]. (2a): Population groups were similar to (2), and all northern clades were excluded before analysis. (2b): Population groups were similar to (2), and the southern clade was excluded before analysis.

	Source of variation	Variance component	% of variation	<i>P</i> value	Fixation indices
(1)	Among groups	0.00994	2.86	0.001	$F_{CT}=0.028$
	Among population within groups	0.00124	0.36	0.154	$F_{SC}=0.003$
	Within populations	0.33538	96.78	0.000	$F_{ST}=0.032$
(2)	Among groups	0.01035	2.99	0.000	$F_{CT}=0.029$
	Among population within groups	0.00059	0.17	0.260	$F_{SC}=0.001$
	Within populations	0.33538	96.84	0.000	$F_{ST}=0.031$
(3)	Among groups	-0.00128	-0.37	0.762	$F_{CT}=-0.003$
	Among population within groups	0.00852	2.48	0.000	$F_{SC}=0.021$
	Within populations	0.33538	97.89	0.000	$F_{ST}=0.025$
(2a)	Among groups	0.01111	5.02	0.000	$F_{CT}=0.050$
	Among population within groups	-0.00048	-0.21	0.570	$F_{SC}=-0.002$
	Within populations	0.21044	95.1	0.000	$F_{ST}=0.048$
(2b)	Among groups	-0.00015	-0.12	0.679	$F_{CT}=-0.000$
	Among population within groups	0.00219	0.81	0.000	$F_{SC}=0.007$
	Within populations	0.28352	99.31	0.000	$F_{ST}=0.007$

513

Fig 1




Fig 2





Fig 3








Fig 5

1



Supplementary Data

Haplotype frequency (number of individuals which showed the haplotype) of partial ND6 in each populations. Haplotype name is abbreviated as *LbN6SS18* into *SS18*.

[illegible]

3. USJ	1	1	1	-	1	-	-	-	1	-	-	-	-	-	-	-
4. FKU	-	-	-	1	4	1	1	1	1	1	-	-	-	-	-	-
5. HCH	-	-	-	-	2	-	-	-	1	-	1	1	1	1	1	1
6. OBM	-	-	-	-	1	-	-	-	-	-	1	-	-	-	-	-
7. CHO	-	-	-	-	2	-	-	-	-	-	-	-	-	-	-	-
8. URG	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
9. OSK	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-
10. SIK	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
11. AMK	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
12. MKR	-	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-
Total	1	1	1	1	10	1	1	1	5	1	2	1	1	1	2	1

[illegible]

