



HOKKAIDO UNIVERSITY

Title	VIP1 is very important/interesting protein 1 regulating touch responses of Arabidopsis
Author(s)	Tsugama, Daisuke; Liu, Shenkui; Takano, Tetsuo
Citation	Plant signaling & behavior, 11(6), e1187358 https://doi.org/10.1080/15592324.2016.1187358
Issue Date	2016-06-02
Doc URL	https://hdl.handle.net/2115/65244
Rights	This is an Accepted Manuscript of an article published by Taylor & Francis Group in Plant Signaling & Behavior on 12/05/2016, available online: http://www.tandfonline.com/10.1080/15592324.2016.1187358 .
Type	journal article
File Information	Plant Signal Behav11-6_e1187358.pdf



VIP1 is Very Important/Interesting Protein 1 regulating touch responses of Arabidopsis

Daisuke Tsugama^{1*}, Shenkui Liu², and Tetsuo Takano³

¹Laboratory of Crop Physiology, Research Faculty of Agriculture, Hokkaido University

Kita 9 Nishi 9 Kita-ku, Sapporo-shi, Hokkaido 060-8589, Japan

²Alkali Soil Natural Environmental Science Center, Northeast Forestry University

No. 26 Hexing Road, Xiangfang District, Harbin 150040, P.R. China

³Asian Natural Environmental Science Center, The University of Tokyo

1-1-1 Midori-cho, Nishitokyo-shi, Tokyo 188-0002, Japan

Addendum to:

Tsugama D, Liu S, Takano T. The bZIP protein VIP1 is involved in touch responses in

Arabidopsis roots. Plant Physiol 2016; in Press.

Submitted: 14 April 2016

Accepted:

Key words: *Arabidopsis thaliana*; bZIP transcription factor; calcium signaling; cell death;

cell wall; hypo-osmotic stress; mechanical stimuli; protein phosphorylation; root cap; root

waving

***Correspondence to:**

Daisuke Tsugama, Email: tsugama@res.agr.hokudai.ac.jp

22 **Abstract**

23 VIP1 (VIRE2-INTERACTING PROTEIN 1) is a bZIP transcription factor in *Arabidopsis*
24 *thaliana*. VIP1 and its close homologs (i.e., Arabidopsis group I bZIP proteins) are present in
25 the cytoplasm under steady conditions, but are transiently localized to the nucleus when cells
26 are exposed to hypo-osmotic conditions, which mimic mechanical stimuli such as touch.
27 Recently we have reported that overexpression of a repression domain-fused form of VIP1
28 represses the expression of some touch-responsive genes, changes structures and/or local
29 auxin responses of the root cap cells, and enhances the touch-induced root waving. This
30 raises the possibility that VIP1 suppresses touch-induced responses. VIP1 should be useful to
31 further characterize touch responses of plants. Here we discuss two seemingly interesting
32 perspectives about VIP1: (1) What factors are involved in regulating the nuclear localization
33 of VIP1?; (2) What can be done to further characterize the physiological functions of VIP1
34 and other Arabidopsis group I bZIP proteins?

35

36 **TEXT**

37 VIP1 (VIRE2-INTERACTING PROTEIN 1) was originally identified as an *Arabidopsis*
38 *thaliana* bZIP protein interacting with the *Agrobacterium tumefaciens* protein VirE2,¹ and has
39 been suggested to play pleiotropic roles.¹⁻¹² Previously we reported that VIP1 exists mainly in
40 the cytosol under steady conditions but transiently accumulates in the nucleus when cells are

41 exposed to hypo-osmotic conditions,⁹ that close homologs of VIP1 (i.e., Arabidopsis group I
42 bZIP proteins¹³) also exhibit such a pattern of subcellular localization,¹¹ and that overexpression
43 of a repression domain-fused form of VIP1 (VIP1-SRDX) enhances touch-induced root
44 waving.¹⁴ Touch as well as the hypo-osmotic conditions seems to induce the nuclear
45 localization of VIP1 (Fig. 1). VIP1 is a novel regulator of touch responses of Arabidopsis,
46 and should be useful to further characterize them. Here we discuss perspectives for further
47 studies on VIP1.

48

49 **What factors are involved in regulating the nuclear localization of VIP1?**

50 Previously a treatment with a microbe associated molecular pattern, flg22, caused VIP1 to be
51 localized to the nucleus,⁶ and a treatment with a gibberellin biosynthesis inhibitor, uniconazole-P,
52 caused the putative tobacco VIP1 ortholog RSG (REPRESSION OF SHOOT GROWTH) to
53 be localized to the nucleus.¹⁵ However, in our experiments, water itself, which was used to
54 dilute flg22 and uniconazole-P stocks in those experiments, can induce the nuclear localization of
55 VIP1 (ref. 10 and unpublished data), thus the effects of these chemicals may have to be
56 re-evaluated.

57

58 VIP1 is thought to accumulate in the nucleus when its 79th serine is phosphorylated by
59 MPK3 (MITOGEN-ACTIVATED PROTEIN KINASE 3),⁶ although this idea has been

60 questioned.¹¹ On the other hand, RSG is thought to accumulate in the nucleus when its
61 phosphorylated 114th serine is either dephosphorylated or replaced by alanine.¹⁵ Although
62 replacing the VIP1 115th serine, which corresponds to the RSG 114th serine, with alanine
63 does not affect the subcellular localization of VIP1,¹¹ our unpublished data suggest that VIP1
64 has multiple putative phosphorylation sites, and that these sites are dephosphorylated when
65 VIP1 is localized to the nucleus. The RSG 114th serine is phosphorylated by the
66 calcium-dependent protein kinase CDPK1 in tobacco,¹⁶ thus a CDPK1 homolog may
67 phosphorylate VIP1 in Arabidopsis. Our unpublished data also suggest that calcium signaling
68 regulates both the nuclear import and the nuclear export of VIP1. It should be interesting to
69 identify protein kinases and protein phosphatases regulating VIP1 phosphorylation states, and
70 to identify the phosphorylation sites in VIP1.

71

72 The mechanosensitive calcium channels MCA1 and MCA2 (MID-COMPLEMENTING
73 ACTIVITY 1 and 2, respectively), the seven-transmembrane domain proteins MLO4 and
74 MLO11 (MILDEW RESISTANCE LOCUS O 4 and 11, respectively), and the receptor
75 kinase FERONIA have been identified as candidate mechanical stress sensors regulating
76 calcium signaling and root tropisms.¹⁷⁻²⁰ *TOUCH2* and *TOUCH4*, which are up-regulated by
77 the FERONIA-mediated touch-responsive signaling,²⁰ are unlikely to be VIP1 target genes.¹⁴
78 However, it would be worth examining whether VIP1 interacts with these proteins and/or acts

under the control of them. To identify novel regulators for VIP1, it would also be helpful and practical to screen for chemicals and genetic mutations that affect the subcellular localization of VIP1.

What can be done to further characterize the physiological functions of VIP1 and other Arabidopsis group I bZIP proteins?

VIP1 and its close homologs in tomato, tobacco, and rice, have been identified in different studies as a regulator of certain physiological responses.^{15,16,21-26} This would support the idea that such VIP1 homologs have important, pleiotropic roles. Thus far only two Arabidopsis group I bZIP protein genes, *VIP1* and *PosF21*, have been associated with physiological roles.^{1-12,14,27} However, at least five of the other group I bZIP protein genes are expressed as highly as *VIP1* and *PosF21*,¹¹ and they could function redundantly. In our recent study, expressing the VIP1-GFP fusion protein suppressed the VIP1-SRDX-induced enhancement of root waving.¹⁴ It would be interesting to examine whether other group I bZIP proteins can also suppress the VIP1-SRDX-induced enhancement of root waving. In our preliminary experiments, the triple mutant that has T-DNA in *VIP1*, *PosF21*, and *bZIP29* (another group I bZIP protein gene) was similar to the wild type in phenotypes including root waving, but knocking out various combinations of the group I bZIP protein genes should also help to elucidate the physiological roles of them.

98 The local auxin responses in the root tip are different between wild-type plants and
99 VIP1-SRDX-overexpressing (VIP1-SRDXox) plants, and this may be attributed to the
100 difference in adhesion and/or removal of the root cap cells. Expression levels of some
101 mechanical stimulus-induced genes that should regulate cell wall properties are lower in
102 VIP1-SRDXox plants than in the wild-type plants, and this may cause the abnormal root cap
103 cell adhesion/removal in VIP1-SRDXox plants.¹⁴ On the other hand, cell death mediated by
104 the NAC (NO APICAL MERISTEM)-family transcription factor SOMBRERO and the S1-P1
105 nuclease-family protein BFN1 (BIFUNCTIONAL NUCLEASE 1) is necessary for the
106 removal of the lateral root cap cells.^{28,29} It would be interesting to characterize the cell wall
107 properties and cell death in the root cap cells of VIP1-SRDXox plants, and to examine
108 genetic and physical interactions between the group I bZIP proteins and the above regulators
109 of cell wall properties and cell death. It would also be important to further evaluate how
110 mechanical stimuli affect cell wall properties and cell death in root cap cells.

111

112 **Acknowledgments**

113 This work was supported by Grant-in-Aid for JSPS Fellows (grant number 25-7247) to D.T.

References

1. Tzfira T, Vaidya M, Citovsky V. VIP1, an Arabidopsis protein that interacts with Agrobacterium VirE2, is involved in VirE2 nuclear import and Agrobacterium infectivity. EMBO J 2001; 20:3596-607.
2. Tzfira T, Vaidya M, Citovsky V. Increasing plant susceptibility to Agrobacterium infection by overexpression of the Arabidopsis nuclear protein VIP1. Proc Natl Acad Sci USA 2002; 99:10435-40.
3. Tzfira T, Vaidya M, Citovsky V. Involvement of targeted proteolysis in plant genetic transformation by Agrobacterium. Nature 2004; 431:87-92.
4. Lacroix B, Vaidya M, Tzfira T, Citovsky V. The VirE3 protein of Agrobacterium mimics a host cell function required for plant genetic transformation. EMBO J 2005; 24:428-37.
5. Li J, Krichevsky A, Vaidya M, Tzfira T, Citovsky V. Uncoupling of the functions of the Arabidopsis VIP1 protein in transient and stable plant genetic transformation by Agrobacterium. Proc Natl Acad Sci USA 2005; 102:5733-8.
6. Djamei A, Pitzschke A, Nakagami H, Rajh I, Hirt H. Trojan horse strategy in Agrobacterium transformation: abusing MAPK defense signaling. Science 2007; 318:453-6.
7. Pitzschke A, Djamei A, Teige M, Hirt H. VIP1 response elements mediate mitogen-activated protein kinase 3-induced stress gene expression. Proc Natl Acad Sci USA 2009; 106:18414-9.

- 133 8. Wu Y, Zhao Q, Gao L, Yu XM, Fang P, Oliver DJ, Xiang CB. Isolation and
134 characterization of low-sulphur-tolerant mutants of *Arabidopsis*. *J Exp Bot* 2010;
135 61:3407-22.
- 136 9. Tsugama D, Liu S, Takano T. A bZIP protein, VIP1, is a regulator of osmosensory
137 signaling in *Arabidopsis*. *Plant Physiol* 2012; 159:144-55.
- 138 10. Lacroix B, Citovsky V. Characterization of VIP1 activity as a transcriptional regulator in
139 vitro and in planta. *Sci Rep* 2013; 3:2440.
- 140 11. Tsugama D, Liu S, Takano T. Analysis of functions of VIP1 and its close homologs in
141 osmosensory responses of *Arabidopsis thaliana*. *PLoS One* 2014; 9:e103930.
- 142 12. Shi Y, Lee LY, Gelvin SB. Is VIP1 important for *Agrobacterium*-mediated transformation?
143 *Plant J* 2014; 79:848-60.
- 144 13. Jakoby M, Weisshaar B, Dröge-Laser W, Vicente-Carbajosa J, Tiedemann J, Kroj T,
145 Parcy F; bZIP Research Group. bZIP transcription factors in *Arabidopsis*. *Trends Plant*
146 *Sci* 2002; 7:106-11.
- 147 14. Tsugama D, Liu S, Takano T. The bZIP protein VIP1 is involved in touch responses in
148 *Arabidopsis* roots. *Plant Physiol* 2016; in Press.
- 149 15. Ishida S, Fukazawa J, Yuasa T, Takahashi Y. Involvement of 14-3-3 signaling protein
150 binding in the functional regulation of the transcriptional activator REPRESSION OF
151 SHOOT GROWTH by gibberellins. *Plant Cell* 2004; 16:2641-51.

- 152 16. Ishida S, Yuasa T, Nakata M, Takahashi Y. A tobacco calcium-dependent protein kinase,
153 CDPK1, regulates the transcription factor REPRESSION OF SHOOT GROWTH in
154 response to gibberellins. *Plant Cell* 2008; 20:3273-88.
- 155 17. Nakagawa Y, Katagiri T, Shinozaki K, Qi Z, Tatsumi H, Furuichi T, Kishigami A, Sokabe
156 M, Kojima I, Sato S, Kato T, Tabata S, Iida K, Terashima A, Nakano M, Ikeda M,
157 Yamanaka T, Iida H. Arabidopsis plasma membrane protein crucial for Ca^{2+} influx and
158 touch sensing in roots. *Proc Natl Acad Sci USA* 2007; 104:3639-44.
- 159 18. Chen Z, Noir S, Kwaaitaal M, Hartmann HA, Wu MJ, Mudgil Y, Sukumar P, Muday G,
160 Panstruga R, Jones AM. Two seven-transmembrane domain MILDEW RESISTANCE
161 LOCUS O proteins cofunction in Arabidopsis root thigmomorphogenesis. *Plant Cell*
162 2009; 21:1972-91.
- 163 19. Yamanaka T, Nakagawa Y, Mori K, Nakano M, Imamura T, Kataoka H, Terashima A,
164 Iida K, Kojima I, Katagiri T, Shinozaki K, Iida H. MCA1 and MCA2 that mediate Ca^{2+}
165 uptake have distinct and overlapping roles in Arabidopsis. *Plant Physiol* 2010; 152:1284-96.
- 166 20. Shih HW, Miller ND, Dai C, Spalding EP, Monshausen GB. The receptor-like kinase
167 FERONIA is required for mechanical signal transduction in Arabidopsis seedlings. *Curr Biol*
168 2014; 24:1887-92.
- 169 21. Torres-Schumann S, Ringli C, Heierli D, Amrhein N, Keller B. In vitro binding of the
170 tomato bZIP transcriptional activator VSF-1 to a regulatory element that controls

171 xylem-specific gene expression. Plant J 1996; 9:283-96.

172 22. Yin Y, Zhu Q, Dai S, Lamb C, Beachy RN. RF2a, a bZIP transcriptional activator of the
173 phloem-specific rice tungro bacilliform virus promoter, functions in vascular development.
174 EMBO J 1997; 16:5247-59.

175 23. Ringli C, Keller B. Specific interaction of the tomato bZIP transcription factor VSF-1
176 with a non-palindromic DNA sequence that controls vascular gene expression. Plant Mol
177 Biol 1998; 37:977-88.

178 24. Fukazawa J, Sakai T, Ishida S, Yamaguchi I, Kamiya Y, Takahashi Y. Repression of shoot
179 growth, a bZIP transcriptional activator, regulates cell elongation by controlling the level
180 of gibberellins. Plant Cell 2000; 12:901-15.

181 25. Dai S, Zhang Z, Chen S, Beachy RN. RF2b, a rice bZIP transcription activator, interacts
182 with RF2a and is involved in symptom development of rice tungro disease. Proc Natl
183 Acad Sci USA 2004; 101:687-92.

184 26. Dai S, Wei X, Alfonso AA, Pei L, Duque UG, Zhang Z, Babb GM, Beachy RN. Transgenic
185 rice plants that overexpress transcription factors RF2a and RF2b are tolerant to rice
186 tungro virus replication and disease. Proc Natl Acad Sci USA 2008; 105:21012-6.

187 27. Van Oosten MJ, Sharkhuu A, Batelli G, Bressan RA, Maggio A. The *Arabidopsis thaliana*
188 mutant *air1* implicates SOS3 in the regulation of anthocyanins under salt stress. Plant Mol
189 Biol 2013; 83:405-15.

28. Farage-Barhom S, Burd S, Sonogo L, Perl-Treves R, Lers A. Expression analysis of the *BFN1* nuclease gene promoter during senescence, abscission, and programmed cell death-related processes. *J Exp Bot* 2008; 59:3247-58.
29. Fendrych M, Van Hautegeem T, Van Durme M, Olvera-Carrillo Y, Huysmans M, Karimi M, Lippens S, Guérin CJ, Krebs M, Schumacher K, Nowack MK. Programmed cell death controlled by ANAC033/SOMBRERO determines root cap organ size in Arabidopsis. *Curr Biol* 2014; 24:931-40.


Figure Legend

Figure 1.

Touch-induced nuclear localization of VIP1.

Transgenic Arabidopsis plants expressing the VIP1-GFP fusion protein were grown for 3 days in darkness. VIP1-GFP signals were observed approximately 3 minutes after their hypocotyls were touched by a needle tip. More than five plants were used, and a representative result is presented. Arrows indicate the approximate position where the hypocotyl was touched.

Scale bar = 200 μ m.



217

218

219