HOKKAIDO UNIVERSITY

Title	Understanding the emergence of collective motion of microtubules driven by kinesins: role of concentration of microtubules and depletion force
Author(s)	Saito, Ai; Farhana, Tamanna Ishrat; Kabir, Arif Md. Rashedul et al.
Citation	RSC advances, 7(22), 13191-13197 https://doi.org/10.1039/c6ra27449h
Issue Date	2017-02-27
Doc URL	https://hdl.handle.net/2115/65443
Rights(URL)	https://creativecommons.org/licenses/by/3.0/
Type	journal article
File Information	Supplementary Information.pdf

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

Electronic Supplementary Material (ESI) for RSC Advances.
This journal is © The Royal Society of Chemistry 2017

Supporting Information

Understanding the emergence of collective motion of microtubules driven by kinesins: role of concentration of microtubules and depletion force

Ai Saito[1,], Tamanna Ishrat Farhana[1,*], Arif Md. Rashedul Kabir[2], Daisuke Inoue[2], Akihiko Konagaya[3], Kazuki Sada[1,2], Akira Kakugo[1,2,#]*

[1]Graduate School of Chemical Sciences and Engineering, Hokkaido University, Sapporo 060-0810, Hokkaido, Japan.

[2]Faculty of Science, Hokkaido University, Sapporo 060-0810, Hokkaido, Japan.

[3]School of Computing, Tokyo Institute of Technology, Yokohama, 226-8501, Japan.

#Corresponding author

E-mail: kakugo@sci.hokudai.ac.jp

Phone/FAX: +81-11-706-3474

[*]These authors contributed equally to this work.



Supporting Figure 1: (a) Schematic representation of analysis of microtubule orientation angle *Θ*. Yellow arrow represents moving direction of microtubules. As shown here, the orientation angle was measured with respect to the X-axis. Scale bar: 5 μm. (b) Circular histogram of orientation angle of of microtubules in absence of methylcellulose at (i) 0 min and (ii) 120 min after ATP addition. The dispersion of orientation angle in circular histogram indicates randomness in moving direction of MT. Here the concentration of microtubules was 15.0 μM



Supporting Figure 2: Relationship between depletion force and concentration of methylcellulose.



Supporting Figure 3: Effect of methylcellulose concentration on the gliding behavior of microtubules. High concentration of methylcellulose increased the probability of snuggling event.



Supporting Figure 4: Effect of methylcellulose concentration on the velocity of gliding microtubules.



Supporting Figure 5: Relationship between buffer viscosity and concentration of methylcellulose.

Caption for the Supporting movie:

Supporting movie 1: Local streams generated by collectively moving microtubules in the presence of methycellulose. Here the concentration of microtubules and methylcellulose were 10 μM and 0.1 wt% respectively.