



Title	Higenamine 4'-O- β -D-glucoside in the lotus plumule induces glucose uptake of L6 cells through β 2-adrenergic receptor
Author(s)	Kato, Eisuke; Inagaki, Yosuke; Kawabata, Jun
Citation	Bioorganic & Medicinal Chemistry, 23(13), 3317-3321 https://doi.org/10.1016/j.bmc.2015.04.054
Issue Date	2015-07-01
Doc URL	https://hdl.handle.net/2115/66355
Rights	©2015. This manuscript version is made available under the CC-BY-NC-ND 4.0 license
Rights(URL)	https://creativecommons.org/licenses/by-nc-nd/4.0/
Type	journal article
File Information	BMC2015_23_3317-3321_Sp_HUSCAP.pdf



Supplementary material

Higenamine 4'-*O*- β -D-glucoside in the lotus plumule induces glucose uptake of L6 cells through β 2-adrenergic receptor

Eisuke Kato ^{a,*}, Yosuke Inagaki ^b, and Jun Kawabata ^a

^a Laboratory of Food Biochemistry, Division of Applied Bioscience, Graduate School of Agriculture, Hokkaido University, Kita-ku, Sapporo, Hokkaido 060-8589, Japan

^b Nippon Supplement, Inc., Applause Tower 18F, 19-19, Chayamachi, Kita-ku, Osaka 530-0013, Japan


*Corresponding author:

Tel/Fax: +81-11-706-2496


E-mail address: eikato@chem.agr.hokudai.ac.jp

This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Acquisition Time (sec)	3.0331	Date	05 Aug 2013 09:13:50	Date Stamp	Mon Aug 05 09:13:09 2013
File Name	D:\NMR\2013-31-2.xls	Frequency (MHz)	270.05	Nucleus	¹ H
Original Points Count	16384	Points Count	16384	Pulse Sequence	NOE
Spectrum Offset (Hz)	1498.9984	Sweep Width (Hz)	5401.76	Temperature (degree C)	26.000
				Receiver Gain	23.00
				Number of Transients	128
				Solvent	METHANOL-d ₄



Supplementary Fig 1. ¹H-NMR spectra of isolated **1** in CD₃OD (270 MHz).



Supplementary Fig 2. ¹H-NMR spectra of isolated (black) and synthetic (red) **1** in D₂O (500 MHz).